



Teck Coal Environmental Office
Bag Service 2000, 421 Pine Avenue
Sparwood, B.C. Canada V0B 2G0

+1 250 425 3331 Tel
www.teck.com

Technical Report Overview

Report: Data Report for the Tributary Evaluation Program (DRAFT)

Overview: This report provides an inventory of all tributaries to the Elk and Fording Rivers that are located within management units (MUs) 1, 2, 3 and 4 (as defined by the Elk Valley Water Quality Plan). This report evaluates the ecological value of these tributaries to the Elk and Fording Rivers and helps to identify tributaries that play a significant role in supporting the health of the ecosystem as a whole.

This report was prepared for Teck by Minnow Environmental Inc.

For More Information

If you have questions regarding this report, please:

- Phone toll-free to 1.855.806.6854
- Email feedbackteckcoal@teck.com

Future studies will be made available at teck.com/elkvalley



**Data Report for the
Tributary Evaluation Program
(DRAFT)**

Prepared For:
Teck Coal Limited
Sparwood, BC

Prepared By:
Minnow Environmental Inc.
Georgetown, ON

June 2016

TABLE OF CONTENTS

1.0	INTRODUCTION	1
1.1	Background	1
1.2	Tributary Evaluation Program Requirements	1
1.3	Tributary Management Plan Requirements.....	3
1.4	Phased TEP Study Design.....	4
1.5	Environmental Monitoring Committee Input	4
2.0	APPROACH AND METHODS	7
2.1	Review of Existing Information	11
2.2	Data Management.....	11
2.3	Watershed Area and Catchment Disturbances	11
2.3.1	Catchment Area	11
2.3.2	Teck Ownership Rights.....	13
2.3.3	Mine-Related and Other Disturbances in Catchment	13
2.3.4	Additional Planned Mining Disturbances and Mitigation/Offsetting.....	14
2.4	Environmental Quality Descriptors	14
2.4.1	Water Quality Monitoring Station	16
2.4.2	Water Quality Index	16
2.4.3	Concentrations of Selenium, Nitrate, Sulphate, Cadmium.....	17
2.4.4	Median Concentrations, Trends, and Loads	19
2.4.5	Surface Water Acute Toxicity.....	21
2.4.6	Sediment Quality.....	21
2.4.7	Calcite	22
2.4.8	Benthic Invertebrate Community Structure	23
2.4.9	Selenium in Periphyton or Invertebrate Tissue	24
2.5	Fish and Aquatic Habitat Information	25
2.5.1	Annual Average Flow.....	25
2.5.2	Ktunaxa Traditional Uses and Cultural Values	25
2.5.3	Stream Length	25
2.5.4	Fish Species Present.....	27
2.5.5	Unique Fish Habitat Features	27
2.5.6	Anthropogenic and Natural Barriers Present	27
2.5.7	Fish Habitat Lengths	27
2.5.8	Riparian Habitat	28
2.5.9	Amphibians and Birds	30
2.5.10	Written Habitat Descriptions	31
2.6	Orthophoto Figures	33
3.0	TRIBUTARIES TO UPPER FORDING RIVER (MU1).....	35
3.1	Upstream Fording River	40
3.2	Henretta Creek	41
3.3	Fish Pond Creek.....	43
3.4	Clode Creek	44
3.5	Lake Mountain Creek	46
3.6	Kilmarnock Creek	49
3.7	Swift Creek	50
3.8	Cataract Creek	51
3.9	Porter Creek.....	53
3.10	Chauncey Creek.....	53

3.11	Ewin Creek	54
3.12	LCO Dry Creek	55
3.13	Greenhills Creek	57
4.0	TRIBUTARIES TO LOWER FORDING RIVER (MU2)	60
4.1	Grace Creek	60
4.2	Line Creek	65
5.0	TRIBUTARIES TO UPPER ELK RIVER (MU3)	69
5.1	Branch F Creek	69
5.2	Wolf Creek	74
5.3	Willow Creek	75
5.4	Wade Creek	77
5.5	Cougar Creek	78
5.6	No-Name Creek and Unnamed Tributary 14	79
5.7	Mickelson Creek	80
5.8	Leask Creek	81
5.9	Wolfram Creek	84
5.10	Thompson Creek	86
6.0	TRIBUTARIES TO MICHEL CREEK (MU4)	88
6.1	Upstream Michel Creek	93
6.2	Corbin Creek	93
6.3	Andy Good Creek	94
6.4	Leach Creek	95
6.5	Carbon Creek	96
6.6	Snowslide Creek	98
6.7	Wheeler Creek	100
6.8	Unnamed Tributary to Michel Creek	101
6.9	Fir Creek	104
6.10	Alexander Creek	104
6.11	Erickson Creek	106
6.12	South Pit Creek	108
6.13	Milligan Creek	109
6.14	Gate Creek	109
6.15	Bodie Creek	110
6.16	Aqueduct Creek	112
7.0	TRIBUTARIES TO MIDDLE ELK RIVER (MU4/MU5)	116
7.1	Grave Creek (including Harmer Creek and EVO Dry Creek)	116
7.2	Six-Mile Creek	124
7.3	Balmer Creek	126
7.4	Fennelon Creek	127
7.5	Goddard Creek	128
7.6	Otto Creek	131
7.7	Transmission Creek	133
7.8	Mine Creek (Hosmer Creek)	134
8.0	REFERENCES	137

LIST OF FIGURES

Figure 1.1:	Location of Teck Coal Operations.....	2
Figure 1.2:	Flow chart for developing and updating the Tributary Evaluation Program.	5
Figure 2.1:	Tributaries of Upper Fording and Upper Elk Rivers (MU1 to MU3).....	9
Figure 2.2:	Tributaries of Michel Creek and Middle Elk River (MU4, MU5).....	10
Figure 3.1:	Draft Upper Fording River Watershed Boundary.	36
Figure 3.2:	Selenium concentration trend over time (2004-2013) in Henretta Creek at the confluence with the Fording River (FR_HC1).....	43
Figure 3.3:	Selenium concentration trend over time (2004-2013) in Greenhills Creek at the confluence with the Fording River (GH_GH1).....	58
Figure 4.1:	Draft Lower Fording River Watershed Boundary.	61
Figure 5.1:	Draft Upper Elk River Watershed Boundary.	70
Figure 6.1:	Draft Michel Creek Watershed Boundary.	89
Figure 6.2:	Selenium concentration trend over time (2004-2013) in Corbin Creek at the confluence with Michel Creek (CM_CC1).....	94
Figure 7.1:	Draft Middle Elk River Watershed Boundary.....	117
Figure 7.2:	Selenium concentration trend over time (2004-2013) at the Harmer Pond outlet into Gate Creek (EV_HC1).	121
Figure 7.3:	Selenium concentration trend over time (2004-2013) in Goddard Creek at the confluence with the Elk River (EV_GC2).....	129

LIST OF PHOTOS

Photo 3.1	Fording River upstream of Henretta Creek (Golder 2014a).....	41
Photo 3.2:	Henretta Lake (Wright et al. 2005).....	42
Photo 3.3:	Clode Creek downstream of Clode Ponds (Golder 2014a).....	45
Photo 3.4:	Culvert at Clode Pond access road in the lower section of Lake Mountain Creek (Golder 2014a).	48
Photo 3.5:	Swift Creek at the Fording River post-flood 2013 (Golder 2014a).	52
Photo 3.6:	Paired culvert crossing under Fording Road in Dry Creek (Teck 2011b).....	56
Photo 4.1:	Typical riffle-glide morphology in Grace Creek (Teck 2011b).....	65
Photo 4.2:	Downstream end of the rock drain in Line Creek (Minnow internal sources).	66
Photo 6.1:	Typical riffle habitat in Leach Creek (Golder 2015).....	96
Photo 6.2:	Carbon Creek cascade barrier (Golder 2015).....	98

Photo 6.3:	Bedrock falls on Snowslide Creek (Golder 2015).	99
Photo 6.4:	Example of steep-gradient cascade barrier in Wheeler Creek (Golder 2015).	101
Photo 6.5:	First rock fall barrier to fish movement on Unnamed Michel Creek Tributary (Golder 2015).	103
Photo 6.6:	Second rock fall barrier to fish movement on Unnamed Michel Creek Tributary (Golder 2015).	103
Photo 6.7:	Gradient barrier defining upstream fish distribution limit in Fir Creek (Golder 2015).	105
Photo 6.8:	Most downstream pond in Erickson Creek headwaters (Lotic 2015a).	107
Photo 6.9:	Moss and calcified streambed in Erickson Creek (Lotic 2015a).	108
Photo 6.10:	Downstream view of the concrete culvert and flap valve for Gate Creek (Lotic 2015a).	111
Photo 6.11:	CPR double culverts in Bodie Creek (Lotic 2015a).	113
Photo 6.12:	Wetland habitat in lower Aqueduct Creek (Lotic 2015a).	114
Photo 7.1:	Upstream view of the 3 m high waterfall below the Grave Lake confluence in Grave Creek (Lotic 2015a).	122
Photo 7.2:	Harmer Pond 0.5 km upstream of the confluence of Harmer Creek with Grave Creek (Lotic 2015a).	123
Photo 7.3:	Channel habitat in lower Six-Mile Creek (Lotic 2015a).	126
Photo 7.4:	View of 2 m tall waterfall at the confluence of Fennelon Creek with the Elk River (Lotic 2015a).	128
Photo 7.5:	Goddard Marsh at the confluence of Goddard Creek and the Elk River (Lotic 2015a).	130
Photo 7.6:	Channelized section of Otto Creek between the rail line and the tailings lagoons (Lotic 2015a).	131
Photo 7.7:	Typical large woody debris/sediment wedge acting as a temporary barrier on Transmission Creek (Golder 2015).	133
Photo 7.8:	Lentic area in Mine Creek, upstream of the confluence with Hosmer Creek (Golder 2015).	135

LIST OF TABLES

Table 2.1	Relevant tributaries according to management unit.	8
Table 2.2:	Teck's in-house information sources related to watershed area and disturbances within catchment.	12
Table 2.3:	Information sources for environment quality data.	15
Table 2.4:	Ranking criteria for observed environmental conditions.	18

Table 2.5:	Level 1 (10%) effect benchmarks identified in the Elk Valley Water Quality Plan.....	19
Table 2.6:	Sources of aquatic habitat information.....	26
Table 2.7:	Tributary features depicted in tributary orthophotos.....	34
Table 3.1:	Watershed area and disturbances for the Upper Fording River (MU1).....	37
Table 3.2:	Environmental quality and descriptors for Upper Fording River (MU1).....	38
Table 3.3:	Aquatic habitat of Upper Fording River (MU1).....	39
Table 4.1:	Watershed area and disturbances for the Lower Fording River (MU2).....	62
Table 4.2:	Environmental quality descriptors for Lower Fording River (MU2).....	63
Table 4.3:	Aquatic habitat of Lower Fording River (MU2).....	64
Table 5.1:	Watershed area and descriptors for tributaries to Upper Elk River (MU3).....	71
Table 5.2:	Environmental quality descriptors for Upper Elk River (MU3).....	72
Table 5.3:	Aquatic habitat of Upper Elk River (MU3).....	73
Table 6.1:	Watershed area and disturbances for tributaries to Michel Creek (MU4).....	90
Table 6.2:	Environmental quality descriptors for tributaries to Michel Creek (MU4).....	91
Table 6.3:	Aquatic habitat of Michel Creek tributaries (MU4).....	92
Table 7.1:	Watershed area and disturbances for Middle Elk River (MU4 and MU5).....	118
Table 7.2:	Environmental quality descriptors for Middle Elk River (MU4 and MU5).....	119
Table 7.3:	Aquatic habitat of Middle Elk River (MU4 and MU5).....	120

APPENDIX A TRIBUTARIES TO UPPER FORDING RIVER (MU1)

APPENDIX B TRIBUTARIES TO LOWER FORDING RIVER (MU2)

APPENDIX C TRIBUTARIES TO UPPER ELK RIVER (MU3)

APPENDIX D TRIBUTARIES TO MICHEL CREEK (MU4)

APPENDIX E TRIBUTARIES TO MIDDLE ELK RIVER (MU4/MU5)

APPENDIX F BIBLIOGRAPHIES

1.0 INTRODUCTION


1.1 Background

Teck Resources Limited (Teck) operates five, open pit, steelmaking coal mines in the Elk River watershed, which are the Fording River Operations (FRO), Greenhills Operations (GHO), Line Creek Operations (LCO), Elkview Operations (EVO), and Coal Mountain Operations (CMO; Figure 1.1). Discharges from the mines to the Elk River watershed are authorized by the British Columbia Ministry of Environment (BCMOE) through permits that are periodically issued under provisions of the *Environmental Management Act*. Permit 107517, issued November 14, 2015, and amended December 23, 2015, specifies the terms and conditions associated with discharges from Teck's five Elk Valley coal mine operations. Section 5 of the Permit requires Teck to develop a Tributary Evaluation Program (TEP) and Tributary Management Plan (TMP), which are described in more detail below.

1.2 Tributary Evaluation Program Requirements

According to Permit 107517, the TEP "is intended to evaluate the ecological value of tributaries to the Elk and Fording Rivers to support identification of tributaries that play a significant role in supporting the health of the ecosystem as a whole." Section 5 of Permit 107517 specifies that the TEP must include the following elements:

1. Inventory of all tributaries to the Elk and Fording Rivers that are located in Management Units (MUs) 1, 2, 3, and 4 (as defined by the Elk Valley Water Quality Plan [EVWQP]) that are affected or potentially influenced by Teck's current and future development plans;
2. Maps of MUs 2, 3, and 4 showing the locations of the tributaries of the Elk and Fording Rivers, and identifying the tributaries that are affected or potentially influenced by Teck's current and future development plans;
3. Collation of existing and readily available data and information on each tributary, including surface-water chemistry, surface-water toxicity, sediment chemistry, sediment-toxicity, calcification, flow, habitat value ranking, benthic invertebrate community structure, and habitat use by fish and/or sensitive aquatic dependent wildlife (i.e., water birds);



<div data-bbox="350 2613 419 2682"></div> <div data-bbox="264 2707 506 2775"></div> <div data-bbox="197 2800 552 2946"><p>MAP INFORMATION Map Projection: UTM Zone 11 NAD 1983 Data Source: Reproduced under licence from Her Majesty the Queen in Right of Canada, Department of Natural Resources Canada, GeoBase and BCGov, GeoBC. All rights reserved. Creation Date: March 2016 Project No.: 2563</p></div>	<div data-bbox="600 2613 903 2862"><ul style="list-style-type: none">Teck Coal Mine OperationsWaterbodyWetlandWatercourseInternational BoundaryProvincial / State BoundaryHighwayTownWater Flow Direction</div>		<div data-bbox="1439 2613 1786 2775"><p>Figure 1.1: Teck Coal Limited Operations within the Elk River Valley, Southeast British Columbia.</p></div> <div data-bbox="1487 2862 1733 2946"></div>
--	---	--	--

4. Evaluation of historical (i.e. conditions relevant to the 1980 timeframe, where available) and current habitat value, based on surface-water quality, sediment quality, extent of calcification, flow, amount of habitat available, habitat types, physical features, connectivity to fish habitat, status of riparian habitat, and habitat use by fish and sensitive aquatic dependent wildlife species;
5. Evaluation of the potential for rehabilitation of aquatic and riparian habitat and potential for improvement of water quality conditions; and,
6. Prioritization of each tributary for ongoing protection and/or restoration based on the evaluation of current ecological value, potential for rehabilitation, and potential to contribute to the objectives of the EVWQP.

Elements #1 through #3 are addressed in this report, which was first submitted to BCMOE on March 31, 2016, and was updated subsequently, as reflected in this version. Elements #4 through #6 will be completed and documented in a written report to be submitted to the EMC by August 31, 2016. The purpose of the TEP, as stated in the permit, is “to provide context for the development of specific management objectives for tributaries included in the Tributary Management Plan”. Thus, the TEP will result in the identification of tributaries that should be targeted for protection from future mine-related degradation, as well as the identification of mine-influenced tributaries that should be targeted for rehabilitation.

1.3 Tributary Management Plan Requirements

Permit 107517 requires Teck to develop and implement a Tributary Management Plan (TMP), after the evaluation of the tributaries in the TEP. The TMP is “intended to incorporate protection and rehabilitation goals¹ for tributaries that will support achieving the area-based objectives of the Elk Valley Water Quality Plan.” A terms of reference and interim TMP report will be submitted to the EMC by March 31, 2016 and October 31, 2016, respectively. The TMP must be submitted to the Director of BCMOE for acceptance by December 31, 2016.

¹ Permit 107517 provides the following clarification related to rehabilitation of historically impacted areas: “The scope of the Tributary Management Plan excludes tributaries that have been permanently removed or severely altered (e.g., covered by waste spoils or other mine infrastructure or dewatered) by mining activities within the Permittee’s current mine permit boundaries. Loss of habitat for such tributaries is governed by requirements under the Federal *Fisheries Act* and the provincial mitigation policy.”

1.4 Phased TEP Study Design

The TEP study design submitted to BCMOE May 29th, 2015, described a phased approach for completing the TEP (Minnow 2015) and fulfilling the Permit associated Permit requirements. It will summarize existing information/data and identify data gaps to support an integrated evaluation for each tributary in the context of protection and rehabilitation goals that will be defined as part of the process (Figure 1.2).

The first phase of the approach (Phase A in Figure 1.2) addresses the TEP requirements #1 and #2 by preparing a tabular inventory and maps of tributaries that are or may be influenced by mining. The next phase (Phase B in Figure 1.2) addresses TEP requirement #3 by describing the ecological characteristics and future mine development plans, to the extent such information is readily available in existing documents, or from Teck's in-house sources, including a summary of current and historical conditions (e.g., water and sediment quality and toxicity, calcite levels, benthic invertebrate community composition, and documented use by fish and other vertebrate species), habitat characteristics, expected future mine-related disturbances (locations and types), and planned mitigation actions for each tributary in each watershed. The information corresponding to Phases A and B of the TEP is presented in this data report. As noted in the TEP study design, subsequent phases will be completed as part of the TEP evaluation report to be submitted to the EMC by August 31, 2016.

1.5 Environmental Monitoring Committee Input

Section 12.2 of the Permit also requires Teck to consider input from the EMC. The EMC consists of representatives from the BCMOE, the Ministry of Energy and Mines, Environment Canada, the Ktunaxa Nation Council (KNC), Interior Health Authority, an Independent Scientist, and Teck. The TEP and TMP include fish and fish habitat components that will inform Teck's Regional Fish Habitat Management Plan, which is currently being developed with the Elk Valley Fish and Fish Habitat Committee (EVFFHC). It is envisioned that the TEP and TMP will eventually be integrated with the Regional Fish Habitat Management Plan that will inform Teck's submissions under Section 35 of the federal *Fisheries Act*.

The EVFFHC includes representatives from Fisheries and Oceans Canada (DFO), the Ministry of Forests, Lands, and Natural Resource Operations (FLNRO), KNC, and Teck, resulting in overlap between the EVFFHC and EMC in terms of both membership and information needs. Therefore, the EVFFHC is participating in EMC meetings during discussions related to the TEP and TMP and providing input regarding deliverables. Input




Figure 1.2: Flow chart depicting phases for developing and updating the Tributary Evaluation Program.

received to date from both the EMC and EVFFHC has been considered in this data report.

The following key items have been addressed in response to input from the EMC regarding the first draft TEP report submitted March 31, 2016:

- Columns for the road density dataset from the Elk Valley Cumulative Effects Management Framework (CEMF) have been added and populated with data received from the Province.
- A column for Ktunaxa traditional uses and cultural values has been added and will be populated when information is received from KNC.
- A column for riparian habitat has been added using the riparian analysis completed for the Baldy Ridge Extension Project.
- The catchment areas and stream lengths presented in the tables and on orthophoto figures have also been updated based on the provincial Corporate Watershed Base dataset, which uses the Fresh Water Atlas (FWA) (TRIM1) data.
- Columns with fish habitat metrics have been added following methods similar to the BCMOE Fish Passage GIS Analysis Methodology and Output Data Specifications Version 2 (Norris and Mount in draft). Version 2 builds on Version 1 of the BC MOE GIS Modelling of Fish Habitat and Road Crossings for the Prioritization of Culvert Assessment and Remediation Version 1 (Mount et al. 2011).

2.0 APPROACH AND METHODS

Information for tributaries is provided in Sections 3.0 to 7.0 of this report and organized according to the Management Unit (MU) in which the tributaries are located. The MUs included in this report are as follows: Upper Fording River Watershed (MU1); Lower Fording River Watershed (MU2); Upper Elk River Watershed (MU3); Michel Creek Watershed (MU4); and Middle Elk River Watershed (MU4/MU5²). The tributaries discussed in this report include all streams within these MUs that flow into the Fording River, Michel Creek, or Elk River that are currently mine-influenced (i.e., the mine footprint extends into a portion of the tributary's catchment) and streams that have not been historically influenced by mining but for which Teck has ownership rights within the catchment area (see Section 2.2.4). All tributaries considered in recent Environmental Assessments (e.g., LCO Phase II, Fording-Swift, Baldy Ridge Extension, Cougar Pit Extension and CMO Phase 2) are also included. The tributaries included in each MU are listed in Table 2.1 and shown on Figures 2.1 and 2.2.


Three data matrices are presented for each group of tributaries (by MU), which correspond to three categories of information: (1) total watershed area and mine-related disturbances within the watershed; (2) environmental quality data (e.g., water, sediment, calcite, and biological data); and (3) fish and aquatic habitat data. The methods and sources associated with this tabulated information are described below. A text description of each tributary is also presented for each tributary, which generally emphasizes information that was available, but not amenable to a comparative data matrix format (e.g., description of mining history, results of habitat surveys). Graphs and photographs are also presented to illustrate selected data or habitat characteristics mentioned in tables or text.

General overviews of information sources and data management practices are presented below in Sections 2.1 and 2.2. The methods and sources of information used to populate the three types of data matrices (listed above) for each MU are described in Sections 2.3 to 2.6 in an order that corresponds to the format of each table.

² The only MU5 tributaries included in the TEP are those that were identified as part of the CMO Phase 2 baseline study (Golder 2015a). The CMO Phase 2 project is no longer proceeding.

Table 2.1 Relevant tributaries according to management unit.

Management Unit (MU)		Tributary Name
MU1	Upper Fording River	Upstream Fording River (above Henretta Creek confluence)
		Henretta Creek
		Fish Pond Creek
		Clode Creek
		Lake Mountain Creek
		Kilmarnock Creek
		Swift Creek
		Cataract Creek
		Porter Creek
		Chauncey Creek
		Ewin Creek
		LCO Dry Creek
		Greenhills Creek
MU2	Lower Fording River	Grace Creek
		Line Creek
MU3	Upper Elk River	Branch F Creek
		Wolf Creek
		Willow Creek (Including Willow South, Willow North and Wilde)
		Wade Creek
		Cougar Creek
		No-Name Creek
		Mickelson Creek
		Leask Creek
		Wolfram Creek
		Thompson Creek
MU4	Michel Creek	Upstream Michel Creek (above Corbin Creek confluence)
		Corbin Creek
		Andy Good Creek
		Leach Creek
		Carbon Creek
		Snowslide Creek
		Wheeler Creek
		Unnamed to Michel Creek
		Fir Creek
		Alexander Creek
		Erickson Creek
		South Pit Creek
		Milligan Creek
		Gate Creek
		Bodie Creek
		Aqueduct Creek
MU4/MU5	Middle Elk River	Grave Creek (including Harmer and EVO Dry creeks)
		Six-mile Creek
		Balmer Creek
		Fennelon Creek
		Goddard Creek
		Otto Creek
		Transmission Creek
		Hosmer Creek (including Mine Creek)



MAP INFORMATION

Map Projection: UTM Zone 11N NAD 1983

Data Source: Reproduced under licence from Her Majesty the Queen in Right of Canada, Department of Natural Resources Canada. All rights reserved.

Creation Date: March 2016

Project No.: 2563

Legend

- Water Flow Direction
- City
- Waterbody
- Teck Coal Mine Operation
- Provincial Boundary

Management Units (MU)

- MU-1
- MU-2
- MU-3
- MU-4
- MU-5
- MU-6

Figure 2.1: Tributaries of Upper Fording and Upper Elk Rivers (MU1 to MU3)




Figure 2.2: Tibutaries of Michel Creek and Middle Elk River (MU4, MU5)

2.1 Review of Existing Information

Information about tributaries in the Elk River watershed was gathered from a variety of sources and reviewed. The information reviewed included scientific literature, consultant reports prepared for Teck and other organizations, government agency reports, and electronic databases maintained by federal and provincial government agencies. Most of the information was derived from sources internal to Teck, whereas the remainder was summarized from recent and historical technical reports. An annotated bibliography of technical reports with tributary information is provided in Appendix F of this report.

2.2 Data Management

Data for the TEP were generated from Teck's ArcGIS and EQUIS databases, with output provided in Excel spreadsheets.

2.3 Watershed Area and Catchment Disturbances

Strictly speaking, "watershed" refers to the high ground or ridge that divides separates waters flowing to adjacent river systems, whereas terms such as "catchment", "basin", and "drainage area" refer to the area bounded by the watershed. However, these terms are often used interchangeably, as is usually the case when referring to the Elk River watershed. Likewise, this document uses the word "watershed" interchangeably with those referring to catchment area, so the term "watershed boundary" has been used to refer to the line dividing adjacent catchments.

The first data matrix presented for each MU identifies the watershed area for each tributary within the MU, the mine-related disturbance footprint (including a list of the types of mine-related disturbances within the tributary catchment), and the area affected by forestry, and roads. The methods and sources used to gather this information are described below. Table 2.2 provides a brief summary of the same information for quick reference.

2.3.1 Catchment Area

The catchment area for each tributary was determined using the Corporate Watershed Base (CWB) dataset provided by the Province. The CWB dataset, formerly known as the TRIM Watershed Atlas, is a watershed atlas that defines watersheds and provides an associated stream and lake network. The CWB dataset adds functionality to TRIM 1:20,000 digital topographic base map data by providing a connected feature-coded stream network, hydrographic information, and associated watershed boundaries. The CWB is derived from TRIM1 mapping. The total catchment area (km²) from the Province's

Table 2.2: Teck's in-house information sources related to watershed area and disturbances within catchment.

Information Type		Source	Definition
Tributary Name		Provincial CWB dataset/Common Names	A stream that flows into the Fording River, Michel Creek, or Elk River that is currently mine influenced (the mine footprint extends into a portion of the tributary's catchment) or Teck has ownership rights within the tributary catchment area.
Total Catchment Area from the Provincial CWB dataset (km2) (1980s plus updates)		Provincial CWB dataset	The total catchment area of the tributary of interest based on the Provincial CWB dataset, which reflects predominantly 1980s conditions and potentially subsequent (unknown) updates.
Total Catchment Area as of September 2015		July/August 2014 LiDAR (scale 1:500) and internal knowledge	The watershed boundary of the tributary of interest, considering current topography if affected by mining as of September 2015.
% Change in Catchment Area		Calculated	= (Catchment area from Provincial CWB dataset minus Sept. 2015 catchment area) / Catchment area from Provincial CWB dataset * 100
Teck Ownership Rights: % of Catchment Area and Type (Coal, Surface, Timber)		Provincial CWB dataset, Provincial parcel fabric layer and Teck's parcel fabric layer	Type is defined as follows: "Coal" indicates crown land in which Teck has coal leases/licences; "Coal and Surface" indicates private land without the timber rights; "Coal, Surface and Timber" indicates private land with timber rights; "Surface and Timber" indicates private land without the coal rights
Mine-Related Disturbances in Area from Provincial CWB dataset (1980s plus updates)	Timing of Mine-Related Disturbances [Historical (H), Current (C), Near Future (NF), Far Future (FF) or Not Expected (N) disturbances]	Provincial CWB dataset, Current disturbance: Teck's dissolved disturbance layer (Scale 1:20,000) Future development: Teck's Development layer (Scale 1:20,000)	FRO: Historical 1971-2012, Current active 2013-2016, Near Future 2017-2021, Far future 2022-2072. GHO: Historical 1978-2012, Current active 2013-2016, Near Future 2017-2021, Far future 2022-2070. LCO Phase 2: Current active 2014-2016, Near Future 2017-2021, Far Future 2022-2057. LCO: Historical 1980-2012, Current active 2013-2016, Near Future 2017 - 2021, Far Future 2022-2032. EVO: Historical 1969-2012; Current Active 2013-2016, Near Future 2017-2021, Far Future 2022-2045. CMO: Historical 1943-2012, Current Active 2013-2016, Near Future 2017
	Total footprint area currently disturbed (km, %)		Total disturbance area within the catchment that has not yet been reclaimed (reclaimed lands represents a small amount of the total disturbance footprint in most cases, as shown in Table 2.1 of the TEP Study Design).
	Reclaimed Land (km, %)	Provincial CWB dataset, Orthophoto Imagery/Teck's Reclaimed Land layer	Land that has been successfully revegetated for one yea
	Pits	Provincial CWB dataset, Teck's Development layer (1:20,000)	Area where mining has occurred, includes highwalls and footwalls. Indicates if pit(s) exist(s) within the catchment area of the tributary.
	Waste rock storage	Provincial CWB dataset, Teck's Development layer (1:20,000)	Dump or spoil, area where waste rock is stored. Indicates if waste rock exists within the catchment area of the tributary.
	Rock drains	Provincial CWB dataset, Teck's Water Management Lines layer (1:20,000)	Indicates if rock drains currently exist within the perennial wetted length of a tributary. Areas where a channel flows through waste rock and has a significant influence on the channel can be natural or man made.
	Channel re-alignment or diversions	Provincial CWB dataset, Teck's Water Management Lines layer (1:20,000)	Indicates if channels have been realigned or diverted within the catchment area of the tributary.
	Settling ponds/Catch basins	Provincial CWB dataset, Teck's Water Management layer (1:20,000)	Body of standing water due to man-made features; used to settle sediments; includes tailings ponds. Total count of ponds that exist within the catchment area of the tributary.
Forestry Disturbances ¹ in Catchment		Provincial CWB dataset, Elk Valley Cumulative Effects Management Framework Equivalent Clearcut Area layer	Area of forestry disturbances within catchment area of the tributary (includes data back to 1956).
Road Disturbances in Catchment	Road density per km ²	Provincial CWB dataset, Elk Valley Cumulative Effects Management Framework road density layers	Influences peak and low flow and water temperature, coarse and fine sediment delivery to stream and access for anglers.
	Stream crossings per km		Stream crossings (culverts, bridges) contribute to fine sediment in streams by exposing soils and acting as points of entry for sediment transport along ditches, as well as acts as barriers to fish and other aquatic organisms.
	Roads within 100 m of a stream		Responsible for the majority of fine sediment delivery that affects water quality, depending on soil texture, road construction and maintenance standards and precipitation.
	% roads on slopes > 60%		Roads on unstable, steep terrain increases the erosion potential through altered surface drainage, diversion of subsurface flow to the surface and reducing structural support of soil provided by roots and other vegetation.
Additional Planned Future Mining Disturbances		Provincial CWB dataset, Teck's Development layer (Scale assumed 1:20,000)	Explanation of known current, near future or far future mining disturbances.
Planned Mitigation/Fish Habitat Offsetting		Provincial CWB dataset, Teck Internal Sources; EVWQP; Fish Habitat Management Planning	Mitigation or offsetting related to water and fish management

CWB dataset is provided in the first matrix and is used for all calculations (i.e., ownership and mine-related, forestry, and road disturbances). The total catchment area from the Provincial CWB dataset is also delineated on the orthophoto figures of tributaries (Appendices A-E). The catchment area for each tributary is based on the Provincial CWB dataset and reflects predominantly 1980s conditions and potentially subsequent (unknown) updates by the province.

The total current catchment area (km² and % change from the Province's CWB catchment area) reflects the current watershed boundary of each tributary as of September 2015. The total current catchment area considers the current topography in areas that have been affected by mining. The total current catchment areas were defined using July/August 2014 LiDAR (scale 1:500) and manually updated, if appropriate, based on mining activities and internal knowledge on water management. The total current catchment area is also delineated on the orthophoto figures (Appendices A to E).

2.3.2 Teck Ownership Rights

Teck has varying types of ownership rights among the tributary catchment areas, which are as follows: "Coal" indicates crown land for which Teck has coal leases/licences; "Coal and Surface" indicates private land without the timber rights; "Coal, Surface and Timber" indicates private land with timber rights; "Surface and Timber" indicates private land without the coal rights. The reference datasets are the provincial parcel fabric layer provided by the BC government and Teck's parcel fabric layer, and were calculated based on the provincial CWB dataset.

2.3.3 Mine-Related and Other Disturbances in Catchment

The timing of mine-related disturbances in each catchment (i.e., historical, current, near future, far future) or, if not expected, were identified by appropriate Teck personnel and are current to the time of reporting. Table 2.2 identifies the years that define historical, current, near future, and far future for each operation.

The total footprint area of mine disturbance area (km² and %) within the CWB catchment area has been calculated from Teck's dissolved disturbance layer (current to time of reporting) and Provincial CWB catchment boundary reference dataset. The total mine footprint area excludes reclaimed lands, which are presented separately in the matrices. Reclaimed lands are defined as land that has been successfully revegetated for one year.

Teck's Development Layer reference dataset (1:20,000) provided information on pits, waste rock storage, rock drains, channel re-alignment/diversions and settling ponds/catch

basins at the time of reporting. These mine-related disturbances are defined in Table 2.2. The first matrix in each of Sections 3.0 to 7.0 of this report lists a “yes” or “no” to indicate if the disturbance corresponding to the column header is present within the provincial CWB catchment (or the number of such disturbances, such as settling ponds, where appropriate).

The forestry disturbance area (km²) in the CWB catchment area were taken from the Equivalent Clearcut Area (ECA) Data Layer which includes include any forestry disturbance within the catchment area of the tributary dating back to 1955 and current to 2015. The data were divided into two timeframes (1955 to 2010 and 2011 to 2015) to distinguish recent from longer-term disturbances. The ECA provincial dataset is from the Province’s Cumulative Environmental Monitoring Framework (CEMF) dataset and includes the area that has been harvested, cleared or burned.

The road disturbances (road density per km², stream crossings per km, roads within 100 m of a stream, % roads on slopes greater than 60%) are taken from the Province’s CEMF dataset and calculated based on the provincial CWB catchment area. Calculations did not include data (i.e., roads and the catchment area) within mine disturbance areas.

2.3.4 Additional Planned Mining Disturbances and Mitigation/Offsetting

The matrices in Sections 3.0 to 7.0 of this report identify any additional near and far future mining-related disturbances known at the time of reporting. The planned mitigation describes the water quality and quantity and fish management strategies that are expected to occur as a result of planned mine developments. This information is provided by Teck and includes information on future mining development as it impacts the affected tributaries. Management strategies include fish habitat offsetting, fish exclusion works, fish salvage activities, water pumping activities, water redirection and calcite management.

2.4 Environmental Quality Descriptors

The second data matrix presented for each MU in Sections 3.0 to 7.0 of this report summarizes information about the environmental quality within each tributary. A description of the information included in the matrix for each MU is presented in the subsections below. Table 2.3 provides a brief summary of the same information for quick reference.

Table 2.3: Information sources for environment quality data.

Information Type	Source	Summary of Approach
Water Quality Monitoring Station	Teck in-house	Teck's internal water quality monitoring location code. "P" -permitted station, "NP" -not required by permit. Years of monitoring are stipulated. Provincial EMS number, if one has been assigned. Taken from EQuIS location table.
Water Quality Index	Windward et al. 2014	Water quality index (WQI) was computed using the method of CCME (2001a, b) based on selenium, nitrate, and sulphate concentrations and site-specific benchmarks established for the EVWQP for protection of aquatic life. WQI values corresponding to CCME categories of good and excellent have been identified as "good" (green); WQI values corresponding to CCME categories of marginal and poor have been conservatively identified as "poor" (pink); fair category (yellow) is consistent with CCME (see text and Table 2.3)
Median Selenium, Nitrate-N, Sulphate and Cadmium Concentrations	Windward et al. 2014	Based on routine monitoring by Teck. Samples collected at least monthly, usually weekly during freshet. In parenthesis, the number of samples with concentrations above the site-specific benchmarks is shown relative to the total number of samples analyzed.
Mean	Windward et al. 2014	Based on same data set as described in cell above.
Median Concentrations (2012) and Trends	Zadjlik and Minnow (2013)	Median concentrations reported for 2012, for comparison to combined 2011-2013 data reported by Windward et al. (2014). Trend size and direction for 2004-2012 (or shorter periods in some cases based on data availability) are shown if trend was significant (p<0.1) and actual p-value is presented.
Total Loads, 2012	Zadjlik and Minnow (2013)	Loads were computed as the product of flow times concentration measured within 3 days of each other. Average loads were computed for each month and used to compute the total annual load.
Water Acute Toxicity	Teck (from laboratory reports)	Percent mortality observed for rainbow trout and <i>Daphnia magna</i> in each quarterly test (Q1-Q4) of 2015 is presented for stations required to do such tests under Permit 107517 (BCMOE 2014). In some cases, there was insufficient flow to collect a sample for testing (NS). In a few cases, no explanation was provided for missing test data (ND).
Sediment Quality (Chemistry, Toxicity)	Minnow 2014b	Data collected in 2013 from tributaries were limited to selected settling ponds and near-field lentic areas to characterize "worst-case" mine-related effects on sediment quality (Minnow 2014b; sampling also included some off-channel areas of main-stem streams, and regional lakes that are not presented in this report). Toxicity tests were performed on samples from only two areas associated with tributaries (Henretta Lake [M-1] and Goddard Marsh [MU-3]; data are presented in matrices).
Calcite (Calcite Index)	Windward et al. 2014	Data were collected in 2013 as part of the regional calcite monitoring program (Robinson and MacDonald 2014) and were taken from Appendix E tables of Windward et al. (2014). Calcite presence (score = 1) or absence (score = 0) was recorded for each of 100 pebbles, along with the degree of concretion based on if the particle was removed with negligible resistance (not concreted; score = 0), noticeable resistance but removable (partially concreted; score = 1), or immovable (fully concreted; score = 2). The Calcite Index sums the average scores for presence/absence and concretion of the 100 pebbles assessed in each area.
Benthic Invertebrate Community Structure (% EPT, % Ephemeroptera)	Windward et al. 2014	Data are from the regional aquatic effects monitoring program (RAEMP) completed in 2012 (Minnow 2014a) and were presented in Appendix E tables of Windward et al. (2014). Samples collected in regional reference areas show that healthy benthic invertebrate communities are dominated by EPT larvae (Ephemeroptera [mayflies], Plecoptera [stoneflies], Trichoptera [caddisflies]) and mine-related effects are reflected in reduced numbers of EPT overall or E specifically (Minnow 2014a). Criteria used to categorize community health as good, fair, or poor are presented in Table 2.3.
Maximum HQ for Selenium in Periphyton or Invertebrate Tissue (Direct effects to invertebrates, fish dietary exposure, bird dietary exposure)	Windward et al. 2014	Computed as the highest value of replicate samples collected at that area divided by a screening value taken from the scientific literature. Values greater than one suggest potential for effects. Criteria used to categorize results as good, fair, or poor are presented in Table 2.3.

Surface water quality, sediment quality, calcite index, and biological data were taken from existing reports and generally reflect sampling completed in 2013 or earlier. Additional data were collected in 2014 and 2015 as part of the regular schedule of the regional and local aquatic effects monitoring programs (RAEMP, LAEMPs) and various supporting studies, but were not consistently available among the environmental quality indicators at the time of reporting. It is desirable for the data presented in the matrices to be temporally comparable among data types, so the newer data were not incorporated into the matrices of this version of the data report. It is considered unlikely that conditions have changed sufficiently among areas since those data were collected to affect relative rankings of tributaries during the evaluation process.

2.4.1 Water Quality Monitoring Station

For many tributaries, water quality is monitored by Teck at a single station near the mouth, and reflects the cumulative upstream mining influences. Therefore, the matrices presenting environmental quality data usually represent conditions at this lower tributary location. Some (typically longer, mine-disturbed) tributaries are monitored at more than one location; and, if associated data were reported by Windward et al. (2014), they are provided for more than one area along the tributary. The matrices present Teck's internal code for identifying water quality stations, followed by a "P" or "NP" to indicate if the station is monitored as permit requirement or is a monitoring station that is not required by permit. The historical period of monitoring is also listed, followed by the provincial Environmental Monitoring System (EMS) number, if one has been assigned.

2.4.2 Water Quality Index

The Canadian Water Quality Index (WQI) results were taken from Appendix D of Windward et al. (2014), with additional discussion provided in Appendix E of the same document. The WQI was developed by the Canadian Council of Ministers of the Environment (CCME 2001a, b) as a tool for simplifying the reporting of water quality data and for providing a broad overview of environmental performance. It was based on an approach used in British Columbia (Rocchini and Swain 1995) and is calculated using three factors:

- Scope – This factor represents the number of that exceed the applicable guidelines.
- Frequency – This factor represents the frequency with which (i.e., number of times) variables are greater than the applicable guidelines.

- Amplitude – This factor represents the magnitude by which guidelines are exceeded.

Details regarding the calculation of these factors and the calculation of the overall index value are provided in the technical report and user's manual (CCME 2001a,b).

Windward et al. (2014) used BCMOE water quality guidelines to compute the WQI, except in the cases of selenium, nitrate, and sulphate (for which site-specific benchmarks were used; see Section 2.4.3), and silver (for which the Canadian Council of Ministers of the Environment [CCME] guideline was used). Site-specific benchmarks for selenium, nitrate, and sulphate were the Level 1 (10% effect) benchmarks derived for the EVWQP (Teck 2014).

WQIs were computed by Windward et al. (2014) for different groups of constituents. Generally, it was found that inclusion of variables that rarely exceed guidelines resulted in a higher WQI than when only those exceeding guidelines more frequently were included. Based on these findings, Windward et al. (2014) considered WQI results based on nitrate, nitrite, sulphate, cobalt, selenium, and uranium to be a conservative approximation of overall water quality at each station. This decision was consistent with WQI guidance (CCME 2001b, a), which recommends that only relevant parameters should be included in the evaluation. The same values are presented in the information matrices in Sections 3.0 to 7.0 of this report.

CCME (2001) uses WQI results to interpret overall water quality according to one of the following five categories: excellent, good, marginal, fair, or poor. A simplified set of categories was used for this report in which the "good" and "excellent" categories of CCME were combined into a category called "good", and CCME "marginal" and "poor" categories were combined into a single category called "poor" (Table 2.4). This was done to be consistent with the other data sets that were classified according to the same three categories of good, fair, or poor (Table 2.4).

2.4.3 Concentrations of Selenium, Nitrate, Sulphate, Cadmium

Nitrate, selenium, sulphate, and cadmium are water quality constituents identified as being of concern with respect to coal mining in the Elk Valley (Teck 2014). Median and mean³ concentrations of these substances were taken from Windward et al. (2014) based

³ Standard deviation was not reported by Windward et al. (2014).

Table 2.4: Ranking criteria for observed environmental conditions.

Data Type	Good	Fair	Poor
Water Quality ^a	$80 \leq \text{WQI} \leq 100$	$65 \leq \text{WQI} \leq 79$	$0 \leq \text{WQI} \leq 64$
Sediment Quality ^a	$80 \leq \text{SeQI} \leq 100$	$65 \leq \text{SeQI} \leq 79$	$0 \leq \text{SeQI} \leq 64$
Benthic Community ^b	EPT ^c proportion $\geq 71\%$ and E ^d proportion $\geq 31\%$.	EPT proportion $\geq 64\%$ and E proportion $\geq 12\%$, and also EPT $< 71\%$ or E $< 31\%$ (i.e., data that did not meet criteria for "good" or "poor" categories).	EPT proportion $< 64\%$ or E proportion $< 12\%$.
Tissue Selenium ^e	Maximum Se HQ ^f ≤ 1.0	Maximum Se HQ > 1.0 , but mean Se HQ ≤ 1.0	Mean Se HQ > 1.0

^a Water and sediment quality categories are based on the Canadian Water Quality and Sediment Quality Indices (WQI and SeQI), as defined by CCME (2001a,b, 2007), described in detail in Appendix D of Windward et al. (2014) and summarized in Appendix E, Section E.1 of the same report. CCME categories have been further simplified for this report by designating results corresponding to CCME categories of "marginal" and "poor" as "poor". Additional explanation in text.

^b Based on data from Minnow (2014), as described in Appendix E, Section E.3 of Windward et al. (2014). Additional explanation in text.

^c EPT - Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies).

^d E - Ephemeroptera (mayflies) only.

^e Based on data in Screening-Level Risk Assessment by Windward (2014). Full report is presented in Appendix C of Windward et al. (2014) and approach for categorizing tissue data is presented in Appendix E, Section E.4 of the same report. Additional explanation in text.

^f HQ - Hazard Quotient computed as a ratio of tissue concentration to relevant benchmark (Windward 2014) (Appendix C).

on routine monitoring by Teck at each station in 2011 through 2013. Samples were collected at a frequency of at least monthly, and usually weekly during freshet. The number of samples with concentrations above the Level 1 site-specific benchmarks derived for the EVWQP (Table 2.5; Teck 2014) are shown in parentheses relative to the total number of samples analyzed.

Table 2.5: Level 1 (10%) effect benchmarks identified in the Elk Valley Water Quality Plan.

Constituent	MU	Concentration	Units
Selenium - total	MU1	70	µg/L
	MU2 to MU6	19	µg/L
Nitrate-N	MU1, MU2	$10^{1.0003[\log(\text{hardness})]-1.52}$	mg/L
	MU3 to MU6	3	mg/L
Sulphate	MU1 to MU5	429	mg/L
	MU6	308	mg/L
Cadmium - dissolved	MU1 to MU6	$10^{0.83(\log(\text{hardness}))-2.53}$	µg/L

2.4.4 Median Concentrations, Trends, and Loads

Surface water data presented by Zadajlik and Minnow (2013) included annual summary statistics for selenium, nitrate, sulphate, and cadmium based on data collected at Teck's monitoring stations for the years 2010 to 2012. Median values for 2012 are presented in the tributary matrices in Sections 3.0 to 7.0 of this report for comparison to values presented by Windward et al. (2014) for the combined years of 2011 through 2013 (Section 2.3.3). In addition, Zadajlik and Minnow (2013) computed the magnitude and direction of historical trends using water data for the period 2004 to 2012, or for a shorter period if data for earlier years of this period were lacking. The period of data used to compute trends for cadmium is shown separately from that associated with computing selenium, nitrate, and sulphate trends because laboratory method detection limit issues precluded use of data from some early years at some stations.

The non-parametric Mann-Kendall test was used by Zadajlik and Minnow (2013) to test for the significance of concentrations and load trends. Sen's slopes were computed to indicate the size and direction of the trend. In the information matrices, blue fill was used to indicate a decreasing trend over time and orange fill indicates an increasing trend. The p-values for the significance of the trends are also presented in parentheses for any trends that were significant at $p < 0.1$. P-values ≤ 0.05 were considered strong evidence to reject the null hypothesis of no trend. Stations having p-value > 0.1 were noted as NT

(no trend). Based on a high proportion of cadmium results being reported near or less than the detection limit, the indicated trends for cadmium should be interpreted with caution.

Decreasing concentration trends were identified by Zadjlik and Minnow (2013) at many non- or minimally-exposed monitoring stations, which are suspected to be the result of improvements in analytical methods and detection limits over time, rather than actual water quality trends. Therefore, decreasing trends are not specifically discussed in Sections 3.0 to 7.0. However, those same patterns (of decreasing trends where mine influence is absent or negligible) lends validity to the increasing trends reported by Zadjlik and Minnow (2013); those increasing trends are mentioned in the text, where applicable.

Loads were computed as the product of synoptically-collected⁴ flow and concentration measurements. Values were averaged within months before computing annual loads. Total annual loads were computed using the composite trapezoid rule applied to within-year Julian days and daily loads to estimate the area under the load/time curve. Estimation at the boundaries (Julian day = {0,365} extrapolated the earliest and latest within-year loads available to the boundaries (i.e., to estimate load contributions for months lacking data). Load estimates for stations having at least nine months of within year load data were considered to yield reasonably good estimates of annual total loads (bold load values in matrices), but load estimates based on less than nine months of data (non-bold load values in matrices) should be interpreted with caution. Loads for cadmium were not reported in the matrices due to large uncertainty of results owing to the high proportion of concentrations reported at less than the detection limit.

To provide context, constituent loads for each station were also expressed in the information matrices in Sections 3.0 to 7.0 of this report as a percentage of the sum of the total loads from all monitored tributaries to the Fording River, Michel Creek, or the Elk River in 2012. Load data were not available for most tributaries that have not been previously influenced by mining, as well as for some mine-influenced tributaries; however, contributions from such tributaries are believed to be small. Nevertheless, the percentages presented in the matrices should be considered approximations of the relative (not absolute) contribution of mine-influenced tributaries to main stem loads.

⁴ If a flow measurement was not available for the same day as a chemical measurement, a flow measured up to 3 days before or after the chemical measurement was used (a median flow was used if more than one flow measurement was available).

2.4.5 Surface Water Acute Toxicity

Percent mortality of rainbow trout and *Daphnia magna* observed in each quarterly test (Q1-Q4, 2015) is presented for stations having acute toxicity test requirements in Permit 107517. There were some instances when there was insufficient flow to collect a sample for testing (reported as NS: No Sample). In a few other cases, no explanation was provided for missing test data (reported as ND: No Data). Not applicable (NA) was shown in the matrices in Sections 3.0 to 7.0 of this report for stations that do not have a permit requirement for surface water acute toxicity testing.

2.4.6 Sediment Quality

Sediment samples were collected from a total of 26 depositional areas in the Elk River watershed in early August 2013, including six mine settling ponds, 12 receiving environment areas (e.g., oxbows, lakes or wetlands), and eight reference areas (e.g., oxbows, lakes or wetlands; Minnow 2014b as summarized by Windward et al. 2014). All sediment samples were analyzed for PAHs, particle size, moisture and TOC (whole sediments) and metals (in both the <1 mm and <0.063 mm size fractions). The number of samples having metals or PAH concentrations above the high sediment quality guidelines is shown in the matrices in Sections 3.0 to 7.0 of this report for samples associated with tributaries. However, data relevant to tributaries are sparse due to lack of depositional habitat in most tributaries (other than constructed settling ponds). Some sediment data were collected in 2011 and reported by Windward et al. (2014) but results were not reported in the matrices in Sections 3.0 to 7.0 in this report based on lack of clarity regarding sampling methods and spatial representativeness of samples.

Sediment samples were also collected in 2013 from one settling pond, three receiving environment areas and two reference areas for toxicity testing (using *C. riparius* and *H. azteca*); data relevant to tributaries are available for only Henretta Lake (MU1) and Goddard Marsh (MU3; Minnow 2014b). Sediment chemistry and toxicity were also evaluated at multiple areas throughout the watershed in 2015 based on a study design developed in consultation with BCMOE and KNC (Minnow et al. 2015) and preliminary results were reported to the EMC in early 2016. However, sampling focused again on depositional areas in the watershed that were mainly side channels of the Fording River, Michel Creek, or Elk River. Data will be available in the RAEMP report, scheduled for completion by May 2017, but Henretta Lake (MU1) Goddard Marsh, and Otto Creek (MU3) are the only areas relevant to tributaries where sediment data were collected (those data are not currently included in the matrices).

2.4.7 Calcite

The data presented in the matrices in Sections 3.0 to 7.0 of this report for calcite were collected in 2013 as part of the regional calcite monitoring program (Robinson and MacDonald 2014) and incorporated in the Appendix E tables of Windward et al. (2014). The same locations were sampled in 2014 and 2015 (Robinson and MacDonald 2015, 2016), but the data were not incorporated into the matrices because they would represent a different time period than for the other environmental quality information that was available (i.e., not directly comparable).

For the regional monitoring program, calcite deposition is measured at one to three 100 m-long areas in numerous reaches defined throughout the watershed, depending on the size of the reach (Robinson and MacDonald 2014). At each 100-m area showing evidence of calcite⁵, a modified Wolman (1954) pebble count procedure is applied involving random selection and measurement of 100 substrate particles throughout each 100-m-long area (and distributed in proportion to the habitat types present⁶). The size of each particle is measured (along the intermediate axis; i.e., perpendicular to the longest axis) and the presence (score = 1) or absence (score = 0) of calcite is recorded. In addition, the degree of concretion is assessed by determining if the particle is removed with negligible resistance (not concreted; score = 0), noticeable resistance but removable (partially concreted; score = 1), or immovable (fully concreted; score = 2).

The results for each 100-m-long area are then expressed as a Calcite Index (CI) based on the following equation:

$$CI = CI_p + CI_c$$

Where:

CI = Calcite Index

$$CI_p = \text{Calcite Presence Score} = \frac{\text{Number of pebbles with calcite}}{\text{Number of pebbles counted}}$$

$$CI_c = \text{Calcite Concretion Score} = \frac{\text{Sum of pebble concretion scores}}{\text{Number of pebbles counted}}$$

⁵ The pebble count is not required in areas with no visible calcite, which are simply assigned a calcite index score of 0.0.

⁶ Riffle, cascade, and pool habitats.

For reaches in which multiple 100-m-long areas are sampled, an average CI is computed for the reach and used to monitor changes over time (Robinson and MacDonald 2014, 2015). Results are presented in the matrices in Sections 3.0 to 7.0 of this report for the calcite monitoring area most closely corresponding to the locations of water and biological sampling.

2.4.8 Benthic Invertebrate Community Structure

The data presented in matrices are from the regional aquatic effects monitoring program (RAEMP) completed in 2012 (Minnow 2014a) and were also reported in the Appendix E tables of Windward et al. (2014). Community samples were collected in September 2012 from 36 reference and 56 mine-exposed lotic areas for assessment of potential mine-related effects on community composition. Reference areas were selected to represent a range of natural habitat characteristics (such as elevation, stream size, catchment area, and catchment gradient) corresponding to those for mine-exposed areas. Samples collected in regional reference areas showed that healthy benthic invertebrate communities are dominated by EPT larvae (Ephemeroptera [mayflies], Plecoptera [stoneflies], Trichoptera [caddisflies]) and mine-related effects tend to be most strongly reflected in reduced proportions of EPT overall, or Ephemeroptera (E) specifically (Minnow 2014a).

Criteria used to categorize community health as good, fair, or poor are presented in Table 2.4 based on evaluation of the data collected in 2012. All areas identified as affected or potentially affected by mining (based on detailed statistical analyses of differences between reference and mine-exposed communities) had EPT and/or E proportions that were less than the 5th percentile of all reference areas combined. Conversely, all but one area having EPT and E values equal to or greater than the reference 5th percentile were judged to be within reference condition (i.e., within the ranges exhibited by most reference areas). Therefore, benthic invertebrate communities were categorized as “good” if both EPT and E proportions were equal to or greater than the corresponding reference 5th percentiles (71% and 30%, respectively). Communities were judged to be “poor” if either EPT or E proportions (or both) were less than the minimum values observed among reference areas (64% and 12% respectively). Communities at all other areas were classified as “fair” (i.e., either EPT or E proportion [or both] was [were] below the reference 5th percentile but above the reference minimum). All areas were re-sampled as part of 2015 RAEMP cycle, for which an interpretive report is scheduled to be completed by May 2017 at which time newer data for tributaries will be available. .

2.4.9 Selenium in Periphyton or Invertebrate Tissue

Screening quotients (SQs)⁷ were calculated to estimate potential effects to invertebrates and to fish and birds that consume invertebrates (i.e., dietary exposure). Values were from Windward (2014), which evaluated the potential for ecological effects associated with tissue contaminant levels measured in the Elk Valley. The Windward (2014) report was included as Appendix C in Windward et al. (2014); the tissue selenium results presented therein were also summarized in Appendix E of Windward et al. (2014) and in the matrices in Sections 3.0 to 7.0 of this report.

The following equation was used to calculate the screening quotient for each receptor:

$$SQ = \frac{\text{Observed tissue selenium concentration}}{SV}$$

Where SV = screening value, taken from the scientific literature

The maximum SQ is presented in the matrices, which represents the highest value of replicate samples collected at an area. When the maximum SQ was greater than 1 (suggesting potential for effects), the mean SQ was also reported in parentheses. SQs presented in matrices are based on benthic invertebrates, which are relatively immobile, which makes the data suitable as an indicator of site-specific conditions.

SQs based on selenium concentrations in fish tissues were also computed by Windward (2014) to indicate potential selenium effects to fish themselves and to wildlife consumers, but the data are not presented in matrices because the mobility of fish makes the data unreliable for indicating risks associated with the specific location of fish capture. Criteria used to categorize results as good, fair, or poor are presented in Table 2.4 and are consistent with those used by Windward et al. (2014).

⁷ The documents cited here presented Hazard Quotients (HQ) that are consistent with a generic tissue-based screening-level risk assessment approach. The terminology in the original document (Windward 2014) was subsequently updated to avoid confusion among some external reviewers that the study was intended to fulfil the scope of a Screening Level Risk Assessment as defined by BC's Contaminated Sites Regulation. Instead, the study was completed to: (1) identify trace elements in aquatic organism tissues that may pose an unacceptable risk to consumers via diet-borne toxicity (and direct tissue-based toxicity for selenium and mercury), and (2) recommend any trace elements that should be considered in future tissue monitoring. The term "hazard quotient", reflected in the original documents cited above, was changed to "screening quotient" in the updated version of the Windward (2014) report, but the data and overall approach did not change. So the newer term (SQ) has been used in this report.

2.5 Fish and Aquatic Habitat Information

The third matrix presented for each MU in Sections 3.0 to 7.0 of this report contains information about the fish and aquatic habitat of each tributary. A description of the information included in this matrix for each MU is presented in the subsections below. Table 2.6 provides a brief summary of the same information for quick reference.

2.5.1 Annual Average Flow

Average annual (2013) daily flows were presented in Table 3-8 of Golder (2014b) in units of m³/day, which were converted to m³/s for presentation in the matrices of this report. Average flow for Grace Creek was taken from Teck (2011a). For tributaries not included in Golder (2014b) or Teck (2011a), average annual flows (m³/s) were calculated by Minnow using available data from Teck (variable months and years). In the latter cases, flow data were typically missing for some or all winter months, so annual averages are likely overestimates and have been prefaced with “<” in the matrices in Sections 3.0 to 7.0 of this report.

2.5.2 Ktunaxa Traditional Uses and Cultural Values

The Ktunaxa traditional uses and cultural values will be inserted into the matrices from reporting provided by KNC.

2.5.3 Stream Length

The total stream length (km) was calculated using the provincial CWB dataset and includes all stream lengths within each CWB catchment area. The provincial CWB dataset uses the Fresh Water Atlas (FWA) stream network. The FWA stream network is derived from Terrain Resource Information Management 1 (TRIM1) stream linework (TRIM II streams are not included). TRIM features are delineated through aerial photo interpretation and have varying degrees of accuracy, particularly when defining smaller streams (Norris and Mount, in draft).

The stream lengths within each CWB catchment area were labelled by Teck by type (i.e., connected, fragmented, altered/destroyed or permanently fragmented) based on Teck's dissolved disturbance layer and known barriers identified in the various environmental assessment (EA) Fish and Fish Habitat Baseline Reports. Definitions of the types and sources are provided in Table 2.6.

Table 2.6: Sources of aquatic habitat information.

Information Type			Source(s)	Definition
Annual Average Daily Flow			From Table 3-8 in Golder 2014b, or computed by Minnow based on Equis data provided by Teck.	Annual average daily flow expressed as m ³ /s
Ktunaxa traditional uses and cultural values			To be supplied by KNC	To be supplied by KNC
Stream Length (km and %)	Total		Provincial CWB dataset	The total stream length within the Provincial CWB catchment area
	Currently Connected to Main Stem (km, %)		Provincial CWB dataset/Fish and Fish Habitat Baseline Reports and/or Historical Reports	The total stream length within the Provincial CWB catchment area that is currently connected to the mainstem.
	Currently Fragmented	Potentially Reversible (km, %)	Provincial CWB dataset/Fish and Fish Habitat Baseline Reports and/or Historical Reports	The total stream length within the Provincial CWB catchment area that is currently fragmented by a known barrier that is not passible by fish. Barriers include a known anthropogenic barriers such as a culvert, spillway, or sediment pond, or a known natural barrier such as a waterfall. There is potential for reconnection.
		Permanently Isolated (km, %)	Provincial CWB dataset/Teck's Dissolved Disturbance Layer	The total stream length within the Provincial CWB catchment area that is permanently isolated because it is located upstream from a permanently altered or destroyed section. It still exists but there is no potential for reconnection.
	Permanently Altered or Destroyed (km, %)		Provincial CWB dataset/Teck's Dissolved Disturbance Layer	The total stream length within the provincial CWB catchment area that is permanently altered or destroyed by mining such as by infill.
Fish and Fish Habitat	Fish Species Present		Fish and Fish Habitat Baseline Reports and/or Historical Reports.	Fish species previously documented as being present within in the tributary.
	Unique Fish Habitat Features			Habitat features within a tributary used for a specific life stage for WCT or BT including lakes or pools for overwintering, gravels for spawning or known redd locations, or known juvenile rearing habitat based on the information available.
	Barriers Present excluding infills (A)(Anthropogenic) (N)(Natural)			A known barrier that is not passible by fish. Barriers include a known anthropogenic (A) barrier such as a culvert, spillway, sediment pond or a known natural (N) barrier such as a waterfall. Partial (P) barriers to some life stages of fish are also noted. A zero is present for tributaries that have no barriers. Barriers do not include infills. Barrier listed in this column do not include gradient barriers.
	Total Fish Habitat (km, %)		Provincial CWB dataset/Fish and Fish Habitat Baseline Reports and/or Historical Reports	Stream length and proportion within the Provincial CWB catchment area (i.e., 1980s plus updates) known or inferred to contain fish.
	Total Non-fish Bearing (km, %)			Sections of stream length within the Provincial CWB catchment are that have a slope greater than 25% for perennial streams and 10% for intermittent streams based on the FWA stream network gradient data and do not contain fish upstream.
	Fish Habitat Currently Connected to Main Stem (km, %)			Stream length and proportion within the Provincial CWB catchment area (i.e., 1980s plus updates) that is connected to the main stem and known or inferred to contain fish.
	Fish Habitat Currently Fragmented	Potentially Reversible (km, %)		Stream length and proportion within the Provincial CWB catchment area (i.e., 1980s plus updates) is known or inferred to contain fish but is isolated by a known barrier to fish passage. There is potential for reconnection.
		Permanently Isolated (km, %)		Sections of stream length within the CWB catchment area that are known or inferred to contain fish that are upstream from a permanently altered or destroyed section.
	Fish Habitat Permanently Altered or Destroyed (km, %)			Sections of stream length within the Provincial CWB catchment area that were known or inferred to contain fish but have been permanently altered or destroyed by mining, such as by infill.
	Riparian Habitat (km ² , %)			Provincial CWB dataset/BRE Riparian Habitat Adjacency and Hydrological Approach dataset
Amphibians and Birds			Minnow 2014a (same information presented by Windward et al. 2014)	Amphibian and bird censuses conducted during the 2012 RAEMP focused on lentic reference and mine-exposed habitats of the watershed. Data presented in matrices were usually associated with mine settling ponds in the lower reaches of tributaries. Observed species included in federal or provincial "at-risk" lists are identified in parentheses; bird species identified are not necessarily "aquatic".

Calculations for each stream length type (km and %) are provided in the matrix. The stream length by type is also delineated on the orthophoto figures. All stream lengths (perennial and intermittent) from the CWB dataset FWA (TRIM1) stream network are included in this report. However, it has been acknowledged by BCMOE that there are errors in the dataset and it has not been ground-truthed. For interior portions of the province, TRIM1 may over-represent the number or magnitude of streams (Norris and Mount in draft), including some ephemeral or intermittent streams that have water in them only at the wettest times of the year. It is often found that there is a 'dry draw' at locations where a stream has been shown on the mapping (Norris and Mount, in draft). See Section 2.5.7 for more details.

2.5.4 Fish Species Present

Fish species present in each tributary were identified mainly from Fish and Fish Habitat Baseline Reports associated with approved or planned mine development projects. Two fish species are present in the Elk River watershed that have special conservation status: westslope cutthroat trout (*Oncorhynchus clarkii lewisi*; blue list) and bull trout (*Salvelinus confluentus*; blue list; BCCDC 2012). The BC population of WCT is also listed as special concern in the *Species at Risk Act* under *Schedule 1 List of Wildlife Species at Risk, Part 4 Special Concern*. The baseline reports summarize historical information that was available at the time the reports were prepared, including fish catch data recorded in the provincial Fisheries Information Summary System (FISS).

2.5.5 Unique Fish Habitat Features

A "yes", "no", or "unknown" was indicated in the unique fish habitat feature column based on the Fish and Fish Habitat Baseline Reports and/or historical reports. This column is subjective and based on professional opinion from the information available and likely requires further refinement. The definition of unique fish habitat features is provided in Table 2.6.

2.5.6 Anthropogenic and Natural Barriers Present

An anthropogenic or natural barrier to fish passage was provided in the matrix if identified in the Fish and Fish Habitat Baseline Reports and/or historical reports. The definition of barrier is provided in Table 2.6. Barriers are also shown on the orthophoto figures.

2.5.7 Fish Habitat Lengths

The total fish habitat length (km) was calculated using the stream length within each CWB catchment area. The stream lengths within each CWB catchment area were labelled as

fish habitat (observed and inferred) or non-fish bearing. Methods used were similar to the BCMOE Fish Passage GIS Analysis Methodology and Output Data Specifications Version 2 (Norris and Mount, in draft), which built on the GIS Modelling of Fish Habitat and Road Crossings for the Prioritization of Culvert Assessment and Remediation Version 1 (Mount et al. 2011). The following sources were used for fish habitat determination: stream slope from the FWA stream network, the Canfor stream classification dataset (Canfor 2012), Fish and Fish Habitat Baseline Reports and/or historical reports, and Provincial historical fish observation data (BCMOE 2014).

A GIS analysis process was run to update the stream base data with the Canfor fish bearing classification, known fish-bearing reaches from the Fish and Fish Habitat Baseline Reports and the known fish observations from the BCMOE database. Stream reaches were then updated manually based on connectivity downstream from fish bearing class reaches and for known conditions on the mine site. Stream reaches with a slope greater than 25% (determined from the FWA stream network data) for more than 100 m of stream length on perennial coded streams were set to non-fish bearing if they did not have fish observations upstream. Stream reaches with a slope greater than 10% (determined from the FWA stream network data) for more than 100 m of stream length on intermittently coded streams were set to non-fish bearing if they did not have fish observations upstream. Definitions and sources are provided in Table 2.6. Calculations for each fish habitat length (km and %) by type (i.e., connected, fragmented, altered/destroyed or permanently fragmented) are provided in the matrix. The fish habitat length by type and the location of gradient barriers of 25% on perennial streams and 10% on intermittent streams are also provided on the orthophoto figures.

2.5.8 Riparian Habitat

Methods from the Baldy Ridge Extension Project for ecosystems environmental assessment for riparian habitat (Golder 2015c) were used to quantify the amounts of riparian habitat within each catchment area presented in the matrices of this report. Data are reported as total square kilometres of riparian habitat and as a proportion of the overall basin area. Riparian habitat amount was calculated using both the hydrologic and stream adjacency approaches. By including both approaches, calculations include riparian habitat not connected to the watercourse (hydrologic approach) and riparian habitat connected to the watercourse (stream adjacency approach) since both types were deemed as riparian habitat in the Baldy Ridge Extension Project Environmental Assessment. Both approaches are described in the following paragraphs.

Hydrologic Approach

The hydrologic approach defined riparian habitat as deciduous floodplain and wet forest ecosystem classes (i.e., site series 110, 111, 111x, 112x [i.e., wet forest] where soil moisture regime is 5 or 6 [MacKillop 2012], and flooded low bench tall shrub types [FI] and flooded middle bench deciduous forest [Fm] types [Mackenzie and Moran 2004]) that intersected a buffer area around streams and waterbodies. Buffers were 200 m for stream orders 7 and 8, 100 m for stream orders 5 and 6, 50 m for stream orders 3 and 4 and all other waterbodies, and no buffer for stream orders 1 and 2 (i.e., the stream must intersect the wet forest or floodplain). Deciduous floodplain and wet forest polygons outside the buffer were not included. Similarly, ecosystems within the buffer that were not wet forests or floodplains were not considered riparian habitat using the hydrologic approach.

The hydrologic approach was selected to capture riparian habitat that is not necessarily intersecting a watercourse, but is still defined as riparian habitat because it exists in low-lying areas that may be periodically inundated when water levels are high, and ecological connectivity with the watercourse is thereby maintained.

Stream Adjacency Approach

The stream adjacency approach applied a variable width buffer to streams, ponds, and lakes to define riparian habitat, as follows:

- stream orders 7 and 8 = 50 m.
- stream orders 4, 5 and 6, waterbodies and wetlands = 30 m
- stream order 3 = 20 m.
- stream orders 1 and 2 = 10 m.

High-elevation streams with Biogeoclimatic Ecosystem Classifications (BEC) of Englemann Spruce- Subalpine Fir (e.g., ESSF dkp, ESSF dkw, ESSF wpm, ESSF wmw) and Undifferentiated Interior Mountain (IMA un) were excluded from the analysis because conditions do not support riparian attributes in the ecological context of providing structure and function to support riparian dependent vegetation and wildlife species. Such areas are sub-alpine to alpine ecosystems with extremely harsh conditions that hamper survival of vegetation and, specifically, riparian assemblages. To better understand the availability of intact riparian systems (Environment Canada 2013), the amount of riparian habitat in areas affected by disturbances that remove vegetation (e.g., mining) was calculated. Riparian habitats near clear-cuts were assumed to be vegetated because of regulations

(Tschaplinski and Pike 2010) and best management practices implemented by industry to maintain natural vegetation in riparian habitat (Tembec 2005).

The adjacency approach was selected to capture habitat that plays a role in riparian and aquatic health or function (e.g., shading, streamside deadfall), but that is not necessarily identified as a riparian ecosystem in the BEC system.

2.5.9 Amphibians and Birds

Breeding amphibian and bird censuses were completed at 32 areas during the 2012 RAEMP, and focused on lentic reference and mine-exposed habitats of the watershed (Minnow 2014). Of all 32 areas evaluated, 18⁸ were associated with tributaries, and most of those were constructed mine settling ponds. The bird species identified were not necessarily "aquatic". Bird or amphibian survey data from EA baseline reports are also provided if specific observation locations were reported in tables or maps of the reports (i.e., tributary catchments in which observations were made could be confirmed). The available breeding amphibian and bird survey results can be considered qualitatively (e.g., indicative of locally available habitat suitable for species that were observed), but should not be used quantitatively in the tributary evaluation.

The habitat matrix for each MU identifies the tributary-related survey areas and lists (in parentheses) any species observed that are included in federal or provincial "at-risk" lists. Overall, three species of amphibians were found (Columbia spotted frog [*Rana luteiventris*], western toad [*Anaxyrus boreas*], and long-toed salamander [*Ambystoma macrodactylum*]; Minnow 2014), of which the western toad is considered to be of "special concern" from a provincial or federal conservation standpoint (BCCDC 2012). A total of 125 bird species were observed among areas of which five (bald eagle⁹, barn swallow, great blue heron - *herodias* subspecies, long-billed curlew and olive-sided flycatcher) have special conservation status in the Kootenay District (BCCDC 2012). Of those five species, only the barn swallow, bald eagle and/or olive-sided flycatcher were observed at one or more tributary sampling areas, typically in low numbers (i.e., 1-4 individuals).

⁸ Some of the surveyed areas lacked suitable amphibian breeding habitat so only a bird census was completed.

⁹ Puzzlingly the bald eagle is generally considered secure/not at risk by other lists (e.g., BC Yellow list and COSEWIC).

2.5.10 Written Habitat Descriptions

Text descriptions of aquatic habitat that are presented in Sections 3.0 to 7.0 of this report were taken mainly from Fish and Fish Habitat Baseline Reports associated with EAs for mine development projects:

- Golder Associates Ltd. 2014a. Fording River Operations Swift Project Environmental Assessment Certificate Application. Annex G - Fish and Fish Habitat Baseline Report.
- Golder Associates Ltd. 2015. Coal Mountain Phase 2 Project. Annex I – Fish and Fish Habitat Baseline Report.
- Lotic Environmental. 2015a. Elkview Operations Baldy Ridge Extension Project - Fish and Fish Habitat Baseline Assessment.
- Lotic Environmental. 2015b. Greenhills Operations Cougar Pit Extension Project - Fish and Fish Habitat Baseline Assessment.
- Teck Coal Limited. 2011b. Line Creek Operations Phase II Project Fish and Fish Habitat Baseline Report.

The baseline reports summarize existing information (e.g., from previous technical studies) and the detailed results of any recent habitat surveys that were undertaken to address gaps in the historical baseline information for tributaries of interest. This report did not attempt to cite all original sources of information that were cited in the baseline reports (i.e., only the baseline report itself is cited herein). Exceptions were cases when additional information was found in historical documents that was not presented in baseline documents, in which case the original source was cited. The list of the reports reviewed are provided in the annotated bibliography in Appendix F of this report. Any information that is used to evaluate tributaries in the next phases of the TEP (i.e., Phase C to E in Figure 1.2) will be verified from original sources, if needed to reduce the uncertainty of the evaluation.

As noted above, the habitat descriptions provided in this report were taken mainly from EA fish and fish habitat baseline reports which followed standard assessment methods. Although completed in the context of fish, the habitat descriptions are relevant to any biota utilizing the stream. Additional habitat data are available in various monitoring reports, such as those associated with benthic invertebrate and periphyton sampling locations of the RAEMP and LAEMP, including stream width, depth, and velocity at the time of sampling, plus riparian and substrate characteristics. Such information was generally

redundant with the fish habitat descriptions; therefore it was mentioned only if data from standard environmental assessment baseline reports were sparse or lacking. The applicable baseline report (or alternative source[s]) are cited. Information presented in the sentences following a citation in Sections 3.0 to 7.0 can be assumed to come from the same source until a new source is cited.

Baseline reports identify the number of “reaches” in each tributary, as determined based on reach break analyses that followed standard provincial procedures (BCFISB 2001). Fish and fish habitat surveys generally took either of two approaches:

- Survey of one portion of the stream within each reach, typically an area of about 100-m in length; and/or
- Evaluation of the entire reach according to Level 1 Fish Habitat Assessment Procedures (FHAP; Johnston and Slaney 1996).

Similar information was collected for both types of survey. When both types of information were reported, the description for the longer (FHAP) survey was used for this report.

The baseline reports typically categorized stream morphology according to the BC Channel Assessment Procedures (BCMOF 1996) into one of three main stream morphology categories:

- Riffle-Pool;
- Cascade-Pool; and
- Step-Pool.

The proportions of specific habitat types (e.g., cascade, riffle, glide, pool) observed throughout the survey area were also recorded. Dominant substrate was categorized using the stream substrate size classification of Johnson and Slaney 1996, as follows:

Substrate Type	Substrate Size (mm)
Fines*	<2
Gravels	2 - 64
Cobbles	64 - 256
Boulders	256 - 4000
Bedrock	>4000

* includes sand, silt, and organics

The above information was summarized in the habitat descriptions presented in Sections 3.0 to 7.0, along with stream gradient, bankfull width, riparian vegetation, instream cover, and any reports of off-channel habitat, to the extent all such information

was found. Conclusions that were made in the baseline reports regarding the availability of habitat suitable for various life stages of fish, including spawning, rearing, adult cover and overwintering habitat have also been summarized in this report. Special note was made of stream features confirmed in the field to be barriers to upstream fish movement or unique habitat features (e.g., side-channels, lentic habitat).

2.6 Orthophoto Figures

Orthophoto figures of each tributary were developed in-house by Teck from sources and definitions provided in Table 2.7.

Table 2.7: Tributary features depicted in tributary orthophotos.

Information Type	Source/Explanation
Bridge	The reference dataset is Teck's Infrastructure layer (1:20,000).
Culvert, Dam	Culverts and dams that occur on permanent streams. The reference dataset is Teck's Water Management Locations layer (1:20,000). Lotic dataset from Fish Studies.
Falls	Provincial dataset: CWB Obstructions layer (1:20,000); Natural waterfall - water in a watercourse or a waterbody that follows a perpendicular or a very steep descent. Artificial waterfall - situations where a steep drop occurs in the refined elevation which is not present in the original raw elevation. Artificial waterfalls can occur at both lake banks and stream confluences. Lotic dataset from Fish Studies: vertical drop of 1 -10m. Additional locations identified by sites captured manually (1:20,000).
Gradient Barrier	10% gradient barriers on intermittent streams and 25% gradient barriers on perennial streams were identified from the FWA Stream Network layer data.
Spillway	Structure used to provide the controlled release of flows from a dam or levee into a downstream area. Lotic dataset from for Fish Studies – 7-10m vertical. Additional locations identified by sites captured manually (1:20,000).
Weir	The reference dataset is Teck's Water Management layer (1:20,000). Lotic dataset from for Fish Studies – vertical drop of 1 to 10m.
Pre-Mining Stream Path	Taken from 1952 orthophotos.
Re-aligned channel	Where channels have been realigned or diverted. The reference dataset is Teck's Water Management Lines layer (1:20,000).
Rock drains	Areas where a channel flows through waste rock and has a significant influence on the channel. Can be natural or man made. The reference dataset is Teck's Water Management Lines layer (1:20,000).
Teck Coal Permit Boundaries	The reference dataset is Teck's Permit layer (1:20,000).


3.0 TRIBUTARIES TO UPPER FORDING RIVER (MU1)

The tributaries in the Upper Fording River Watershed (MU1) are shown in Figure 3.1, and include:

- Upstream Fording River,
- Henretta Creek,
- Fish Pond Creek,
- Clode Creek,
- Lake Mountain Creek,
- Kilmarnock Creek,
- Swift Creek,
- Cataract Creek,
- Porter Creek,
- Chauncey Creek,
- Ewin Creek,
- LCO Dry Creek, and
- Greenhills Creek.

Matrices for these tributaries provide data related to watershed area and disturbances (Table 3.1), environmental quality descriptors (Table 3.2) and fish and aquatic habitat (Table 3.3). An orthophoto figure for each tributary is found in Appendix A. Each tributary is briefly described in the following subsections.

Westslope cutthroat trout is the only fish species present in the Upper Fording River, which is defined as the portion of the Fording River extending from the headwaters to Josephine Falls.
















  <p>MAP INFORMATION Datum: NAD 1983 Map Projection: Zone 11N Data Source: Department of Natural Resources Canada. All rights reserved. Creation Date: Sept 2015</p>	<p>Legend</p> <ul style="list-style-type: none">  Water Flow Direction  Upper Fording Watershed  Watershed Boundary  Falls  Community  Waterbody  Streams  Mine Development Areas  Teck Coal Mine Operation  Provincial Boundary 	<p>Figure 3.1. DRAFT Upper Fording River Watershed Boundary</p>  <p>1:125,000</p>
---	---	---

Table 3.1: Watershed area and disturbances associated with Upper Fording River (MU1). Data sources are described in Section 2.0.

Tributary Name	Catchment Area				Stressors																			Mine Alterations/Disturbances/Planned Mitigation							
	Total Catchment Area from Provincial Corporate Watershed Base Dataset (km ²) (1980s plus updates)	Total Catchment Area as of September 2015 (km ²)	% Change in Total Catchment Area	Teck Ownership Rights: % of Catchment Area and Type (Coal, Surface, Timber)	Mine-Related Disturbances in Catchment as of September 2015										Forestry Disturbances in Catchment				Road Disturbances in Catchment												
					Timing of Disturbance: Historical (H), Current (C), Near Future (NF), Far Future (FF) or Not Expected (N)	Total Footprint Area Currently Disturbed (excludes reclaimed lands)		Reclaimed Land Area		Pit(s) (yes/no)	Waste Rock Storage (yes/no)	Rock Drain(s) (yes/no)	Channel Re-alignment or Diversion (yes/no)	Settlement Ponds/Catch Basins (number)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (km ²)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (%)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (km ²)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (%)	Road Density per km ²	Stream Crossings per km	Roads Within 100 m of a Stream	% Roads on Slopes Greater than 60%									
						km ²	%	km ²	%																						
Additional Planned Near/Far Future Mining Disturbances																							Planned Mitigation/Fish Habitat Offsetting								
Upstream Fording River (above Henretta Creek confluence)	38.8	38	-2	53% Coal, 2% Coal and Surface	H,C	1.0	3	0.2	1	Y	Y	Y	N	0	2.0	5	0.9	2	0.8	0.6	0.3	0.0	There are no additional near or far future mine disturbances planned. From a regional water quality management perspective, there is no planned mitigation.								
Henretta Creek	48.7	49.4	1	36% Coal	H,C	3.7	8	1.2	2	Y	Y	N	Y	0	0.0	0	0.0	0	0.6	0.8	0.4	0.0	There are no additional near or far future mine disturbances planned. From a regional water quality management perspective, there is no planned mitigation.								
Fish Pond Creek	0.1	0.1	0	100% Coal	H	0.0	0	0.0	0	N	Y	N	Y	0	0.0	0	0.0	0	3.0	10.0	3.0	0.0	Fish Pond Creek is a potential area for fish habitat offsetting as part of the regional fish habitat management plan that is being developed in consultation with DFO, FLNRO, and KNC representatives. From a regional water quality management perspective, there are no planned mitigation.								
Clode Creek	8	12.9	61	100% Coal	H,C,NF,FF	5.2	65	0.1	1	Y	Y	Y	Y	1	0.0	0	0.0	0	5.7	6.4	2.3	1.4	Mining currently occurring within the permit boundary will continue into the near and far future. From a regional water quality management perspective, the EVWQP identifies that water from Clode Creek will be redirected to the North AWTF as part of the Fording-Swift Project, which is currently planned for 2022.								
Lake Mountain Creek	13	11.5	-12	88% Coal, 22% Coal and Surface, 2% Coal, Surface, Timber	H,NF,FF	3.5	27	1.0	8	Y	Y	Y	Y	1	1.6	12	2.7	21	2.8	1.3	0.9	0.0	Gradual watershed fragmentation will occur as a result of mining (e.g., dewatering, pits, waste rock storage, clean water diversions, and collection of mine-influenced water) as part of the Fording-Swift Project. There are no mitigation options available for Lake Mountain Creek. From a regional water quality management perspective, the EVWQP identifies that water from North Spoil watershed, which will include a portion of flow redirected from Lake Mountain Creek, will be directed to the North AWTF, which is currently planned for 2022.								
Kilmarnock Creek	44.6	41.3	-7	68% Coal	H, C, NF,FF	13.6	30	1.9	4	N	Y	Y	Y	4	3.6	8	0.0	0	1.4	1.4	0.5	0.3	Mining currently occurring within the permit boundary will continue into the near and far future. From a regional water quality management perspective, the EVWQP identifies that water from Kilmarnock Creek will be redirected to the South AWTF, which is currently planned for 2018.								
Swift Creek	9.1	8	-12	71% Coal, 24% Coal and Surface, 4% Coal, Surface and Timber	H,NF,FF	4.5	49	1.0	11	Y	Y	Y	Y	1	0.5	5	0.0	0	5.2	1.7	1.5	0.3	Mining and water management activities (e.g., dewatering, pits, waste rock placement, clean water diversions, and collection of mine-influenced water) will occur as part of the Fording-Swift Project. No fish/aquatic habitat mitigation is planned for the Swift Creek watershed. From a regional water quality management perspective, the EVWQP identifies that water from Swift Creek will be redirected to the South AWTF, which is currently planned for 2018.								
Cataract Creek	6.6	1.8	-73	29% Coal, 73% Coal and Surface	C,NF,FF	5.9	89	0.3	5	N	Y	Y	N	1	0.1	2	0.0	0	3.7	2.9	0.7	0.0	Mining and water management activities (e.g., dewatering, pits, waste rock placement, clean water diversions, and collection of mine-influenced water) will occur as part of the Fording-Swift Project. No fish/aquatic habitat mitigation is planned for the Cataract Creek watershed. From a regional water quality management perspective, the EVWQP identifies that water from Cataract Creek will be redirected to the South AWTF, which is currently planned for 2018.								
Porter Creek	3.4	2.4	-29	100% Coal and Surface	H,C,NF,FF	2.3	68	0.6	18	N	Y	Y	N	1	0.5	15	0.0	0	0.3	0.0	0.2	0.0	Mining currently occurring within the permit boundary will continue into the near and far future. No fish/aquatic habitat mitigation is planned for the Porter Creek watershed. From a regional water quality management perspective, there is no planned mitigation.								
Chauncey Creek	35.6	35.6	0	52% Coal, 1% Coal and Surface	N	0.0	0	0.0	0	N	N	N	N	0	1.8	5	0.3	1	0.7	1.2	0.4	0.0	NA The EVFFHC is in the process of developing a solution for the Chauncey Creek culverts. From a regional water quality management perspective, there is no planned mitigation.								
Ewin Creek	86.7	86.7	0	31% Coal, 1% Coal and Surface	N	0.0	0	0.0	0	N	N	N	N	0	11.4	13	1.7	2	1.0	1.3	0.5	0.1	NA No fish/aquatic habitat mitigation is planned for the Ewin Creek watershed. From a regional water quality management perspective, there is no planned mitigation.								
LCO Dry Creek	28.4	28.4	0	21% Coal, 31% Coal and Surface	C, NF,FF	1.2	4	0.0	0	N	Y	Y	N	3	0.9	3	1.3	5	2.1	2.9	1.1	0.3	The LCO Dry Creek Sediment Pond System was commissioned in 2015 to support mining within the Dry Creek catchment area. Mining and water management activities (e.g., pits, waste rock placement, dewatering, and collection of min-influenced water) to occur as part of the LCO Phase II Project. Condition 14 of the LCO Phase II Environmental Assessment Certificate requires Teck to develop, in consultation with FLNRO, KNC and DFO, an instream flow needs study design for Dry Creek. The EVFFHC is in the process of developing a solution for the LCO Dry Creek culverts at the road and railway crossing. From a regional water quality management perspective, the Dry Creek Water Management Plan has been developed. The initial implementation plan indicates an AWTF and conveyance of bypass water and treated discharge to the Fording River by 2028. The initial implementation plan will be reviewed and revised as additional monitoring and modeling work is conducted under the EVWQP.								
Greenhills Creek	16.5	14.4	-13	100% Coal and Surface	H, C, NF,FF	6.3	38	2.1	13	N	Y	Y	N	7	3.3	20	0.2	1	3.1	2.9	1.5	0.3	Mining currently occurring within the permit boundary will continue into the near and far future. No fish/aquatic habitat mitigation is planned for the Greenhills Creek watershed. From a regional water quality management perspective, the initial implementation plan indicates an AWTF diverting a portion of upper Greenhills Creek by 2026. The initial implementation plan will be reviewed and revised as additional monitoring and modeling work is conducted under the EVWQP. Greenhills Creek has been identified as a priority stream for potential calcite management under the EVWQP.								

NA - Not Applicable

Table 3.2: Environmental quality descriptors for Upper Fording River (MU1). Data sources are described in Section 2.0.

Tributary Name	Track Surface Water Quality Monitoring Station (Code, Permitted-P or Non-Permitted-NP, Years Data Available, and [EMS# if applicable])	Surface Water Chemistry (2011-2013) (from Windward et al. 2014) ^a								Median Concentrations and Trends Blue- and orange-highlighted cells indicate decreasing and increasing trends, respectively (from Zadajlik and Minnow 2013).								Total Loads (2012) (from Zadajlik and Minnow 2013) Bold load values are associated with 9+ months of data. Non-bold values should be interpreted with caution.						Surface Water Acute Toxicity (2015) (Q1-Q4)		Sediment Quality (2013) (Minnow 2014b)		Toxicity Survival and growth of <i>Hyalella azteca</i> and <i>Chironomus riparius</i>	Calcite (2013) (from Windward et al. 2014)	Benthic Invertebrate Community Structure (2012) (Windward et al.		Max SQ ^c for Selenium in Periphyton or Invertebrate Tissue (mean SQ in brackets if max SQ is >1) (Windward et al. 2014)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
		Water Quality Index (WQI) (see Section 2.0 and Table 2.4 for explanation about how WQI was computed and definitions for colour categories)		Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Dissolved Cadmium (ug/L) *Not included in WQI		Data Period for Selenium, Nitrate, and Sulphate Trends		Selenium (µg/L)		Nitrate-N (mg/L)														Sulphate (mg/L)		Cadmium (mg/L)		Rainbow trout (% Mortality)		Daphnia magna (% Mortality)		Chemistry		Colour categories explained in Table 2.4 and text																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
		Median [# Samples >Level 1 BM 19 ug/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 BM of 481 mg/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above			Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Data period for Cadmium Trend	Median Concentration 2012 (ug/L)	Trend (% year) (p-value) - colour codes defined above	Selenium (ton/year)	% of Total Tributary Load	Nitrate (ton/year)	% of Total Tributary Load	Sulphate (ton/year)	% of Total Tributary Load	Chemistry			Direct Effects to invertebrates (periphyton consumption)		Fish Dietary Exposure (Invertebrate Consumption)		Bird Dietary Exposure (Invertebrate Consumption)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							

NA - Not Applicable; ND - No Data; NT - No trend

^a Based on monitoring data and site-specific targets established in EVWQP for nitrate, sulphate, cadmium and selenium (see Appendices A and D in Windward et al. 2014).

^b EPT - Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies).

^c HQ - Hazard Quotient computed as a ratio of tissue concentration to relevant benchmark (Windward 2014). Values above 1 indicate potential risk.

^d Quarterly monitoring data will be summarized in December 2015 version for locations monitored quarterly under Permit 107517.

^e Data for this location may be available but were not assessed by Windward et al. (2014).

Table 3.3: Aquatic habitat of Upper Fording River (MU1). Data sources are described in Section 2.0.

Tributary Name	Annual Average Daily Flow (m ³ /s)	Ktunaxa traditional uses and cultural values	Stream Length based on Province's CWB Layer									Fish and Fish Habitat																Riparian Habitat within Catchment Area		Amphibian census location, date (and any observed species-at-risk)	Bird census location, date (and any observed species-at-risk)
			Total (km) (1980s plus Updates)	Currently Connected to Main Stem		Currently Fragmented		Permanently Altered or Destroyed		Fish Species Present	Unique Fish Habitat Features (Yes/No/Unknown)	Barrier(s) Present (excludes infills)	Total Fish Habitat (1980s plus Updates)		Total Non-fish Bearing (1980s plus Updates)		Fish Habitat Currently Connected to Main Stem		Fish Habitat Currently Fragmented		Fish Habitat Permanently Altered or Destroyed										
						Potentially Reversible	Permanently Isolated												Potentially Reversible	Permanently Isolated											
				km	%	km	%	km	%				km	%	km	%	km	%	km	%	km	%	km	%	km ²	%					
Upstream Fording River (above Henretta Creek confluence)	0.49		78.7	78.7	100	0.0	0	0.0	0	0.0	0	WCT	U	0	19.8	25	58.9	75	19.8	100	0.0	0	0.0	0	0.0	0	2.4	6	ND	ND	
Henretta Creek	0.67		86.8	76.6	88	0.0	0	4.3	5	5.9	7	WCT	Y	P	24.0	28	62.8	72	22.2	93	0.0	0	0.0	0	1.7	7	2.0	4	Henretta Creek (HECK), May 2012	Henretta Creek (HECK), June 2012	
Fish Pond Creek	<0.23 ^a		0.9	0.8	89	0.0	0	0.0	0	0.1	11	WCT	Y	0	0.9	100	0.0	0	0.8	89	0.0	0	0.0	0	0.1	11	0.1	100	ND	ND	
Clode Creek	0.08		15.3	0.8	5	0.0	0	5.7	37	8.8	58	WCT	No	A	4.8	31	10.5	69	0.8	17	0.0	0	0.0	0	4.0	83	0.2	3	Clode Pond, May 2012	Clode Pond, June 2012 (barn swallow)	
Lake Mountain Creek	0.15		23.9	0.2	1	0.0	0	20.7	87	3.0	13	WCT	Y	A	6.5	27	17.4	73	0.2	3	0.0	0	5.7	88	0.6	9	1.1	8	Lake Mountain Lake, May 2012	Lake Mountain Lake, June 2012 (olive-sided flycatcher)	
Kilmarnock Creek	0.70		98.9	0.0	0	0.0	0	59.6	60	39.4	40	WCT	U	A	31.6	32	67.3	68	0.0	0	0.0	0	15.5	49	16.1	51	1.2	3	ND	Settling pond, June 2012 (bald eagle, barn swallow, olive-sided flycatcher)	
Swift Creek	0.04		15.8	0.1	1	0.6	4	5.7	36	9.4	59	WCT	No	A, N	0.1	1	15.7	99	0.1	100	0.0	0	0.0	0	0.0	0	0.2	2	Settling Pond, May 2012 (western toad); Upper catchment (olive-sided flycatcher [2013], western toad [2014])	Swift Creek (SWCK), June 2012	
Cataract Creek	0.05		15.2	0.0	0	0.6	4	0.0	0	14.6	96	WCT	No	N	0.0	0	15.2	100	0.0	0	0.0	0	0.0	0	0.0	0	0.1	2	ND	Cataract Creek, June 2012, July 2013 (olive-sided flycatcher)	
Porter Creek	0.04		5.8	0.4	7	0.8	14	0.4	7	4.2	72	WCT	U	A	0.7	12	5.1	88	0.4	57	0.3	43	0.0	0	0.0	0	0.1	3	ND	ND	
Chauncey Creek	<0.87 ^a		102.4	1.0	1	101.4	99	0.0	0	0.0	0	WCT	U	A	21.9	21	80.6	79	1.0	5	20.8	95	0.0	0	0.0	0	3.2	9	Chauncey Creek, May 2012	Chauncey Creek, June 2012	
Ewin Creek	ND		207.4	207.4	100	0.0	0	0.0	0	0.0	0	WCT	U	P	45.8	22	161.6	78	45.8	100	0.0	0	0.0	0	0.0	0	6.9	8	ND	ND	
LCO Dry Creek	0.28		78.9	1.3	2	57.7	73	19.8	25	0.0	0	WCT	U	A	10.2	13	68.7	87	1.3	13	7.3	72	1.5	15	0.0	0	2.4	8	Off-channel lentic habitat, May 2012 (western toad)	Dry Creek, June 2012 (olive-sided flycatcher)	
Greenhills Creek	0.16		34.7	0.5	1	22.3	64	2.5	7	9.4	27	WCT	Y	A	12.4	36	22.3	64	0.5	4	9.8	79	0.2	2	2.0	16	1.4	8	Settling pond, May 2012	Settling pond, June 2012 (barn swallow)	

ND - no data; WCT - westslope cutthroat trout (Blue-list conservation status)

^a Available data set is missing flows for some low-flow months.

3.1 Upstream Fording River

About 3% of the total catchment area of the Upper Fording River upstream from the confluence with Henretta Creek has been disturbed by mining activities (Table 3.1). The mining disturbance footprint is limited to vicinity of the confluence of the Fording River with Henretta Creek (Appendix Figure A.1). No additional mining disturbances are planned for the near or far future. Of the total stream length of 78.7 km, 100% is connected to the main stem (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

About 400 m of the Fording River upstream from Henretta Creek was enhanced in 1991 to improve the riffle-glide habitat, increase pool habitat, and provide cover through installation of large woody debris (Fording Coal 1991). The Upper Fording River upstream of the Henretta Creek confluence was heavily influenced by the 1995 and 2013 floods including realignment of the channel in 2013. Flood repair works occurred in fall 2015, realigning the channel back to the pre-flood location.

Water quality, benthic invertebrate community, and tissue selenium concentrations were categorized as “good” based on Windward et al. (2014) but calcite data were not reported (Table 3.2).

A 530-m area of stream was surveyed in September 2011, located 4.5 km upstream from the confluence with Henretta Creek (Golder 2014a). The surveyed area consisted of 51% riffle, 31% cascade, 15% glide, and 3% pool habitat, with mean stream gradient of 3% (Photo 3.1). Substrates were primarily cobble, gravel, and boulder. The mean stream gradient was 3% and average bankfull width was 5.2 m. Average mid-channel depth was 0.2 m. Cover was present at only about 8% of the surveyed area and included overhanging vegetation and boulders, with large woody debris in some glides and pools. Riparian vegetation was dominated by young mixed forest or shrubs. Isolated pockets contained substrate suitable for spawning. The Fording River upstream from mining provides suitable habitat for all life stages of WCT when water flows are adequate to support fish. No barriers to fish migration were observed in or near the surveyed area (Golder 2014a).



Photo 3.1 Fording River upstream from Henretta Creek (Golder 2014a).

3.2 Henretta Creek

About 8% of the total catchment area of Henretta Creek has been disturbed by mining activities, which occurred from 1991 to 1997 (Table 3.1). No additional mining disturbances are planned near or far future. Of the total stream length of 86.8 km, 88% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by roads, but not forestry activities (Table 3.1).

Henretta Creek was diverted in the early 1990s into a series of culverts to allow for mining of the Henretta Pit (Wood and Berdusco 1992). Once mining was complete, new channels were constructed at the inlet and outlet of the pit and water was allowed to flow into the pit, creating Henretta Lake and a reclaimed-channel and flow-through lake system (Photo 3.2). Instream reclamation was completed in 1999. In August 2002, V-weirs, small stone lines, wing deflectors, spawning gravel placement, and boulder clusters were constructed in the reclaimed channel downstream from Henretta Lake to increase habitat diversity. The reclaimed channel and lake were heavily influenced by the 2013 floods (Cope et al. 2013).



Photo 3.2: Henretta Lake (Wright et al. 2005).

Water quality, benthic invertebrate community, and tissue selenium concentrations were characterized as “good”, based on Windward et al. (2014), but calcite data were not reported (Table 3.2). However, increasing trends in the concentrations of selenium, nitrate, and sulphate have been reported (Zadjlik and Minnow 2013; Figure 3.2).

Fish and fish habitat were monitored for seven years (1999 to 2006) along the reclaimed channel (Golder 2014a). Surveys conducted in the early 2000s reported that the reclaimed portions of stream were dominated by riffles (55%) and glides (45%) with substrate of cobble and large gravel, gradient of 1.5%, and average depth of about 0.5 m. The substrate was considered too large for WCT spawning and lacking in habitat complexity and resting cover, with the exception of one relatively deep glide and one pool downstream from Henretta Lake (Golder 2014a). Henretta Lake (area of 3.3 ha, mean depth 4.2 m), which is located about 800 m upstream from the Fording River confluence, is a high use area for WCT overwintering (Cope et al. 2014). WCT have been captured in all reaches of Henretta Creek but adult densities are considered to be low upstream from Henretta Lake (Golder 2014a).




Figure 3.2: Selenium concentration trend over time (2004-2013) in Henretta Creek at the confluence with the Fording River (FR__HC1).

Fish and fish habitat were monitored for seven years (1999 to 2006) along the reclaimed channel (Golder 2014a). Surveys conducted in the early 2000s reported that the reclaimed portions of stream were dominated by riffles (55%) and glides (45%) with substrate of cobble and large gravel, gradient of 1.5%, and average depth of about 0.5 m. The substrate was considered too large for WCT spawning and lacking in habitat complexity and resting cover, with the exception of one relatively deep glide and one pool downstream from Henretta Lake (Golder 2014a). Henretta Lake (area of 3.3 ha, mean depth 4.2 m), which is located about 800 m upstream from the Fording River confluence, is a high use area for WCT overwintering (Cope et al. 2014). WCT have been captured in all reaches of Henretta Creek but adult densities are considered to be low upstream from Henretta Lake (Golder 2014a).

Breeding amphibian and bird surveys completed at Henretta Lake in 2012 did not identify any species at risk (Minnow 2014a; Table 3.3).

3.3 Fish Pond Creek

The catchment area of Fish Pond Creek has not been disturbed by mining activities (Table 3.1). There is a diversion ditch that collects and diverts all runoff from the spoil above Fish Pond Creek to the Clode Settling Pond system (Teck internal sources). No additional mining disturbances are planned for the near or far future. Of the total stream length of 0.9 km, 89% is connected to the main stem whereas the remainder has been lost (Table 3.3). The catchment has been disturbed by roads but not forestry activities (Table 3.1). Fish Pond Creek is considered part of the Clode Creek watershed by the CWB dataset.

Fish Pond Creek was targeted for enhancement in the 1980s through 1991 to compensate for effects of mining on Henretta Creek (Wood and Berdusco 1999). This included the construction of overwintering and pocket pools, the addition of spawning gravels, and the addition of various types of riparian vegetation and cover.

No data were presented by Windward et al. (2014) respecting water quality, benthic invertebrate community, calcite levels, or tissue selenium concentrations (Table 3.2).

Fish Pond Creek flows east of and parallel to the Fording River through a wide, flat valley bottom, which is believed to have groundwater influence based on stable, cool water temperatures (Golder 2014a). Fish and fish habitat information was collected to evaluate the effectiveness of enhancement actions in Fish Pond Creek between 1995 and 2002 (i.e. Allan 1996; Amos et al. 2002; Amos and Wright 2000, 2002). The head pond has historically been used by cutthroat trout for overwintering and was deepened in the 1980s and enhanced with natural cover (i.e., root wads and rock groupings; Wood and Berdusco 1999). The creek is considered to provide some of the highest overall habitat quality in the Fording River watershed for all life stages of westslope cutthroat trout, with consistently high population estimates and numerous redds observed (Golder 2014a). Extensive flooding in June 2013 compromised much of the habitat within the creek (Cope et al. 2014).

3.4 Clode Creek

About 65% of the total catchment area of Clode Creek has been disturbed by mining activities (Table 3.1; Photo 3.3). Mining is continuing within the permit boundary and is planned to continue in the near and far future. Of the total stream length of 15.3 km, 5% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by roads but not forestry activities (Table 3.1).

The lower portion of the creek has been moved from its original confluence with the Fording River. On topographic maps based on aerial imagery from 1972, Clode Creek entered the Fording River where the Eagle Sediment Ponds are currently located. The CWB dataset has Clode Creek flowing into Fish Pond Creek. However, Clode Creek has been diverted into the Clode Settlement Pond system adjacent to Fish Pond Creek. The remaining, lower portion of the creek consists of the East Settling Pond (180m x 34m, mean depth 2 m), the Main Settling Pond (150m x 280m; mean depth 2-3 m; collectively referred to as Clode Ponds), and the 100-m section of Clode Creek that discharges from the Main Settling Pond to the Fording River (Golder 2014a). The Clode Ponds were

initially constructed as a settling ponds for removing suspended sediment from water that was impacted by the mining of Clode Pit, which commenced in 1971 (Cope et al. 2014; Golder 2014a).



Photo 3.3: Clode Creek downstream from Clode Ponds (Golder 2014a).

Water quality has been characterized as “poor” based on Windward et al. (2014), with increasing trends in the concentrations of selenium and nitrate (Zajdlik and Minnow 2013; Table 3.2). No data were reported by Windward et al. (2014) for calcite, benthic invertebrate community characteristics or tissue selenium concentrations (Table 3.2).

Active water treatment is planned for Clode Creek, the North Spoil and Swift Pit in 2022 (Teck 2014). Treated water will discharge to the Fording River.

A survey in 2013 reported that the 60-m portion of Clode Creek connecting to the Fording River had mean gradient of 3% and riffle and glide habitats with groundwater inflow and cobble-gravel-fine substrate (Cope et al. 2014, Golder 2014a; Photo 3.3). The West Exfiltration and Grassy Creek channels flowing west and south of the Main Pond, respectively, are mainly groundwater fed and may also convey seepage from the Main

Pond. The area around the Clode Ponds and outflow channels is sometimes referred to as “Clode Flats”. Prior to 1980, large aggregations of fluvial migratory westslope cutthroat trout were identified as overwintering within the groundwater influenced Clode Flats area. Radio tagged fish (n=6) were documented over-wintering at the confluence of Clode Creek and West Exfiltration Creek in both 2013 and 2014 (Cope et al. 2014). Spawning surveys have also identified westslope cutthroat trout redds in lower Clode Creek.

Clode Pond consistently retains water, and vegetation has grown over a large portion of the bottom sediments, making it an ideal rearing ground for westslope cutthroat trout (Wood and Berdusco 1999). After construction of the Clode Ponds, westslope cutthroat trout moved freely between the Fording River and the ponds (Cope et al. 2014). In 2003, the Turnbull South Pit and Spoil Development Application was submitted to reviewing agencies and approved for construction. In the application, it was determined that the East Pond, Main Pond and connecting portion of Clode Creek were going to be directly impacted by spoil (Golder 2014a). In 2004, screens were installed at the seven outflow culverts from the Main Pond to deter fish passage, and a large-scale fish salvage operation was completed (2004 and 2005), which moved almost 6,000 trout from the ponds into the upper Fording River and Fish Pond Creek. However, the salvage did not capture all fish and some culvert screens subsequently failed (which have since been repaired). Also, fish likely re-entered the pond during the record flood in June 2013. Therefore, some westslope cutthroat trout remain in the ponds (Golder 2014a).

Barn swallow was the only species at risk identified in breeding amphibian and bird surveys completed at Clode Pond in 2012 (Minnow 2014a; Table 3.3).

3.5 Lake Mountain Creek

Lake Mountain Creek flows out of the southern end of Lake Mountain Lake and flows in a southeasterly direction into the Fording River (Golder 2014a). About 27% of the total catchment area of Lake Mountain Creek has been disturbed by mining (Table 3.1). Specific portions of the upper Lake Mountain Creek watershed, referred to as the Lake Mountain Spoil, are included in a Memorandum of Agreement (MOA) that documents historically authorized mining activities affecting watercourses in Teck’s area of operations in the Elk Valley (Teck et al. 2011). Gradual watershed fragmentation will occur in the future due to mining (e.g., dewatering, pits, waste rock storage, clean water diversions, and collection of mine-influenced water) as part of the Fording-Swift Project (Table 3.1). Of the total stream length of 23.9 km, 1% is connected to the main stem whereas the

remainder is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

The lower portion of the creek has been moved from its original confluence with the Fording River. On topographic maps based on aerial imagery from 1972, Lake Mountain Creek flowed into the North Tailings Pond. The lower portion of Lake Mountain Creek (~800 m) was realigned about 30 years ago as part of historical mining operations. The CWB dataset has Lake Mountain Creek flowing into the Fording River after it was diverted around the North Tailings Pond, which is consistent with its current confluence with the Fording River.

Water quality, benthic invertebrate community, and tissue selenium concentrations have been characterized as “good” based on Windward et al. (2014), but no calcite data were reported (Table 3.2).

Culverts and a rock drain, which are barriers to upstream fish movement divide Lake Mountain Creek into four reaches (Photo 3.4). Fish and fish habitat surveys were conducted on Lake Mountain Creek in July 2010, with winter surveys also occurring in 2011 (Golder 2014a). A survey of about 100 m of the most downstream reach indicated the habitat is predominantly cascade (40%), riffle (30%), and glide (29%), with mean stream gradient of 6%. The dominant substrates are cobble and gravel, respectively, with a limited amount of boulder. Average bankfull channel width was 4.3 m (Golder 2014a).

The next upstream reach (Reach 2) includes the North Greenhills catchment basin, which is an excavated settling pond, about 160 m long by 80 m wide, with maximum depth of about 1.6 m. The 100-m section of Reach 2 that was surveyed consisted mainly of cascade habitat (62%), with lesser amounts of glide (19%), riffle (12%), and pool (7%) habitats. Mean stream gradient was 7% and the dominant substrate types were cobble, boulder, and bedrock (Golder 2014a).

The third reach of Lake Mountain Creek extends downstream from a rock drain south of Lake Mountain Lake. The 109-m section that was surveyed in 2010-11 consisted of 47% cascade, 32% riffle, and 21% glide. The mean stream gradient was 6% and the substrate was dominated by cobble, with lesser amounts of gravel or finer particles. Average bankfull channel width in Reach 3 was 4.5 m (Golder 2014a).



Photo 3.4: Culvert at Clode Pond access road in the lower section of Lake Mountain Creek (Golder 2014a).

Reach 4 of Lake Mountain Creek flows for about 600 from Lake Mountain Lake to the rock drain (Golder 2014a). This portion of stream includes cascades, glides and riffles, with some pool habitat and mean gradient of 5%. Lake Mountain Lake is situated in a coniferous forest with grass and sedge lining the shallow sloped shoreline. The maximum water depth is 2.65 m, and average water depth is 1.14 m (Golder 2014a).

Westslope cutthroat trout is the only species of fish present in Lake Mountain Creek (Golder 2014a). There is overwintering habitat available and suitable habitat is also available for all life stages of westslope cutthroat trout during the open-water season, especially in the lower reaches. Pockets of substrate suitable for spawning by westslope cutthroat trout are present. Continued presence of fish in the fragmented reaches suggests that they contain habitat that is adequate to support all life stages and functions. The lake may also support westslope cutthroat trout, although open water fish surveys conducted in 2010 and 2011 using fyke nets, minnow traps, and angling did not yield any

fish. Under-ice dissolved oxygen levels were low in the lake (0.08 to 1.51 mg/L) indicating low likelihood for successful fish overwintering (Golder 2014a).

Olive-sided flycatcher was the only species at risk identified in breeding amphibian and bird surveys completed at Lake Mountain Lake in 2012 (Minnow 2014a; Table 3.3).

3.6 Kilmarnock Creek

About 30% of the total catchment area of Kilmarnock Creek has been disturbed by mining activities (Table 3.1). Mining is currently occurring within the permit boundary and will continue in the near and far future (Table 3.1). All of the stream length of 98.9 km is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

The upper reaches of Kilmarnock Creek (and the most upstream portion of Brownie Creek) remain relatively undisturbed apart from adjacent forest harvesting (Table 3.1). The lower reaches have been disturbed by mining and forestry (Fording Coal 1985; Golder 2014a,b). Kilmarnock Creek flows through a rock drain, as well as a settling pond system that was constructed in 1989. This is a conventional settling pond, where a portion of the total flow infiltrates to ground and the pond is empty for most of the year.

Water quality and benthic invertebrate community was characterized as “poor” based on Windward et al. (2014), with increasing trends in the concentrations of selenium, nitrate and sulphate (Zadjlik and Minnow 2013; Table 3.2). Tissue selenium concentrations have been characterized as “good” (Windward et al. 2014; Table 3.2). High levels of calcite have been reported (calcite index: 2.2).

The EVWQP committed to the construction of Active Water Treatment Facilities (AWTF) to manage water associated with the existing and future coal mining development (Golder 2014d). The FRO South AWTF will be commissioned in 2018 and will include treatment of water from Kilmarnock Creek.

Historically (e.g., 1952), the channel geomorphology of Kilmarnock Creek was meandering, transitioning from a confined to wide valley, with braiding on the alluvial fan where the tributary met with Fording River (Golder 2014b). Riparian vegetation was primarily established forest. Avalanche paths connected directly to the channel in upper reaches but less so in lower reaches where the valley widened. Prior to 1980, large aggregations of fluvial migratory westslope cutthroat trout were identified within Kilmarnock Creek (Cope et al. 2014). The lower portion of the creek prior to the construction of the sediment pond system was characterized by shallow riffles and

moderately deep glides (Fording Coal 1985). Some overwintering habitat was present in Kilmarnock Creek prior to the construction of the sediment pond system (Fording Coal 1985).

The lower section of Kilmarnock Creek has been substantially altered and portions are buried. As the creek approaches the mine footprint it braids prior to discharging into a lake formed against the edge of a waste rock spoil that has buried the stream channel. This permitted habitat loss occurred in the late 1980's as part of the Eagle Mountain Project. (Cope et al. 2014). An isolated population of westslope cutthroat trout remains in Kilmarnock Creek upstream from the rock drain (Golder 2014a). There are no fish present in the Kilmarnock settling pond system. About 3,870 m of Brownie Creek and associated streams were assessed in November 2002 and June 2003 (Edeburn 2003 and Berdusco 2003). Despite habitat suitable for overwintering (Edeburn 2003), no fish were captured or observed (Berdusco 2003). The remainder of Brownie Creek and its tributaries were considered non-fish bearing due to a gradient barrier to upstream fish migration and a lack of suitable habitat in 2002 (Edeburn 2003).

Bald eagle, barn swallow and olive-sided flycatcher are species at risk that were identified in breeding amphibian and bird surveys completed at Kilmarnock settling ponds in 2012 (Minnow 2014a; Table 3.3).

3.7 Swift Creek

About 49% of the total catchment area of Swift Creek has been disturbed by mining activities (Table 3.1). Additional mining and water management activities will occur in the future as part of the Fording-Swift Project (Table 3.1). Of the total stream length of 15.8 km, 1% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

Water quality and the benthic invertebrate community was characterized as “poor” based on Windward et al. (2014), with increasing trends in the concentrations of selenium, nitrate and sulphate (Zadjlik and Minnow 2013; Table 3.2). Tissue selenium concentrations were characterized as “good” (Table 3.2). High levels of calcite were reported (calcite index: 2.2).

The EVWQp committed to the construction of AWTFs to manage water associated with the existing and future coal mining development (Golder 2014d). The FRO South AWTF will be commissioned in 2018 and will include treatment of water from Swift Creek.

The remnant portion of Swift Creek includes cascade and glide habitats (Golder 2014a). Prior to a flood in 2013, Swift Creek flowed over a bedrock falls and flowed south along the base of a slope for about 300 m before entering the Fording River. Historical surveys reported the presence of fish downstream but not upstream from the falls (RL&L 1995; Golder 2014a). In May-June 2010, no substrate suitable for spawning was observed in the downstream portion of stream (due to calcified substrate) nor were spawning westslope cutthroat trout or constructed redds observed but fish were captured in July of the same year (Golder 2014a).

During the 2013 flood event, Swift Creek avulsed its banks and flowed overland for 22 m into the Fording River (Photo 3.5), leaving the former channel dry other than a few isolated pools (Golder 2014a). This has resulted in 5-m-high waterfall 22 m upstream from the Fording River that represents a barrier to upstream movement by all fish life stages. A survey in October 2013 (post-flood) documented average wetted width and mid-channel depth of 3.9 m and 0.13 m, respectively. The substrate was heavily calcified. The shallow braided habitat downstream from the waterfall likely limits use of the lower creek by adult westslope cutthroat trout but could serve as rearing and overwintering habitat for juvenile and young-of-year westslope cutthroat trout. The reach upstream from the waterfall was considered non-fish-bearing (Golder 2014a).

Western toad was the only species at risk identified in breeding amphibian and bird surveys completed at the Swift Creek settling pond in 2012 (Minnow 2014a; Table 3.3). An olive-sided flycatcher (July 2013) and a western toad (July 2014) were also observed in the headwaters of the Swift Creek basin (Matrix Solutions 2015).

3.8 Cataract Creek

About 89% of the total catchment area of Cataract Creek has been disturbed by mining activities (Table 3.1). Additional mining and water management activities will occur as part of the Fording-Swift Project (Table 3.1). All of the total stream length of 15.2 km is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

Water quality, benthic invertebrate community and tissue selenium concentrations have been characterized as “poor” based on (Windward et al. 2014), and increasing trends in the concentrations of selenium, nitrate, sulphate and cadmium have been reported (Zadjlik and Minnow 2013; Table 3.2). High levels of calcite were reported (calcite index: 3.0).



Photo 3.5: Swift Creek at the Fording River post-flood 2013 (Golder 2014a).

The EVWQP committed to the construction of AWTFs to manage water associated with the existing and future coal mining development (Golder 2014d). The FRO South AWTF will be commissioned in 2018 and will include treatment of Cataract Creek water.

Fish sampling was conducted in May 2010 and habitat mapping was completed for a 100-m section of stream in the lower watershed in July 2010 (Golder 2014a). The surveyed section was located 125 m upstream from a 15-m-high waterfall located at the confluence of Cataract Creek with the Fording River, which is considered a barrier to fish migration. Therefore, the creek is non-fish-bearing, which was confirmed by lack of fish capture during 842 seconds of electrofishing in May 2010. Sections of the creek that were surveyed had predominantly cascade habitat (63%), with some glide (22%), riffle (8%) and pool (7%) habitat. The maximum depth of pools was 0.45m. The mean stream gradient was 11% (Golder 2014a).

Olive-sided flycatcher is a species at risk that was identified in breeding amphibian and bird surveys completed in 2012 (Minnow 2014) and in 2013 (Matrix Solutions 2015).

3.9 Porter Creek

About 68% of the total catchment area of Porter Creek has been disturbed by mining activities (Table 3.1). Mining is currently occurring within the permit boundary and will continue in the near and far future (Table 3.1). Of the total stream length of 5.8 km, 7% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

Water quality was characterized as “fair”, the benthic invertebrate community was characterized as “poor”, and tissue selenium concentrations were characterized as “good” based on Windward et al. (2014; Table 3.2). An increasing trend in the concentration of sulphate was reported, while no trends were observed in the concentrations of selenium, nitrate, or cadmium (Zadjlik and Minnow 2013; Table 3.2). Low levels of calcite have been reported in Porter Creek (calcite index: 0.9).

The settling pond was likely built as part of the Greenhills Surface Coal Mining Project after 1981. The portion of the creek between the settling pond and the Fording River is assumed to be fish-bearing because no barriers are present. No fish were captured in 325 m of channel that was electrofished in September 1979 and field observations indicated the creek had low fish potential (BC Research 1981).

3.10 Chauncey Creek

None of the catchment area of Chauncey Creek has been disturbed by mining activities and is not expected to be in the future (Table 3.1). Of the total stream length of 102.4 km, 1% is connected to the mainstem and the remainder is fragmented (Table 3.3). The catchment has also been disturbed by forestry (Table 3.1) and road construction (Golder 2014b).

Water quality and the benthic invertebrate community were characterized as “good” based on Windward et al. (2014; Table 3.2). Tissue selenium concentrations were not reported for Chauncey Creek but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”; Table 3.2). No calcite was observed in the creek.

Five sections of Chauncey Creek were surveyed in August 1995 to identify the number of reaches, stream channel widths and gradients, fish distribution and identification of species present (Russell and Oliver 1996). In the downstream section of the creek, about 200 m downstream from the highway and near the confluence with the Fording River, the gradient was low at 1% and the average channel width was 8.5 m. The surveyed section

was predominantly pool/glide features with abundant gravel substrate suitable for westslope cutthroat trout. A second area was sampled 1 km upstream from a logging road bridge crossing, and was characterized by a slightly steeper gradient (3%), a narrower channel width (6.9 m), equal proportion of riffle, pool and run habitats, and substrates of boulder and gravel. The upper sections of the creek increased in gradient from 2% to 20%, contained intermittent subsurface flow with increasing substrate size. Riparian vegetation was typically mixed forest with some sections dominated by shrubs (e.g., willow, alder, fir and spruce; Russell and Oliver 1996).

The culvert at the Chauncey Creek crossing of the Fording River Road is a barrier to westslope cutthroat trout migration (Russell and Oliver 1996). Relatively high densities of westslope cutthroat trout are found in the Fording River in the vicinity of Chauncey Creek where the habitat is dominated by deep pools, and high juvenile densities have been observed in Chauncey Creek downstream from the culvert (Cope et al. 2014). By comparison very low densities of both juvenile and mature westslope cutthroat trout are present in upper Chauncey Creek. Restoring the connectivity of the mainstem Fording River to the tributary habitat of Chauncey Creek represents an opportunity for habitat improvement in the upper Fording River watershed (Cope et al. 2014).

No species at risk were identified in breeding amphibian and bird surveys conducted at an area in lower Chauncey Creek in 2012 (Minnow 2014a; Table 3.3).

3.11 Ewin Creek

None of the catchment area of Ewin Creek has been disturbed by mining activities (Table 3.1). No mining is planned for the future (Table 3.1). Of the total stream length of 207.4 km, 100% is connected to the main stem (Table 3.3). The catchment has been disturbed by forestry activities (Table 3.1) and road construction (Golder 2014b).

The benthic invertebrate community of Ewin Creek was characterized as “good” based on Windward et al. (2014; Table 3.2). Water quality and tissue selenium concentrations were not reported by Windward et al. (2014) but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). No calcite was observed.

Ewin Creek was surveyed in August 1995 to identify the number of reaches, stream channel widths and gradients, fish distribution and identification of species present (Russell and Oliver 1996). The first surveyed section was located about 30 m upstream from the FRO access road bridge, lying within the floodplain of the Fording River. It had a mean gradient of 1% and an average channel width of 7.1 m. This reach was

predominantly pool/run complex with a streambed dominated by cobble with some gravel suitable for spawning (Russell and Oliver 1996). A second survey area was about 500 m downstream from the tributary locally referred to as the Ewin Side Draw, near the confluence with Toddhunter Creek. The stream channel in this area was confined, with a steeper average gradient (3%) and an average width of 4.7 m. It had equal distribution of riffles, runs and pools, with composed predominantly of cobble and gravel. A third section the stream was surveyed upstream from a series of beaver ponds leading to a marshier habitat. The sampled area upstream from the marshland had 2% gradient, and an average channel width of 4.2 m. This section of the creek had signs of extreme flooding (2 m), including significant bedload movement and debris torrents. The habitat was predominantly riffle with some runs and pools. A fourth section of stream 200 m upstream from a former bridge crossing was also surveyed and determined to be highly impacted by a previous flood at the lower end. The channel gradient was low (1%) and the width was wide (13.4 m) due to the creek overflowing onto the adjacent logging road. At the upper end of the surveyed section the impact of flooding was less, and the reach was much steeper with a 15% gradient and a step-pool morphology dominated by cobble and boulders and a channel width of 6 m. A final section was surveyed closer to the headwaters, adjacent to a landing in a logging block. This section had a gradient of 21% and was completely subsurface flow, although some evidence of spring freshet was found. Riparian vegetation throughout the watershed consisted of mixed forest (willow, alder, spruce; Russell and Oliver 1996).

Past surveys indicated westslope cutthroat trout are absent or present only in low densities in Ewin Creek (Russell and Oliver 1996; Cope et al. 2014) despite abundant potential spawning and rearing habitat, especially in the lower reaches (Hooton et al. 1971), including undercut banks, log and debris jams, boulders and pools. This may be related to low annual mean temperature (7.2 °C) compared to the temperatures preferred by westslope cutthroat trout (9-13°C; Cope et al. 2014). Also, the culvert where the Fording River Road crosses Ewin Creek is considered to be a partial barrier to fish migration. One juvenile fish was documented to move through the culvert in 2015 (Cope et al. 2016, in draft). Tributaries to Ewin Creek (Toddhunter Creek, Ewin Side Draw, unnamed tributaries) were also surveyed and no fish were captured (Russell and Oliver 1996).

3.12 LCO Dry Creek

About 4% of the total catchment area of LCO Dry Creek has been disturbed by mining (Table 3.1). Additional mining and water management activities will occur as part of the

LCO Phase II Project. Of the total stream length of 78.9 km, 2% is connected to the main stem whereas the remainder is fragmented (Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

LCO's Dry Creek flows north from its headwaters to the Fording River. An unnamed tributary (often referred to as the East Tributary), joins Dry Creek from the east about 5 km upstream from the confluence with the Fording River. A railway and two road crossings are located along the lower portion of Dry Creek. The railway and one road crossing are paired culvert crossings (Photo 3.6), whereas the second road crossing is a clear-span bridge.



Photo 3.6: Paired culvert crossing under Fording Road in Dry Creek (Teck 2011b).

Recent water quality and the benthic invertebrate community were characterized as “good” (Windward et al. 2014; Table 3.2). Windward et al. (2014) did not report any tissue selenium concentrations for Dry Creek (Table 3.2). No calcite was observed in the creek.

Despite a considerable amount of fishing effort in field surveys conducted in 2009 (i.e., total 16,021 seconds electrofishing effort over a total distance of 5,856 m) only 28 fish were captured in Dry Creek (Teck 2011b). Low fish densities were also reported by BC Research (1981) in surveys conducted downstream from the FRO access road in July and September 1979 and June 1980. Fish presence appears to be limited to the lower 4.2 km of the creek (i.e., mainly Reaches 1 to 3) where habitat is suitable for use by westslope cutthroat trout for feeding as well as some suitable spawning habitat. Stream gradient in this portion of creek averages about 3% and is dominated by riffle and glide habitats, with some pools and cascades. Substrate was predominantly cobble, with variable amounts of gravel, sand, and boulders. Bankfull and mid-channel water depths (September 2009) were about 0.45 m and 0.2 m, respectively. About 20% of the stream has cover in the form of undercut banks, overhanging vegetation and large woody debris. In the reaches farther upstream and in the East Tributary, stream size is smaller, water depths were shallower, there was more cascade habitat with fewer pools and glides, and less cover (Teck 2011b).

The two culvert crossings under the Canadian Pacific railway line and the FRO access road, may be barriers to fish movement under some flow conditions (Teck 2011b). A 250-m section of channel about 3.5 km upstream from the Fording River goes dry during low-flow periods as does a section of the East Tributary just upstream from the confluence with Dry Creek. A 1-m-high debris-created waterfall, located about 4.5 km upstream from the Fording River, presents another potential barrier to upstream movement of fish. Seasonally dry sections and the debris-created waterfall may account for the apparent absence of fish in the upper reaches of Dry Creek and the East Tributary. Most of Dry Creek was frozen to the bottom in February 2010 or lacking sufficient water depth to support fish.

Breeding amphibian and bird surveys conducted in an off-channel lentic area of Dry Creek in 2012 reported the presence of two species at risk: western toad and olive-sided flycatcher (Minnow 2014a; Table 3.3).

3.13 Greenhills Creek

About 38% of the total catchment area of Greenhills Creek has been disturbed by mining activities (Table 3.1). Mining is currently occurring within the permit boundary will continue into the near and far future (Table 3.1). Of the total stream length of 34.7 km, 1% is connected to the main stem whereas the remainder is fragmented or has been lost

(Table 3.3). The catchment has also been disturbed by forestry activities and roads (Table 3.1).

Based on samples collected near the mouth of the creek, water quality, benthic invertebrate community and tissue selenium concentrations were characterized as “poor”, with the exception of bird dietary selenium exposure (i.e., invertebrate consumption), which was characterized as “good” (Windward et al. 2014; Table 3.2). Increasing trends in the concentrations of selenium, nitrate, sulphate and cadmium were reported (Zadjlik and Minnow 2013; Figure 3.3) and low levels of calcite have been observed (calcite index: 0.4; Active water treatment is scheduled to begin in 2026, treating water from GHO’s West Spoil (influencing Thompson, Leask, and Wolfram Creeks) as well as a portion of upper Greenhills Creek (Teck 2014). Treated effluent will discharge to Thompson Creek.




Figure 3.3: Selenium concentration trend over time (2004-2013) in Greenhills Creek at the confluence with the Fording River (GH_GH1).

In the environmental assessment associated with development of GHO, BC Research (1981) identified that the lower 2 km of Greenhills Creek was formerly used extensively by cutthroat trout for spawning and rearing. During mine development, a settling pond (Greenhills Pond) was constructed, which is located about 600 m upstream from the Fording River. A dam spillway and culverts located immediately downstream from the pond are currently barriers to the upstream migration of fish. Fish surveys conducted in 1970 and 1980 prior to construction of the Greenhills Pond showed lower densities upstream from the road culvert compared to downstream, suggesting that fish movement was impaired at the road culvert (BC Research 1981). A study of the creek in July 2007

(KNRC 2007) reported that, upstream from the pond, the main stem of the creek is fish-bearing for about 4.9 km and westslope cutthroat trout have free access between the creek and the pond (KNRC 2007). Over this reach, the gradient is about 4% and adequate habitat is available to support all life-stages of fish, including spawning and overwintering. Further upstream, the steep gradient, including a falls and long cascade, act as a barrier to upstream fish movement, so no westslope cutthroat trout are present. A tributary to the west (Gardine Creek) offers another 1.8 km of fish-bearing habitat suitable for all life stages of westslope cutthroat trout, with gradients between 4% and 7%. Portions of the stream both upstream and downstream from Greenhills Pond are affected by calcite deposition (Robinson and MacDonald 2014).

Barn swallow, is a species at risk that was observed near the Greenhills Pond in breeding bird and amphibian surveys conducted in 2012 (Minnow 2014a; Table 3.3).

4.0 TRIBUTARIES TO LOWER FORDING RIVER (MU2)

The tributaries in the Lower Fording River Watershed (MU2) are shown in Figure 4.1, and include:

- Grace Creek; and
- Line Creek.

Matrices for these tributaries provide data related to watershed area and disturbances (Table 4.1), environmental quality descriptors (Table 4.2) and fish and aquatic habitat (Table 4.3). An orthophoto figure for each tributary is found in Appendix B. Each tributary is briefly described in the following sections.



4.1 Grace Creek

None of the catchment area of Grace Creek has been disturbed by mining activities (Table 4.1; Photo 4.1). No mining disturbances are planned for the future (Table 4.1). Of the total stream length of 45 km, 31% is connected to the main stem whereas the remainder is fragmented (Table 4.3). The catchment has been disturbed by forestry activities and roads (Table 4.1).

Water quality was characterized as “good” based on Windward et al. (2014; Table 4.2). Benthic invertebrate community data and tissue selenium concentrations were not reported for Grace Creek but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). Low levels of calcite were reported (calcite index: 0.31; Table 4.2).

The lower portion (~2.6 km) of Grace Creek is dominated by glide and riffle habitat with gravel substrate and mean stream gradient of about 2.2% (Teck 2011b). In October 2010, mean bankfull channel width and depth were 3.15 m and 0.34 m, with an average wetted width and mid-channel depth of 2.71 m and 0.21 m. Reaches farther upstream have steeper gradient (about 5-8%; Photo 3.7) with predominantly riffle and cascade habitats and substrate dominated by cobble, gravel, or sand. Stream cover types include large and small woody debris, undercut banks, and overhanging vegetation.

Two corrugated steel pipe culvert crossings culverts are located several kilometres upstream from the Fording River (Teck 2011b). The lower culvert, associated with a rail road crossing, is 100 m long with a gradient of 5% and an outflow height of greater than 1 m. The second is a road crossing located upstream from the rail crossing that includes



MAP INFORMATION
 Datum: NAD 1983 Map Projection: Zone 11N
 Data Source: Department of Natural
 Resources Canada. All rights reserved.
 Creation Date: Sept 2015

Legend

- | | |
|----------------------|--------------------------|
| Water Flow Direction | Waterbody |
| Lower Fording River | Streams |
| Watershed Boundary | Mine Development Areas |
| Falls | Teck Coal Mine Operation |
| Community | Provincial Boundary |

**Figure 4.1. DRAFT
 Lower Fording River
 Watershed Boundary**

Teck

1:125,000

Table 4.1: Watershed area and disturbances associated with Lower Fording River (MU2). Data sources are described in Section 2.0.

Tributary Name	Catchment Area				Stressors																	Mine Alterations/Disturbances/Planned Mitigation		
	Total Catchment Area from Provincial Corporate Watershed Base Dataset (km ²) (1980s plus updates)	Total Catchment Area as of September 2015 (km ²)	% Change in Total Catchment Area	Teck Ownership Rights: % of Catchment Area and Type (Coal, Surface, Timber)	Mine-Related Disturbances in Catchment as of September 2015										Forestry Disturbances in Catchment				Road Disturbances in Catchment				Additional Planned Near/Far Future Mining Disturbances	Planned Mitigation/Fish Habitat Offsetting
					Timing of Disturbance: Historical (H), Current (C), Near Future (NF), Far Future (FF) or Not Expected (N)	Total Footprint Area Currently Disturbed (excludes reclaimed lands)		Reclaimed Land Area		Pit(s) (yes/no)	Waste Rock Storage (yes/no)	Rock Drain(s) (yes/no)	Channel Re-alignment or Diversion (yes/no)	Settlement Ponds/Catch Basins (number)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (km ²)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (%)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (km ²)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (%)	Road Density per km ²	Stream Crossings per km	Roads Within 100 m of a Stream	% Roads on Slopes Greater than 60%		
Grace Creek	18.6	18.6	0	44% Surface and Coal, 1% Coal, Surface and Timber	N	0.0	0	0.0	0														N	N
Line Creek	138.0	138.0	0	37% Coal, 6% Surface and Coal, 18% Surface and Coal and Timber	H, C, NF, FF	18.7	14	3.9	3	Y	Y	Y	N	4	16.6	12	0.0	0	1.7	1.3	0.7	0.1	Mining currently occurring within the permit boundary in the Line Creek catchment area will continue into the near and far future.	Line Creek is a potential area for fish habitat offsetting as part of the regional fish habitat management plan that is being developed in consultation with DFO, FLNRO, and KNC representatives. From a regional water quality management perspective, the West Line Creek AWTF is currently operating downstream of the rock drain in Line Creek. Fish were removed from the Contingency Ponds in 2015, permanent exclusion is planned. A fish salvage is planned in 2016 for MSAN Ponds to remove fish. Water is being pumped from the Horseshoe pit as part of the routine flood management plan to the rock drain in Line Creek. Water is being pumped from the BRS pit to maintain daily operations to the No-name drainage where it infiltrates to the West pit then the rock drain in Line Creek.

NA - Not Applicable

Table 4.2: Environmental quality descriptors for Lower Fording River (MU2). Data sources are described in Section 2.0.

Tributary Name	Tack Surface Water Quality Monitoring Station (Code, Permitted-P or Non-Permitted-NP, Years Data Available, and [EMS# if applicable])	Surface Water Chemistry (2011-2013) (from Windward et al. 2014) ^a										Median Concentrations and Trends Blue- and orange-highlighted cells indicate decreasing and increasing trends, respectively (from Zadjlik and Minnow 2013).										Total Loads (2012) (from Zadjlik and Minnow 2013) Bold load values are associated with 9+ months of data. Non- bold values should be interpreted with caution.						Surface Water Acute Toxicity (2015) (Q1-Q4)		Sediment Quality (2013) (Minnow 2014b)		Calcite (2013) (from Windward et al. 2014)	Benthic Invertebrate Community Structure (2012) (Windward et al.		Max SQ ^c for Selenium in Periphyton or Invertebrate Tissue (mean SQ in brackets if max SQ is >1) (Windward et al. 2014)	
		Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Dissolved Cadmium (ug/L) *Not included in WQI		Data Period for Selenium, Nitrate, and Sulphate Trends		Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Cadmium (mg/L)		Rainbow trout (% Mortality)													Daphnia magna (% Mortality)		Chemistry	Toxicity Survival and growth of <i>Hyalella azteca</i> and <i>Chironomus riparius</i>
		Median [# Samples >Level 1 BM 19 ug/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 BM of 481 mg/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean			Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Data period for Cadmium Trend	Median Concentration 2012 (µg/L)	Trend (% year) (p-value) - colour codes defined above	Selenium (ton/year)	% of Total Tributary Load	Nitrate (ton/year)	% of Total Tributary Load	Sulphate (ton/year)	% of Total Tributary Load										
		Water Quality Index (WQI) (see Section 2.0 and Table 2.4 for explanation about how WQI was computed and definitions for colour categories)	Median [# Samples >Level 1 BM 19 ug/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 BM of 481 mg/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark			Mean	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Data period for Cadmium Trend	Median Concentration 2012 (µg/L)	Trend (% year) (p-value) - colour codes defined above	Selenium (ton/year)	% of Total Tributary Load	Nitrate (ton/year)	% of Total Tributary Load	Sulphate (ton/year)	% of Total Tributary Load									
Grace Creek	LC_GRCK (P), 2004, 2008-2015 [E288275]	100	2.16 [0 of 38]	2.01	0.0365 [0 of 38]	0.0443	47.9 [0 of 38]	45.1	0.01 [0 of 38]	0.0101	2008-2012	1.9	-3.9 (0.0173)	0.03	NT	46.20	NT	2008-2012	0.01	-11.6 (0.0016)	ND	ND	ND	ND	ND	ND	ND	ND	0.31	ND	ND	ND	ND	ND		
Line Creek	LC_LC3	43	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.51	8.0	2.0	0.5	0.6	0.5			
	LIDSL	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.00	49	32	0.6	0.7	0.5				
	LC_LC4 (P), 1990-2015 [200044]	57	42.5 [61 of 81]	38.5	7.34 [18 of 81]	7.5	163 [0 of 81]	145	0.0597 [0 of 80]	0.0929	2004-2012	33	6.3 (<0.0001)	5.65	10.9 (<0.0001)	113.00	5.6 (<0.0001)	2006-2012	0.17	NT	3	18	436	15	9,306	16	ND	ND	ND	ND	0.00	72	28	0.8	1.0	0.7

ND - No Data; NT - No trend

^a Based on monitoring data and site-specific targets established in EVWQP for nitrate, sulphate, cadmium, and selenium (see Appendices A and D in Windward et al. 2014).

^b EPT - Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies).

^c HQ - Hazard Quotient computed as a ratio of tissue concentration to relevant benchmark (Windward 2014). Values above 1 indicate potential risk

^d Quarterly monitoring data will be summarized in December 2015 version for locations monitored quarterly under Permit 107517

^e Data for this location may be available but were not assessed by Windward et al. (2014)

Table 4.3: Aquatic habitat of Lower Fording River (MU2). Data sources are described in Section 2.0.

Tributary Name	Annual Average Daily Flow (m³/s)	Ktunaxa traditional uses and cultural values	Stream Length based on Province's CWB Layer										Fish and Fish Habitat																Riparian Habitat within Catchment Area		Amphibian census location, date (and any observed species-at-risk)	Bird census location, date (and any observed species-at-risk)
			Total (km) (1980s plus Updates)	Currently Connected to Main Stem		Currently Fragmented				Permanently Altered or Destroyed		Fish Species Present	Unique Fish Habitat Features (Yes/No/Unknown)	Barrier(s) Present (excludes infills)	Total Fish Habitat (1980s plus Updates)		Total Non-fish Bearing (1980s plus Updates)		Fish Habitat Currently Connected to Main Stem		Fish Habitat Currently Fragmented				Fish Habitat Permanently Altered or Destroyed							
						Potentially Reversible		Permanently Isolated													Potentially Reversible		Permanently Isolated									
				km	%	km	%	km	%	km	%				km	%	km	%	km	%	km	%	km	%	km	%	km	%	km	%		
Grace Creek	0.18		45.0	14.1	31	31.0	69	0.0	0	0.0	0	WCT	U	A	15.2	34	29.8	66	8	53	7.3	48	0.0	0	0.0	0	2.1	11	ND	ND		
Line Creek	2.06		263.3	109.4	42	0.0	0	112.9	43	41.0	16	WCT, BT, MW	Y	A	52	20	211.3	80	32.2	62	0	0	7.6	15	12.2	23	7.4	5	Settling pond, May 2012.	Line Creek (LI8), June 2012 (Bald eagle).		

ND - No Data; ID - Insufficient Data; WCT - westslope cutthroat trout (Blue-list conservation status); BT - bull trout (Blue-list conservation status); MW - mountain whitefish.



Photo 4.1: Typical riffle-glide morphology in Grace Creek (Teck 2011b).

a pair of culverts about 33 m long, with an outflow height of 0.75 m. The culverts at both the road and rail crossings are considered to be barriers to upstream fish movement (Teck 2011b).

No species other than westslope cutthroat trout have been captured or observed in Grace Creek (Teck 2011b). In June 2004, westslope cutthroat trout were captured downstream from the rail crossing culvert, where there is suitable habitat for all life-stages during the open water season. However, no fish were captured at two areas sampled upstream from the culverts even though a survey in October 2010 confirmed that there is suitable habitat for westslope cutthroat trout (Teck 2011b).

4.2 Line Creek

About 14% of the total catchment area of Line Creek has been disturbed by mining activities (Table 4.1). Mining is currently occurring within the permit boundary in the Line Creek catchment area and will continue into the near and far future. Of the total stream length of 263 km, 42% is connected to the main stem whereas the remainder is

fragmented or has been lost (Table 4.3). The catchment has also been disturbed by forestry activities and roads (Table 4.1).

A rock drain associated with LCO is located about 9 km upstream from the Fording River and covers about 3.8 km of Line Creek (Photo 4.2). West Line Creek, which flows into Line Creek near the downstream end of the rock drain, has been heavily influenced by mining and Teck recently constructed an AWTF to reduce selenium and nitrate loads from West Line Creek into Line Creek. Near the site of the AWTF and downstream from the AWTF outfall, South Line Creek (which is not mine-influenced) also flows into Line Creek. Mine-related pits and spoils are situated in the Line Creek watershed primarily upstream from the confluence with South Line Creek and coal is transported to a processing plant near the Fording River by a conveyor that parallels Line Creek (Arnett and Robinson 2010). For about 2.5 km upstream from the rock drain, Line Creek flows through historical and active mining areas, as well as through past forestry cutblocks. Farther upstream, the headwaters of Line Creek and Tornado Creek each flow for about 4.5 km through areas that have been influenced historically by forestry but not mining before joining together at the upstream end of the mine.



Photo 4.2: Downstream end of the rock drain in Line Creek (Minnow unpublished).

Water quality in Line Creek downstream from LCO was characterized as “poor” based on Windward et al. (2014), and increasing trends in concentrations of selenium, nitrate and sulphate were reported (Zadjlik and Minnow 2013; Table 4.2). The benthic invertebrate community was characterized as “fair” while tissue selenium concentrations were characterized as “good” (Windward et al. 2014). No calcite was observed.

Downstream from LCO, Line Creek flows for about 9 km to its confluence with the Fording River. The creek is low gradient with riffle/glide habitat, except for the portion that flows through a canyon where the habitat includes cascade/pool segments and two falls (Teck 2011b). One of the falls was altered in 1989 to improve fish passage and thereafter neither was considered a barrier to fish movement (Teck 2011b). There were two landslides in the canyon section in June 2015 that blocked the creek, but removal of the slide material and restoration of the creek has been completed.

Numerous surveys of fish and fish habitat in the Line Creek watershed have been conducted over the past 30 years and continue to be completed (Allan 1987; Allan 1991; Allan 1993; Allan 1994; Allan 1995; Goltz 1997; Goltz 1998; Goltz 1999; Allan 2000; Allan 2001; Robinson and Wright 2005; Berdusco et al. 2007; Berdusco and Arnett 2008; Arnett and Berdusco 2009). Line Creek is the only tributary of the Fording River where bull trout and mountain whitefish, as well as westslope cutthroat trout have been reported (Teck 2011b). It is considered a regionally important stream for bull trout spawning, which likely relates to relatively high proportion of groundwater contributing to total streamflow, providing stable thermal conditions throughout the incubation period (Robinson et al. 2014). Adult bull trout overwinter in the Elk River and migrate into Line Creek to spawn (in late September), whereas juveniles rear in Line Creek for several years before migrating out to the Elk River (Teck 2011b). Adult whitefish move into Line Creek from the Fording and Elk rivers to feed in the spring but move out in the fall. The westslope cutthroat trout population in Line Creek is a resident population with all life stages occurring throughout the year. Bull trout and cutthroat trout have also been observed in the lower reaches of South Line Creek. Upstream from the Line Creek rock drain, fish salvages have been conducted periodically since 2004 to relocate bull trout and westslope cutthroat trout to habitats downstream from the rock drain, but it is likely that remnant populations of both bull trout and westslope cutthroat trout remain (Teck 2011b).

Following a comprehensive sampling program in Line Creek after three consecutive years (2007-2009) and comparisons to historical data, Arnett and Robinson (2010) reported that “the absence of pre-mining fish population and spawning data precluded any comparisons of pre- to post-mining conditions. However, based on 15 years of fish population estimate

data and 20 years of bull trout spawning data, it was inferred that while fish populations and bull trout spawning seem to be variable between years, the changes observed appear to be fluctuating independently of mining activity and were more strongly correlated to environmental variables (e.g., June and September discharge, bull trout escapement, water temperature)". The same authors also concluded that water flow (i.e., dilution of mine-related constituents) and sediment loads from South Line Creek may be ameliorating effects of mining on biota in lower Line Creek (Arnett and Robinson 2010).

No amphibians were observed in a survey completed near the settling ponds (Minnow 2014a). A breeding bird survey completed on Line Creek downstream from LCO identified, bald eagle, a species at risk (Minnow 2014a; Table 4.3).

5.0 TRIBUTARIES TO UPPER ELK RIVER (MU3)

Tributaries in the Upper Elk River Watershed (MU3) are shown in Figure 5.1, and include:


- Branch F Creek;
- Wolf Creek;
- Willow Creek;
- Wade Creek;
- Cougar Creek;
- No-Name Creek and Unnamed Tributary 14;
- Mickelson Creek;
- Leask Creek;
- Wolfram Creek; and
- Thompson Creek.

Matrices for these tributaries provide data related to watershed area and disturbances (Table 5.1), environmental quality descriptors (Table 5.2) and fish and aquatic habitat (Table 5.3). An orthophoto figure for each tributary is found in Appendix C. Each tributary is briefly described in the following sections.

5.1 Branch F Creek

Branch F Creek is located on the west side of GHO about 14 km north of Elkford, BC, along the Round Prairie Forest Service Road (FSR). There is currently no influence from mining, but there are expected to be mine-related disturbances within the catchment in the future as part of the Cougar Pit Expansion Project (Table 5.1). All of the total stream length of 6 km is fragmented (Table 5.3). The catchment has also been disturbed by forestry activities and roads (Table 5.1).

Water quality was characterized as “good” based on Windward et al. (2014), but the authors did not report any biological or calcite data for this stream (Table 5.2). No water quality trends were reported for Branch F Creek (Zadjlik and Minnow 2013).



MAP INFORMATION
Datum: NAD 1983 Map Projection: Zone 11N
Data Source: Department of Natural Resources Canada. All rights reserved.
Creation Date: Sept 2015

Legend

Water Flow Direction	Waterbody
WaterNetwork	Streams
Upper Elk River	Mine Development Areas
Watershed Boundary	Teck Coal Mine Operation
Falls	Provincial Boundary
Community	

**Figure 5.1. DRAFT
Upper Elk River
Watershed Boundary**

Teck

1:125,000

Table 5.1: Watershed area and descriptors associated with tributaries to Upper Elk River (MU3). Data sources are described in Section 2.0.

Tributary Name	Catchment Area				Stressors																		Mine Alterations/Disturbances/Planned Mitigation			
	Total Catchment Area from Provincial Corporate Watershed Base Dataset (km ²)	Total Catchment Area as of September 2015 (km ²)	% Change in Total Catchment Area	Teck Ownership Rights: % of Catchment Area and Type (Coal, Surface, Timber)	Mine-Related Disturbances in Catchment as of September 2015										Forestry Disturbances in Catchment				Road Disturbances in Catchment				Additional Planned Near/Far Future Mining Disturbances	Planned Mitigation/Fish Habitat Offsetting		
					Timing of Disturbance: Historical (H), Current (C), Near Future (NF), Far Future (FF) or Not Expected (N)	Total Footprint Area Currently Disturbed (excludes reclaimed lands)		Reclaimed Land Area		Pit(s) (yes/no)	Waste Rock Storage (yes/no)	Rock Drain(s) (yes/no)	Channel Re-alignment or Diversion (yes/no)	Settlement Ponds/Catch Basins (number)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (km ²)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (%)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (km ²)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (%)	Road Density per km ²	Stream Crossings per km	Roads Within 100 m of a Stream	% Roads on Slopes Greater than 60%				
						km ²	%	km ²	%																	
Branch F Creek	1.3	1.3	0	8% Surface and Coal, 54% Surface and Coal and Timber	NF, FF	0.0	0	0.0	0	Y	N	N	N	0	0.4	31	0.0	0	1.2	1.5	0.8	0.0	There is near and far future disturbances from the potential Phase 8 development.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation, however, potentially a settlement pond may be needed in the future.		
Wolf Creek	1.0	1.0	0	10% Surface and Coal, 40% Surface and Coal and Timber	NF, FF	0.0	0	0.0	0	Y	N	N	N	1	0.4	40	0.0	0	0.3	1.0	0.3	0.0	There is near and far future disturbances from the proposed CPX project.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is a settlement pond being built in 2016.		
Willow Creek (Including Willow South, Willow North and Wilde)	2.6	3.1	19	77% Surface and Coal, 40% Surface and Coal and Timber	H, C, 'NF, FF	0.1	4	0.0	0	Y	N	N	Y	2	1.4	54	0.0	0	1.1	3.2	0.6	0.1	Mining currently occurring within the permit boundary will continue into the near and far future as part of the CPX project.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, one settlement pond currently exists and another settlement pond being built in 2016.		
Wade Creek	0.7	0.6	-14	100% Surface and Coal	H, C, NF, FF	0.2	29	0.0	0	Y	N	N	Y	1	0.2	29	0.0	0	0.2	2.0	0.2	0.0	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation.		
Cougar Creek	1.8	1.6	-11	89% Surface and Coal	H, C, NF, FF	0.3	17	0.0	0	Y	N	N	N	0	0.8	44	0.3	17	0.9	1.3	0.4	0.0	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation.		
No-Name Creek (and Unnamed Tributary 14)	1.1	1.1	0	55% Surface and Coal and Timber	N	0.0	0	0.0	0	N	N	N	N	0	0.4	36	0.2	18	2.5	3.6	1.5	0.0	NA	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation.		
Mickelson Creek	3.0	2.6	-13	93% Surface and Coal and Timber	H, C, NF, FF	0.7	24	0.0	0	Y	Y	Y	Y	2	0.7	23	0.3	10	2.7	3.0	1.0	0.0	Mining currently occurring within the permit boundary will continue into the near and far future as part of the CPX project.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation. Water is being pumped into the Mickelson Ponds. Water pumping mitigation is being implemented.		
Leask Creek	1.8	3.7	106	94% Surface and Coal	H, C, NF, FF	1.5	83	0.0	0	N	Y	Y	Y	1	0.1	6	0.1	6	3.3	6.7	2.0	0.0	Mining currently occurring within the permit boundary will continue into the near and far future as part of the CPX project.	Fish barrier installation needed on pond outlet to ensure no fish connection. No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation. Control structure has been installed to prevent debris slides. Water is being pumped into the Leask Ponds. Water pumping mitigation is being implemented.		
Wolfram Creek	3.8	5.6	47	92% Surface and Coal	H, C, NF, FF	3.2	84	0.0	0	N	Y	Y	Y	1	0.5	13	0.3	8	3.0	1.7	0.7	0.0	Mining currently occurring within the permit boundary will continue into the near and far future as part of the CPX project.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation. Control structure has been built to prevent debris slides. Water is being pumped into the Wolfram Ponds. Water pumping mitigation is being implemented.		
Thompson Creek	11.5	10.5	-9	81% Surface and Coal	H, C, NF, FF	4.2	37	1.0	9	Y	Y	Y	Y	3	2.9	25	0.9	8	3.0	1.2	0.4	0.1	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned . From a regional water quality management perspective, there is no planned mitigation.		

NA - Not Applicable

Table 5.2: Environmental quality descriptors for Upper Elk River (MU3). Data sources are described in Section 2.0.

Tributary Name	Treck Surface Water Quality Monitoring Station (Code, Permitted-P or Non-Permitted-NP, Years Data Available, and [EMS# if applicable])	Surface Water Chemistry (2011-2013) (from Windward et al. 2014) ^a									Median Concentrations and Trends Blue- and orange-highlighted cells indicate decreasing and increasing trends, respectively (from Zadjlik and Minnow 2013).									Total Loads (2012) (from Zadjlik and Minnow 2013) Bold load values are associated with 9+ months of data. Non- bold values should be interpreted with caution.						Surface Water Acute Toxicity (2015) (Q1-Q4)		Sediment Quality (2013) (Minnow 2014b)		Calcite (2013) (from Windward et al. 2014)	Benthic Invertebrate		Max SQ ^c for Selenium in Periphyton or Invertebrate Tissue (mean SQ in brackets if max SQ is >1) (Windward et al. 2014)			
		Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Dissolved Cadmium (ug/L) *Not included in WQI		Data Period for Selenium, Nitrate, and Sulphate Trends	Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Cadmium (mg/L)		Community Structure (2012) (Windward et al. 2014)												Colour categories explained in Table 2.4 and text					
		Median [# Samples >Level 1 BM 19 ug/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 BM of 481 mg/L out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median [# Samples >Level 1 hardness-dependent BM out of Total # Samples] - grey shade indicates more than 10% of samples exceeded benchmark	Mean		Water Quality Index (WQI) (see Section 2.0 and Table 2.4 for explanation about how WQI was computed and definitions for colour categories)	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Data period for Cadmium Trend	Median Concentration 2012 (µg/L)	Trend (% year) (p-value) - colour codes defined above	Selenium (ton/year)	% of Total Tributary Load	Nitrate (ton/year)	% of Total Tributary Load	Sulphate (ton/year)	% of Total Tributary Load	Rainbow trout (% Mortality)	Daphnia magna (% Mortality)	Chemistry	Toxicity Survival and growth of <i>Hyalella azteca</i> and <i>Chironomus riparius</i>	Calcite Index	% EPT ^b	% Ephemeroptera	Direct Effects to Invertebrates (periphyton consumption)	Fish Dietary Exposure (Invertebrate Consumption)	Bird Dietary Exposure (Invertebrate Consumption)
Branch F Creek	GH_BR_F (P), 2012-2015 [E287437]	100	0.595 [0 of 6]	0.597	0.198 [0 of 6]	0.215	3.79 [0 of 6]	4.2	0.022 [0 of 6]	0.0253	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND		
Wolf Creek	GH_Wolf (P), 2009-2015 [E287436]	100	0.75 [0 of 32]	0.739	0.136 [0 of 32]	0.356	12.7 [0 of 33]	14.8	0.0102 [0 of 21]	0.0104	2009- 2012	0.64	NT	0.08	NT	17.00	NT	2009- 2012	0.02	NT	0.0	0.0	0.4	0.0	11	0.0	ND	ND	ND	ND	ND	ND	ND	ND	ND	
Willow Creek (Including Willow South, Willow North and Wilde)	GH_Willow (P), 2009-2015 [E287434]	100	0.97 [0 of 55]	1.19	0.0938 [0 of 55]	0.376	10.8 [0 of 56]	17.4	0.0123 [0 of 31]	0.0127	2009- 2012	0.89	-8.8 (0.0318)	0.12	NT	19.98	19.9 (0.0037)	2009- 2012	0.02	NT	0.0	0.0	0.7	0.0	26	0.0	ND	ND	ND	ND	ND	ND	ND	ND	ND	
Wade Creek	GH_Wade (P), 2009-2015 [E287433]	100	1.26 [0 of 32]	1.26	0.655 [0 of 32]	0.664	20.6 [0 of 33]	18.7	0.021 [0 of 21]	0.0198	2009- 2012	1.2	NT	0.47	24.2 (0.0006)	20.80	NT	2009- 2012	0.03	NT	0.0	0.0	1	0.0	13	0.0	0, 0, 0, 0	73, 7, 0, 0	ND	ND	ND	ND	ND	ND	ND	
Cougar Creek	GH_Cougar (P), 2009-2015 [E287432]	100	0.55 [0 of 17]	0.574	0.0877 [0 of 17]	0.124	10.9 [0 of 18]	10.7	0.0305 [0 of 10]	0.0322	2009- 2012	0.55	NT	0.07	NT	11.00	-6.3 (0.0478)	2009- 2012	0.05	NT	0.0	0.0	0.1	0.0	4	0.0	ND, ND, ND, NS	ND, ND, ND, NS	ND	ND	ND	ND	ND	ND	ND	
No-Name Creek	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND		
Mickelson Creek	GH_MC1 (P), 1993-2015 [0200388]	100	1.27 [0 of 36]	1.72	0.165 [0 of 36]	0.532	42.1 [0 of 37]	48.3	0.03 [0 of 23]	0.0293	2005- 2012	0.87	NT	0.19	NT	71.10	4 (0.0073)	2009- 2012	0.05	NT	0.0	0.0	1	0.0	89	0.2	ND, ND, ND, 0	ND, ND, ND, 0	ND	ND	ND	ND	ND	ND	ND	
Leask Creek	GH_LC2 (P), 2005-2015 [E257796 is GH_LC1-to ground]	27	31.1 [20 of 39]	33.9	22.8 [34 of 34]	25.4	146 [0 of 35]	170	0.0173 [0 of 24]	0.0219	2005- 2012	38	27.2 (<0.0001)	29.05	24.7 (<0.0001)	227.00	16.5 (<0.0001)	2008- 2012	0.06	23 (0.0376)	0.0	0.3	22	0.7	168	0.3	ND	ND	ND	ND	ND	ND	ND	ND	ND	
Wolfram Creek	GH_WC2 (P), 2005- 2015 [E257795 is GH_WC1- to ground]	16	33 [33 of 40]	38.5	21.4 [40 of 40]	21.8	336 [8 of 42]	346	0.011 [0 of 28]	0.0811	2005- 2012	44	25.9 (<0.0001)	17.95	29.7 (<0.0001)	399.00	16.1 (<0.0001)	2008- 2012	0.29	25.5 (0.0633)	0.1	0.8	43	1	694	1	NA	NA	ND	ND	0.27	12	1.9	1.3 (0.7)	1.5 (0.8)	1.1 (0.6)
Thompson Creek	GH_TC1 (P), 2006-2015 [E102714]	12	82.4 [43 of 43]	84.7	8.81 [42 of 43]	10.4	446 [20 of 46]	450	0.022 [0 of 29]	0.026	2006- 2012	116	34 (<0.0001)	12.60	44.2 (<0.0001)	509.00	19.8 (<0.0001)	2008- 2012	0.04	NT	0.3	2	32	1	1,274	2	0, 0, 0, 0	0, 0, 0, 0	ND	ND	ND	ND	ND	ND	ND	
	GH_TC2 (P), 1994-2015 [E207436]	13	ND	ND	ND	ND	ND	ND	ND	ND	2004- 2012	97	19.7 (<0.0001)	13.10	23.2 (<0.0001)	498.00	16.8 (<0.0001)	2008- 2012	0.03	NT	ND	ND	ND	ND	ND	ND	0, 0, 0, 0	0, 0, 0, 0	ND	ND	ND	68	1.1	2.7 (1.6)	3.2 (1.9)	2.4 (1.4)

ND - No Data; NS - No Sample; NT - No trend

^a Based on monitoring data and site-specific targets established in EVWQP for nitrate, sulphate, cadmium, and selenium (see Appendices A and D in Windward et al. 2014).

^b EPT - Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies).

^c HQ - Hazard Quotient computed as a ratio of tissue concentration to relevant benchmark (Windward 2014). Values above 1 indicate potential risk.

^d Quarterly monitoring data will be summarized in December 2015 version for locations monitored quarterly under Permit 107517.

Table 5.3: Aquatic habitat of Upper Elk River (MU3). Data sources are described in Section 2.0.

Tributary Name	Annual Average Daily Flow (m³/s)	Ktunaxa traditional uses and cultural values	Stream Length based on Province's CWB Layer									Fish and Fish Habitat																Riparian Habitat within Catchment Area		Amphibian census location, date (and any observed species-at-risk)	Bird census location, date (and any observed species-at-risk)
			Total (km) (1980s plus Updates)	Currently Connected to Main Stem		Currently Fragmented				Permanently Altered or Destroyed		Fish Species Present	Unique Fish Habitat Features (Yes/No/Unknown)	Barrier(s) Present (excludes infills)	Total Fish Habitat (1980s plus Updates)		Total Non-fish Bearing (1980s plus Updates)		Fish Habitat Currently Connected to Main Stem		Fish Habitat Currently Fragmented				Fish Habitat Permanently Altered or Destroyed						
						Potentially Reversible		Permanently Isolated													Potentially Reversible		Permanently Isolated								
				km	%	km	%	km	%	km	%				km	%	km	%	km	%	km	%	km	%	km	%	km	%	km	%	km²
Branch F Creek	<0.011 ^a		6.0	0.0	0	6.0	100	0.0	0	0.0	0	None	No	N	0.0	0	6.0	100	0.0	0	0.0	0	0.0	0	0.0	0	0.2	15	ND	ND	
Wolf Creek	<0.015 ^a		5.6	0.0	0	5.6	100	0.0	0	0.0	0	None	No	N	0.0	0	5.6	100	0.0	0	0.0	0	0.0	0	0.0	0	0.2	20	ND	ND	
Willow Creek (Including Willow South, Willow North and Wilde)	<0.023 ^a		12.8	2.4	19	9.7	76	0.0	0	0.7	5	EB present in Reach 1 in Sept 2013.	No	A, N	0.0 (10m section at mouth)	0	12.8	100	0.0	0	0.0	0	0.0	0	0.0	0	0.5	19	ND	Varied locations, July 2013 (olive-sided flycatcher)	
Wade Creek	<0.007 ^a		3.6	2.3	64	0.0	0	0.0	0	1.3	36	None	No	A, N	0.0	0	3.6	100	0.0	0	0.0	0	0.0	0	0.0	0	0.1	14	Upper watershed, July 2014 (long toed salamander)	Between Wade and Cougar, July 2013 (olive-sided flycatcher)	
Cougar Creek	<0.012 ^a		7.7	6.1	79	0.0	0	0.0	0	1.6	21	None	No	A, N	0.5	6	7.2	94	0.5	100	0.0	0	0.0	0	0.0	0	0.3	17	ND		
No-Name Creek	ND		2.9	2.9	100	0.0	0	0.0	0	0.0	0	None	No	0	0.0	0	2.9	100	0.0	0	0.0	0	0.0	0	0.0	0	0.2	18	ND	ND	
Mickelson Creek	<0.029 ^a		9.3	0.0	0	0.9	10	4.3	46	4.1	44	None	No	A	0.0	0	9.3	100	0.0	0	0.0	0	0.0	0	0.0	0	0.3	10	Upper watershed, July 2014 (western toad)	Varied locations, July 2013 (olive-sided flycatcher)	
Leask Creek	0.023		9.0	0.2	2	0.0	0	2.2	24	6.6	73	None	No	A	0.0	0	9.0	100	0.0	0	0.0	0	0.0	0	0.0	0	0.2	11	Leask Creek Settling Pond (LEPD), May 2012.	Leask Creek Settling Pond (LEPD), June 2012.	
Wolfram Creek	0.039		11.3	0.0	0	1.9	17	0.2	2	9.2	81	None	No	A	0.0	0	11.3	100	0.0	0	0.0	0	0.0	0	0.0	0	0.2	5	Wolfram Settling Pond (WOPD), May 2012.	Wolfram Settling Pond (WOPD), June 2012.	
Thompson Creek	0.091		21.2	2.8	13	5.9	28	0.0	0	12.5	59	WCT, BT, EB	U	A, N	5.4	25	15.8	75	2.3	43	3.0	56	0.0	0	0.1	2	0.8	7	Varied locations, 2012, 2014 (western toad)	Varied locations, 2012, 2013 (olive-sided flycatcher)	

ND - no data; WCT - westslope cutthroat trout (Blue-list conservation status); BT - bull trout (Blue-list conservation status); EB - eastern brook trout.

^a Available data set is missing flows for some low-flow months.

In September 2013, the creek was characterized as dry from the Elk River to the FSR (Lotic 2015b; Photo 5.1). Additional visits in February 2014, and July 2014 again found the channel dry, so no biological sampling occurred. Branch F Creek has an average gradient of 15.3% and an average channel width of 0.80 m, and there is a gradient barrier (slope greater than 30% for 30 m) located at its confluence with the Elk River (Lotic 2015b). The stream is considered non-fish bearing (Lotic 2015b).



Photo 5.1: Dry creek bed in Branch F Creek (Lotic 2015b).

5.2 Wolf Creek

There is currently no influence from mining within the catchment, but there are expected to be mine-related disturbances in the future as part of the Cougar Pit Expansion Project (Table 5.1). All of the total stream length of 5.6 km is fragmented (Table 5.3). The catchment has also been disturbed by forestry activities and roads (Table 5.1).

Water quality was characterized as “good” based on Windward et al. (2014), but the authors did not report any biological or calcite data for this stream (Table 5.2). No trends

were observed for mine-related water quality constituents (Zadjlik and Minnow 2013; Table 5.2).

Wolf Creek has average gradient of 25.7% and average channel width of 1.43 m (Lotic 2015b). In September 2013, most of Wolf Creek was dry, except for some standing water present over a short section downstream from the Round Prairie FSR (Photo 5.2). There was also a dry 5-m-high cascade/fall at the confluence of Wolf Creek and the Elk River that would serve as a barrier to fish movement. Wolf Creek was also dry in February 2014 and was largely subsurface between the FSR and Elk River in July 2014. As a result no biological sampling was conducted. Wolf Creek is considered to be non-fish bearing (Lotic 2015b).



Photo 5.2: Dry creek bed in Wolf Creek (Lotic 2015b).

5.3 Willow Creek

Willow has two branches: Willow Creek North and Willow Creek South. About 4% of the total catchment area of Willow Creek has been disturbed by mining activities (Table 5.1). Mining is currently occurring within the permit boundary will continue into the near and far

future as part of the Cougar Pit Expansion Project. Of the total stream length of 12.8 km, 19% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 5.3). There has also been historical disturbance from forestry and roads (Table 5.1)

Water quality in Willow Creek was characterized as “good” based on Windward et al. (2014), and although a decreasing trend was reported for selenium, nitrate concentrations showed an increasing trend (Zadjlik and Minnow 2013; Table 5.2). Windward et al. (2014) did not report any biological or calcite data for this stream (Table 5.2).

Field surveys of fish and fish habitat were conducted in September 2013, February 2014, and July 2014 (Lotic 2015b). A 2-m-high cascade/waterfall was reported at the Willow Creek and Elk River confluence, which was considered a barrier to upstream fish migration during low flow periods. Upstream from this barrier, was a 10-m section with riffle-pool morphology, average gradient of 3%, average bankfull width of 1.57 m, and dominant substrate of cobble and boulders. At the upstream end of this section was a second barrier (25% gradient for 20 m). Upstream from this second barrier Willow Creek North had step-pool morphology and average gradient of 18.5%. Willow Creek South was dry during two summer surveys and was considered to have insufficient water during the winter to provide fish habitat. The riparian vegetation was mixed forest (Lotic 2015b).

There was only a 10 m fish bearing section in Willow Creek North bordered by the two-metre-high cascade/waterfall on its downstream end (Photo 5.3) and the gradient barrier of 25% for 20 m at its upstream end (Lotic 2015b). The cascade/waterfall at the Willow Creek and Elk River confluence was considered a barrier to upstream fish migration through all but high flow periods. It is likely that the one fish captured in Reach 1 was stranded above the first barrier when the Elk River receded following the spring 2013 record flood. There was no suitable salmonid spawning habitat as the cobble substrate was too large. Pools and glides for holding and feeding areas for adults were lacking. Overall, only the lower 10-m of Willow Creek North has potential for use by sub-adult life-stages of fish during the open-water season, and there is connectivity with the Elk River only during high water flows. The remainder of the creek is non-fish bearing (Lotic 2015b).

Two olive-sided flycatchers were reported in the Willow Creek basin in July 2013 (Matrix Solutions 2015).



Photo 5.3: Two-meter-high waterfall at the confluence of Willow Creek with the Elk River (Lotic 2015b).

5.4 Wade Creek

About 29% of the total catchment area of Wade Creek has been disturbed by mining activities (Table 5.1). Mining currently occurring within the permit boundary will continue into the near and far future. Of the total stream length of 3.6 km, 64% is connected to the main stem whereas the remainder has been lost (Table 5.3). There has also been historical disturbance from forestry activities and roads (Table 5.1).

Water quality has been characterized as “good” based on Windward et al. (2014), although nitrate concentrations have been increasing over time (Zadjlik and Minnow 2013; Table 5.2). Neither benthic invertebrate community nor calcite data were reported by Windward et al. (2014).

Fish and fish habitat surveys were conducted in Wade Creek in September 2013, February 2014 and July 2014 (Lotic 2015b). It was characterized as having a step-pool morphology, with average gradient of 25% and bankfull width of 1.2 m. The dominant

substrate was cobble and boulders. The riparian vegetation was mixed forest (Lotic 2015b).

There was abundant cover in Wade Creek provided by large woody debris, and moderate cover was provided by small woody debris, boulders and undercut banks (Lotic 2015b). However, a gradient barrier of 48% slope for 30 m was found immediately upstream from the confluence of Wade Creek with Elk River which was determined to be a barrier to upstream fish migration at low flows (Lotic 2015b).

Wade Creek was classified as non-fish bearing following two consecutive years of sampling (September 2013, July 2014) during which no fish were caught or observed (Lotic 2015b). The February 2014 overwintering survey indicated that Wade Creek did not provide fish overwintering habitat (Lotic 2015b).

An olive-sided flycatcher was observed between Wade and Cougar Creeks in July 2013 (Matrix Solutions 2015). A long-toed salamander was also observed near Wade Creek in July 2014 (Matrix Solutions 2015).

5.5 Cougar Creek

About 17% of the total catchment area of Cougar Creek has been disturbed by mining activities (Table 5.1). Mining is currently occurring within the permit boundary and will continue into the near and far future. Of the total stream length of 7.7 km, 79% is connected to the main stem whereas the remainder has been lost (Table 5.3). The catchment has also been disturbed by forestry activities and roads.

Water quality was classified as “good” based on Windward et al. (2014; Table 5.2), but biological and calcite data were not reported by Windward et al. (2014).

Cougar Creek has an average gradient of 24% (Lotic 2015b). There is a gradient barrier of 37% slope for 30 m immediately upstream from its confluence with the Elk River. In September 2013, Cougar Creek was reported as dry from the Elk River up to the FSR (Photo 5.4). It was also dry during surveys in February 2014 and July 2014. Therefore, no biological sampling was completed. The combination of dry conditions and gradient barrier preclude fish presence at any time of year. Cougar Creek is classified as non-fish bearing (Lotic 2015b).

An olive-sided flycatcher was observed between Wade and Cougar Creeks in July 2013 (Matrix Solutions 2015).



Photo 5.4: Dry creek bed in Cougar Creek (Lotic 2015b).

5.6 No-Name Creek and Unnamed Tributary 14

Although they are separate creeks, No Name Creek and Unnamed Tributary 14 are both combined within the same watershed boundary defined in Figure 5.1. This catchment is not currently influenced by mining, nor is it expected to be in the future (Table 5.1). All of the total stream length of 2.9 km is connected to the main stem (Table 5.3). The catchment has also been disturbed by forestry activities and roads (Table 5.1).

Water quality, calcite, and biological data were not reported for either No-Name Creek or Unnamed Creek by Windward et al. (2014; Table 5.2).

In September 2013, No Name Creek was identified as an intermittently dry channel (i.e., water depth less than 0.02 m) from the Elk River up to the Round Prairie FSR and was too shallow for electrofishing (Lotic 2015b). Subsequent surveys found the creek dry in February 2014 almost completely dry in July 2014. Unnamed Tributary had no visible channel when surveyed in October 2013 and only standing water upstream from the road in July 2014. The gradient of both creeks was reported as about 16%. No biological

sampling was completed. No barriers to fish migration were observed, but based on low to no flow the creeks were considered non-fish bearing. There were also no records of fish presence/absence in the provincial fisheries database for either stream (Lotic 2015b).

5.7 Mickelson Creek

About 24% of the total catchment area of Mickelson Creek has been disturbed by mining activities (Table 5.1). Mining is currently occurring within the permit boundary will continue into the near and far future as part of the Cougar Pit Expansion Project. The catchment has also been disturbed by forestry activities and roads (Table 5.1). All of the total stream length of 9.3 km is fragmented or has been lost (Table 5.3). The stream flows into a newly constructed sedimentation pond that has no surface connection to the Elk River (Lotic 2015b). The water from the pond decants to Leask Creek and Leask Pond, which also has no overland connection to the Elk River. The catchment has also been disturbed by forestry activities and roads (Table 5.1, Photo 5.5).



Photo 5.5: Extensive logging in Mickelson Creek (Lotic 2015b).

Water quality was characterized as “good” based on Windward et al. (2014), although sulphate concentrations have been increasing slightly over time (4% per year; Table 5.2). Calcite, benthic invertebrate community, and tissue selenium data were not reported by Windward et al. (2014).

A survey conducted September 2013 characterized the stream as having a step-pool morphology, with average gradient of 10.6% (Lotic 2015b). The dominant substrate was cobble and gravel. The average bankfull width in September 2013 was 0.97 m. Calcite was absent or present in low concentrations near the Round Prairie FSR (CI range: 0.00 - 0.02). The riparian vegetation was mature coniferous forest (Lotic 2015b).

Mickelson Creek provided trace amounts of cover for fish with small woody debris and overhanging vegetation (Lotic 2015b). Shallow depths are not suitable salmonid spawning and the sedimentation pond that Mickelson Creek flows into is sometimes dry. A survey in 1979, prior to construction of the settling pond, also reported seasonal channel dryness (BC Research 1981). A 1-m-high waterfall created by a man-made weir was reported about 15 m upstream from the Round Prairie FSR, and is considered a permanent migratory fish barrier (Lotic 2015b; Photo 5.6). In 2014 the area was logged, and the flow had been greatly reduced from 2013 levels, with sections of subsurface flow. There were no records of fish presence/absence in the provincial fisheries database for Mickelson Creek (Lotic 2015b). No fish were captured or observed during sampling in September 2013, and no fish inventory sampling could be completed for Mickelson Creek in 2014, due to low water. Mickelson Creek was classified as non-fish bearing due to the lack of fish caught during September 2013 sampling and its lack of connectivity to the Elk River (Lotic 2015b).

Olive-sided flycatchers (July 2013) and western toads (July 2014) were observed in the Mickelson Creek catchment (Matrix Solutions 2015).

5.8 Leask Creek

About 83% of the total catchment area of Leask Creek has been disturbed by mining activities (Table 5.1). Mining currently occurring within the permit boundary will continue into the near and far future as part of the Cougar Pit Expansion project. Of the total stream length of 9 km, 2% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 5.3). The catchment has also been disturbed by forestry activities and roads (Table 5.1).



Photo 5.6: One metre tall waterfall on Mickelson Creek upstream from the Round Prairie Forest Service Road (Lotic 2015b).

The lower portion of Leask Creek consists of two man-made, rectangular sedimentation ponds (2,784 and 2,002 m²) that lack connectivity to the Elk River (Photo 5.7). The downstream pond is dry for most of the year and the water depth of the upper pond fluctuates dramatically (Lotic 2015b). There were no macrophytes observed growing in the ponds. There is a culvert crossing between the two ponds at the Round Prairie FSR, and there is another culvert upstream from the second pond.

Water quality was categorized as “poor” based on Windward et al. (2014) and increasing trends in concentrations were reported for selenium, nitrate, sulphate, and cadmium (Zadjick and Minnow 2013; Table 5.2). No calcite index was reported for Leask Creek by Windward et al. (2014), but Lotic (2015b) reported low amounts of calcite in the second pond (CI range: 0.0 - 0.51).



Photo 5.7: Downstream sedimentation pond at Leask Creek (Lotic 2015b).

Active water treatment is scheduled to begin in 2026, treating water from GHO's West Spoil (influencing Thompson, Leask, and Wolfram Creeks) as well as Greenhills Creek (Teck 2014). Treated effluent will discharge to Thompson Creek.

Upstream from the ponds the creek consisted of a step-pool morphology with average gradient of 12%, and average bankfull width of 2.92 m (Lotic 2015b). The dominant substrate was cobble and gravel with and moderate calcite was detected (CI range: 0.51 - 1.8; Lotic 2015b).

Upper Leask Creek provides trace amounts of cover by small and large woody debris, undercut banks, and overhanging vegetation (Lotic 2015b). However, the culverts associated with the ponds may be barriers to fish movement in low flow (if fish were present). There were no records of fish capture in the provincial fisheries database for Leask Creek. Fish surveys were conducted in the upper portion of Leask Creek in July and September 2012, but no fish were captured. As a result, Leask Creek was classified as non-fish bearing (Lotic 2015b).

No species at risk were observed during breeding amphibian and bird surveys at the Leask Creek setting ponds in 2012 (Minnow 2014a; Table 5.3).

5.9 Wolfram Creek

Wolfram Creek branches into North Wolfram Creek and South Wolfram Creek about 290 m upstream from the Round Prairie Forestry Side Road (FSR).

About 84% of the total catchment area of Wolfram Creek has been disturbed by mining activities (Table 5.1). Mining is currently occurring within the permit boundary and will continue into the near and far future as part of the Cougar Pit Expansion project. All of the total stream length of 11.3 km is fragmented or has been lost (Table 5.3). The catchment has also been disturbed by forestry activities and roads (Table 5.1).

Water quality in the lower creek has been categorized as “poor”, and increasing trends in concentrations have been reported for selenium, nitrate, sulphate, and cadmium (Zadjlik and Minnow 2013; Table 5.2). The benthic invertebrate community was categorized as poor and tissue selenium concentrations were categorized as fair. Low amounts of calcite have been reported. Active water treatment is scheduled to begin in 2026, treating water from GH0’s West Spoil (influencing Thompson, Leask, and Wolfram Creeks) as well as Greenhills Creek (Teck 2014). Treated effluent will discharge to Thompson Creek.

Fish and habitat surveys were conducted in 1994, and more recently in July and September 2012 (Lotic 2015b). The most downstream section of Wolfram Creek consists of two man-made sedimentation ponds (about 2,958 and 1,843 m²; Photo 5.8). The downstream pond is dry most of the year. The water depth of the ponds fluctuates substantially throughout the year, and there were no macrophytes observed growing in the ponds. The culvert crossing at the FSR may be a barrier to fish movement in low flow (Lotic 2015b). There was no overland connectivity between the ponds and the Elk River. The stream extends upstream for 290 m before splitting into the north and south branches. It has a step-pool morphology and an average gradient of 10.6%. The dominant substrate was cobble and gravel and the average bankfull width was 4.36 m (Lotic 2015b).



Photo 5.8: Upstream view of downstream sedimentation pond on Wolfram Creek (Lotic 2015b).

Due to the infiltration of water to ground in the ponds and lack of surface channel connection with the Elk River¹⁰, the ponds in Wolfram Creek are considered non-fish bearing (Lotic 2015b). A survey conducted in 1979, prior to settling pond construction, also reported seasonal dryness in Wolfram Creek (BC Research 1981). The upper section of the creek has high sediment deposition due to eroding banks, and moderate instream cover from large woody debris and overhanging vegetation. No suitable salmonid spawning habitat was present due to the calcite deposition and fine sediment depositions caused by bank erosion. Deep pools and glides for holding and feeding areas for adults were lacking. No fish were captured or observed in surveys completed in July

¹⁰ There may be overland connection to the Elk River during high flow periods but there is no defined channel, and there is also a culvert at the ponds that is a barrier to fish passage.

and September 2012. Therefore, upper Wolfram Creek is also classified as non-fish bearing (Lotic 2015b).

No species at risk were observed during breeding amphibian and bird surveys at the Leask Creek setting ponds in 2012 (Minnow 2014a; Table 5.3).

5.10 Thompson Creek

Upper (South) Thompson Creek flows northwest before it is joined by North Thompson Creek and then flows west into a side channel of the Elk River. About 37% of the total catchment area of Thompson Creek has been disturbed by mining activities (Table 5.1). Mining currently occurring within the permit boundary will continue into the near and far future. Of the total stream length of 21.2 km, 13% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 5.3). The catchment has also been disturbed by forestry activities and roads (Table 5.1).

Water quality was characterized as “poor” based on Windward et al. (2014), and increasing concentration trends were reported for selenium, nitrate and sulphate (Zadjlik and Minnow 2013; Table 5.2). The benthic invertebrate community and tissue selenium concentrations were also categorized as “poor” (Windward et al. 2014).

Active water treatment is scheduled to begin in 2026, treating water from GHO’s West Spoil (influencing Thompson, Leask, and Wolfram Creeks) as well as Greenhills Creek (Teck 2014). Treated effluent will discharge to Thompson Creek.

Most of Thompson Creek is high gradient and dominated by boulder substrates (RL&L 1994). The lowest portion of the downstream reach has low gradient and subject to intermittent water flows during fall months. Subsurface flow was reported in 1979 300 m downstream from the road crossing in 1979 (BC Research 1981). A series of waterfalls located on North Thompson Creek, just upstream from the confluence with upper Thompson Creek, is a barrier to fish passage. There is also a barrier on Upper Thompson creek 150 m upstream of its confluence with North Thompson Creek. There are three settling ponds (Lower Thompson Creek Pond, Upper Pond and North Pond). The Upper and North ponds are located upstream from the falls that act as a barrier to fish movement. Lower Thompson Creek Pond has three cells constructed within the creek and adjacent low lying area. The system has a bypass channel (RL&L 1994).

Fish species found in Thompson Creek include bull trout, eastern brook trout, and westslope cutthroat trout (RL&L 1994). Bull trout and westslope cutthroat trout were also reported in Thompson Creek during a fish survey in 1979, prior to mine development (BC

Research 1981). The creek contains some suitable habitat for fish rearing, feeding, and spawning (RL&L 1994). Cutthroat trout are present year round, and the presence of adults in spawning condition, and young-of-the-year downstream of the barriers indicates the population may be resident. Fish can move in and out of the Lower Thompson Creek Pond, although the spillway at the outlet may be a partial barrier during low flows (Teck Internal Sources). The largest cell of the pond may provide overwinter habitat (RL&L 1994).


Western toad and olive-sided flycatcher, which are species at risk, were identified during breeding amphibian and bird surveys at the Thompson Creek setting ponds in 2012 (Minnow 2014a; Table 5.3). Surveys completed in 2013 and 2014 reported olive-sided flycatchers, western toads, long-toed salamanders, and a Columbia spotted frog in the catchment of Thompson Creek (Matrix Solutions 2015).

6.0 TRIBUTARIES TO MICHEL CREEK (MU4)

Tributaries in the Michel Creek Watershed (MU4) are shown in Figure 6.1, and include:

- Upstream Michel Creek;
- Corbin Creek;
- Andy Good Creek;
- Leach Creek;
- Carbon Creek;
- Snowslide Creek;
- Wheeler Creek;
- Unnamed Tributary to Michel Creek;
- Fir Creek
- Alexander Creek
- Erickson Creek
- South Pit Creek
- Milligan Creek
- Gate Creek
- Bodie Creek; and
- Aqueduct Creek.

Matrices for these tributaries provide data related to watershed area and disturbances (Table 6.1), environmental quality descriptors (Table 6.2) and fish and aquatic habitat (Table 6.3). An orthophoto figure for each tributary is found in Appendix D. Each tributary is briefly described in the following sections.



MAP INFORMATION
Datum: NAD 1983 Map Projection: Zone 11N
Data Source: Department of Natural Resources Canada. All rights reserved.
Creation Date: Sept 2015

Legend

Water Flow Direction	Waterbody
Michel Creek	Streams
Watershed Boundary	Mine Development Areas
Falls	Teck Coal Mine Operation
Community	Provincial Boundary

Figure 6.1. DRAFT Michel Creek Watershed Boundary

1:150,000

Table 6.1: Watershed area and disturbances associated with tributaries to Michel Creek (MU4). Data sources are described in Section 2.0.

Tributary Name	Catchment Area				Stressors																	Mine Alterations/Disturbances/Planned Mitigation		
	Total Catchment Area from Provincial Corporate Watershed Base Dataset (km ²) (1980s plus updates)	Total Catchment Area as of September 2015 (km ²)	% Change in Total Catchment Area	Teck Ownership Rights: % of Catchment Area and Type (Coal, Surface, Timber)	Mine-Related Disturbances in Catchment as of September 2015										Forestry Disturbances in Catchment				Road Disturbances in Catchment				Additional Planned Near/Far Future Mining Disturbances	Planned Mitigation/Fish Habitat Offsetting
					Timing of Disturbance: Historical (H), Current (C), Near Future (NF), Far Future (FF) or Not Expected (N)	Total Footprint Area Currently Disturbed (excludes reclaimed lands)		Reclaimed Land Area		Pit(s) (yes/no)	Waste Rock Storage (yes/no)	Rock Drain(s) (yes/no)	Channel Re-alignment or Diversion (yes/no)	Settlement Ponds/Catch Basins (number)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (km ²)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (%)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (km ²)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (%)	Road Density per km ²	Stream Crossings per km	Roads Within 100 m of a Stream	% Roads on Slopes Greater than 60%		
						km ²	%	km ²	%															
Upstream Michel Creek (above Corbin Creek confluence)	35.0	33.6	-4	22% Coal, 3% Coal and Surface	H	1.9	5	0.6	2	N	N	N	N	2	0.0	0	0.0	0	1.3	1.2	0.7	0.0	Active mining is expected to be completed by end of 2017 and closure plans will be implemented.	Closure planning will take into consideration fish and fish habitat and water quality.
Corbin Creek	30.4	32.1	6	34% Coal, 14% Surface and Coal	H, C, NF	7.6	25	1.2	4	Y	Y	Y	Y	12	0.0	0	0.0	0	0.5	0.2	0.2	0.0	Active mining is expected to be completed by end of 2017 and closure plans will be implemented.	Closure planning will take into consideration fish and fish habitat and water quality. Corbin Creek has been identified as a is priority stream for potential calcite management under the EVWQP.
Andy Good Creek	34.5	34.4	0	14% Coal	H	0.3	1	0.2	1	N	Y	N	N	0	0.0	0	0.0	0	0.6	1.1	0.4	0.0	NA	Closure planning will take into consideration fish and fish habitat and water quality.
Leach Creek	126.6	126.6	0	10% Coal, 2% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	5.6	4	3.7	3	1.3	1.2	0.5	0.1	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Carbon Creek	10.5	10.5	0	42% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	0.1	1	1.2	11	1.7	2.0	0.8	0.4	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Snowslide Creek	5.6	5.6	0	100% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	0.4	7	0.6	11	3.3	2.0	1.0	0.7	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Wheeler Creek	32.5	32.5	0	1% Coal, 77% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	5.5	17	2.4	7	3.2	2.7	1.2	0.5	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Unnamed to Michel Creek	5.4	5.4	0	24% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	1.6	30	0.3	6	3.6	1.9	1.3	0.0	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Fir Creek	11.6	11.6	0	9% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	5.7	49	0.0	0	2.7	1.0	1.0	0.0	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Alexander Creek	186.4	186.4	0	10% Surface and Coal, 14% Surface and Timber	N	0.0	0	0.0	0	N	N	N	N	0	13.7	7	3.6	2	1.2	0.9	0.5	0.0	NA - Teck Conservation Land	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Erickson Creek	32.5	32.1	-1	90% Surface and Coal, 38% Surface and Timber	H, C, NF, FF	8.4	26	1.4	4	N	Y	Y	N	1	0.9	3	0.0	0	1.4	1.7	0.8	0.3	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, the initial implementation plan indicates Erickson Creek being diverted for AWTF in two phases, Phase 1 by 2020 and Phase 2 by 2024; the initial implementation plan will be reviewed and revised as additional monitoring and modeling work is conducted under the EVWQP. Erickson Creek has been identified as a priority stream for potential calcite management under the EVWQP.
South Pit Creek	0.9	1.6	78	100% Surface and Coal	H, C, NF	0.5	56	0.3	33	Y	Y	Y	Y	2	0.0	0	0.0	0	1.3	2.5	0.5	0.0	Mining currently occurring within the permit boundary will continue into the near future with reclamation.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Milligan Creek	5.6	4.8	-14	100% Surface and Coal	H, C, NF	1.8	32	1.2	21	N	Y	N	Y	2	0.0	0	0.0	0	2.5	1.3	0.4	0.0	Mining currently occurring within the permit boundary will continue into the near future with reclamation.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Gate Creek	5.0	4.4	-12	100% Surface and Coal	H, C, NF, FF	2.8	56	1.6	32	Y	Y	Y	Y	2	0.0	0	0.0	0	5.2	1.8	0.9	0.8	Mining currently occurring within the permit boundary will continue into the near and far future.	Fish salvage and exclusion for Gate Pond and fish-bearing portion of the channel to occur in 2016. No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, the initial implementation plan indicates Gate Creek being diverted for AWTF in Phase 1 by 2020; the initial implementation plan will be reviewed and revised as additional monitoring and modeling work is conducted under the EVWQP. Water is currently being pumped from the Natal Pit to the Gate and Bodie Systems till 2019.
Bodie Creek	10.5	1.4	-87	100% Surface and Coal	H, C, NF, FF	10.3	98	1.0	10	Y	Y	Y	Y	2	0.0	0	0.0	0	4.5	5.0	3.5	0.0	Mining currently occurring within the permit boundary will continue into the near and far future.	Fish salvage and exclusion for Bodie Pond and fish-bearing portion of the channel to occur in 2016. No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, the initial implementation plan indicates Gate Creek being diverted for AWTF in Phase 1 by 2020; the initial implementation plan will be reviewed and revised as additional monitoring and modeling work is conducted under the EVWQP. Water is currently being pumped from the Natal Pit to the Gate and Bodie Systems till 2019.
Aqueduct Creek	7.0	5.7	-19	96% Surface and Coal	H, C, NF, FF	2.3	33	0.6	9	Y	Y	N	Y	1	0.0	0	0.0	0	6.0	3.0	1.2	0.1	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation. Sediment Pond System was built on Aqueduct Creek in 2015. Additional constuction will occur in 2016.

NA - Not Applicable

Table 6.2: Environmental quality descriptors for tributaries to Michel Creek (MU4). Data sources are described in Section 2.0.

Tributary Name	Teck Surface Water Quality Monitoring Station (Code, Permitted-P or Non-Permitted-NP, Years Data Available, and [EMS# if applicable])	Surface Water Chemistry (2011-2013) (from Windward et al. 2014) ^a									Median Concentrations and Trends Blue- and orange-highlighted cells indicate decreasing and increasing trends, respectively (from Zadajlik and Minnow 2013).										Total Loads (2012) (from Zadajlik and Minnow 2013) Bold load values are associated with 9+ months of data. Non-bold values should be interpreted with caution.						Surface Water Acute Toxicity (2015) (Q1-Q4)		Sediment Quality (2013) (Minnow 2014b)		Calcite (2013) (from Windward et al. 2014)	Benthic Invertebrate Community Structure (2012) (Windward et al. 2014)		Max SQ ^c for Selenium in Invertebrate Tissue (mean SQ in brackets if max SQ is >1) (Windward et al. 2014)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
		Water Quality Index (WQI) (see Section 2.0 and Table 2.4 for explanation about how WQI was computed and definitions for colour categories)	Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Dissolved Cadmium (ug/L) *Not included in WQI		Data Period for Selenium, Nitrate, and Sulphate Trends	Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Cadmium (mg/L)			Selenium (ton/year)	% of Total Tributary Load	Nitrate (ton/year)	% of Total Tributary Load	Sulphate (ton/year)	% of Total Tributary Load	Rainbow trout (% Mortality)	Daphnia magna (% Mortality)	Chemistry	Toxicity Survival and growth of <i>Hyalella azteca</i> and <i>Chironomus riparius</i>		Calcite Index	% EP ^b	% Ephemeroptera	Direct Effects to Invertebrates	Fish Dietary Exposure (Invertebrate Consumption)	Bird Dietary Exposure (Invertebrate Consumption)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
			Median (# Samples >Level 1 BM 19 ug/L out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median (# Samples >Level 1 hardness-dependent BM out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median (# Samples >Level 1 BM of 481 mg/L out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median (# Samples >Level 1 hardness-dependent BM out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean		Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Data period for Cadmium Trend	Median Concentration 2012 (µg/L)	Trend (% year) (p-value) - colour codes defined above																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								

ND - No Data; NT - No trend

^a Based on monitoring data and site-specific targets established in EVWQP for nitrate, sulphate, cadmium, and selenium (see Appendices A and D in Windward et al. 2014).

^b EPT - Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies)

^c HQ - Hazard Quotient computed as a ratio of tissue concentration to relevant benchmark (Windward 2014). Values above 1 indicate potential risk

^d Quarterly monitoring data will be summarized in December 2015 version for locations monitored quarterly under Permit 107517

^e Data for this location may be available but were not assessed by Windward et al. (2014).

Table 6.3: Aquatic habitat of Michel Creek tributaries (MU4). Data sources are described in Section 2.0

Tributary Name	Annual Average Daily Flow (m³/s)	Ktunaxa traditional uses and cultural values	Stream Length based on Province's CWB Layer									Fish and Fish Habitat																Riparian Habitat within Catchment Area		Amphibian survey location, date (and any observed species-at-risk)	Bird census location, date (and any observed species-at-risk)
			Total (km) (1980s plus Updates)	Currently Connected to Main Stem		Currently Fragmented				Permanently Altered or Destroyed		Fish Species Present	Unique Fish Habitat Features (Yes/No/Unknown)	Barrier(s) Present (excludes infills)	Total Fish Habitat (1980s plus Updates)		Total Non-fish Bearing (1980s plus Updates)		Fish Habitat Currently Connected to Main Stem		Fish Habitat Currently Fragmented				Fish Habitat Permanently Altered or Destroyed						
						Potentially Reversible		Permanently Isolated													Potentially Reversible		Permanently Isolated								
				km	%	km	%	km	%	km	%				km	%	km	%	km	%	km	%	km	%	km	%	km				
Upstream Michel Creek (above Corbin Creek confluence)	0.27		84.8	80.4	95	1.3	2	0.3	0	2.8	3	WCT, EB	U	0	28.9	34	55.9	66	28.9	100	0.0	0	0.0	0	0.0	0	3.7	11	ND	ND	
Corbin Creek	0.38		63.0	2.3	4	3.5	6	42.5	67	14.8	23	WCT, MW, EB, LSU, LNC	U	A	16.1	26	46.9	74	2.0	12	3.1	19	4.7	29	6.3	39	1.2	4	ND	ND	
Andy Good Creek	ND		75.4	75.4	100	0.0	0	0.0	0	0.0	0	WCT, EB	U	0	13.1	17	62.4	83	13.1	100	0.0	0	0.0	0	0.0	0	2.4	7	ND	ND	
Leach Creek	ND		267.1	267.1	100	0.0	0	0.0	0	0.0	0	WCT, BT, MW, EB	Y	0	67.3	25	199.8	75	67.3	100	0.0	0	0.0	0	0.0	0	11.5	9	ND	ND	
Carbon Creek	0.17		21.1	4.6	22	16.5	78	0.0	0	0.0	0	WCT, EB	Y	N	3.9	18	17.2	82	3.9	100	0.0	0	0.0	0	0.0	0	0.9	9	ND	ND	
Snowslide Creek	0.09		11.1	3.6	32	7.5	68	0.0	0	0.0	0	WCT, EB	Y	N	3.6	32	7.5	68	3.6	100	0.0	0	0.0	0	0.0	0	0.5	9	ND	ND	
Wheeler Creek	0.52		60.8	26.4	43	34.4	57	0.0	0	0.0	0	WCT, BT, EB	U	N	6.5	11	54.3	89	6.5	100	0.0	0	0.0	0	0.0	0	2.8	9	Upper catchment, 2013 (western toad)	ND	
Unnamed to Michel Creek	ND		10.1	0.1	1	9.9	98	0.0	0	0.0	0	WCT	U	N	0.1	1	9.9	98	0.1	100	0.0	0	0.0	0	0.0	0	0.5	9	ND	Upper catchment, 2013 (olive-sided flycatcher)	
Fir Creek	ND		19.1	7.1	37	12.0	63	0.0	0	0.0	0	WCT, EB	Y	N	0.2	1	18.9	99	0.2	100	0.0	0	0.0	0	0.0	0	1.2	10	ND	ND	
Alexander Creek	3.27		309.4	309.4	100	0.0	0	0.0	0	0.0	0	WCT, BT, RB, EB	U	P	72.9	24	236.5	76	72.9	100	0.0	0	0.0	0	0.0	0	12.5	7	ND	Alexander Creek u/s Michel (AL4), June 2012 (olive-sided flycatcher).	
Erickson Creek	0.39		82.4	0.3	0	28.1	34	36.1	44	17.8	22	WCT, BT, EB	No	N	18.7	23	63.6	77	0.3	2	6.7	36	4.5	24	7.3	39	2.2	7	ND	Varied locations, 2013 (olive-sided flycatcher)	
South Pit Creek	0.01		1.3	0.0	0	0.6	46	0.0	0	0.7	54	None	No	A, N	0.0	0	1.3	100	0.0	0	0.0	0	0.0	0	0.0	0	0.1	11	ND	ND	
Milligan Creek	<0.020 ^a		7.2	4.4	61	0.0	0	1.0	14	1.8	25	None	No	A	0.0	0	7.2	100	0.0	0	0.0	0	0.0	0	0.0	0	0.3	5	ND	ND	
Gate Creek	0.03		5.9	0.4	7	0.7	12	0.5	8	4.3	73	WCT, LSU, LNC	No	A	1.2	20	4.7	80	0.4	33	0.2	17	0.0	0	0.6	50	0.2	4	ND	ND	
Bodie Creek	0.01		12.5	0.7	6	0.7	6	0.0	0	11.0	88	WCT, MW, LSU, LNC	No	A	1.2	10	11.3	90	0.7	58	0.3	25	0.0	0	0.2	17	0.1	1	Bodie Creek Settling Pond (BOPD), May 2012.	Bodie Creek Settling Pond (BOPD), June 2012.	
Aqueduct Creek	0.02		6.9	2.2	32	0.9	13	1.6	23	2.1	30	WCT, BT, EB, LSU	U	A	2.5	36	4.3	62	2.0	80	0.0	0	0.0	0	0.5	20	0.3	4	ND	Upper catchment, 2013 (olive-sided flycatcher)	

ND - no data; WCT - westslope cutthroat trout (Blue-list conservation status); BT - bull trout (Blue-list conservation status); MW - mountain whitefish; LSU - longnose sucker; LNC - longnose dace; RB - rainbow trout.

^a Available data set is missing flows for some low-flow months.

6.1 Upstream Michel Creek

A small amount (5%) of the total catchment area of Michel Creek upstream from Corbin Creek has been disturbed by mining activities (Table 6.1). Active mining is expected to be completed by 2017 (Table 6.1). Of the total stream length of 84.8 km, 95% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).

Based on samples collected upstream from CMO influence, water quality and tissue selenium concentrations were characterized as “good” (Windward et al. 2014; Table 6.2). The benthic invertebrate community was categorized as “fair” based on relatively low percentage of Ephemeroptera compared to other reference areas in the region (Table 6.2). No calcite was observed.

A reach surveyed for fish and fish habitat upstream from CMO influence in 2007-2008 had an average channel width from 5.1 m to 10.4 m and wetted width from 2.4 m to 5.9 m (NorthSouth Consultants Inc. 2007, 2008). Gradient within the reach surveyed ranged from 1% to 3%. A survey of a representative 10-m reach in 2012 noted bankfull width of 6 m, gradient of 3%, habitat of riffle (75%) and glide (25%), with dominant substrate of cobble (Minnow 2014). Riparian vegetation consisted of coniferous trees, ferns and grasses, and shrubs (Minnow 2014). Fish sampling in September 2007 and June and August in 2008 captured westslope cutthroat trout and eastern brook trout (NorthSouth Consultants Inc. 2007, 2008).

6.2 Corbin Creek

About 25% of the total catchment area of Corbin Creek has been disturbed by mining activities but active mining is expected to be completed by 2017 (Table 6.1). Of the total stream length of 63 km, 4% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).

Water quality was categorized as “poor” (Windward et al., 2014) and increasing trends in selenium, nitrate, and sulphate concentrations were reported (Zadjlick and Minnow 2013; Table 6.2; Figure 6.2). High levels of calcite have been observed (calcite index: 1.95), and the benthic invertebrate community was characterized as “poor”. Tissue selenium concentrations were characterized as “good” (Windward et al. 2014).




Figure 6.2: Selenium concentration trend over time (2004-2013) in Corbin Creek at the confluence with Michel Creek (CM_CC1).

Golder (2015) reported stream gradient of 2% and channel width of 6.3 m. An assessment of a small section of the creek (30 m) near the confluence with Michel Creek in September 2012 identified habitat of riffles (50%) and glides (50%), with gradient of 1.5% and bankfull width of 6.4 m (Minnow 2014). Substrate was predominantly cobble and riparian vegetation was predominantly ferns and grasses with some shrubs and coniferous trees (Golder 2015a).

Historical fish surveys identified westslope cutthroat trout, bull trout, eastern brook trout, and mountain whitefish in Corbin Creek (Golder 2015a). About 1.5 km upstream from the confluence with Michel Creek is an offline settling pond system that is connected to a man-made channel containing fish. A fish salvage was recently conducted in the man-made channel, relocating about 700 fish, mostly westslope cutthroat trout and eastern brook trout with some longnose sucker and longnose dace (M. Robinson, Lotic Environmental, pers. comm.).

6.3 Andy Good Creek

A small amount (1%) of the total catchment area of Andy Good Creek has been disturbed by mining activities (Table 6.1). All of the total stream length of 75.4 km is connected to the main stem (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality, benthic invertebrate community and tissue selenium concentrations were characterized as “good” (Windward et al. 2014; Table 6.2). No calcite was observed (Table 6.2).

An unspecified stream distance surveyed for fish and fish habitat near the mouth of the creek in August 2010 had bankfull width of 25 m and gradient of 4% (Interior Reforestation 2011). The dominant substrate materials were cobble and gravel. Riparian vegetation was not reported.

Habitat was also documented at a 100-m section of the creek about 1 km upstream from the confluence with Michel Creek in September 2012 (Minnow 2014). The surveyed area consisted of riffle habitat with bankfull width of 14.6 m and gradient of 1.5%. Riparian vegetation included shrubs and coniferous trees, with some grass and ferns (Minnow 2014).

Westslope cutthroat trout and eastern brook trout have been captured in Andy Good Creek (Interior Reforestation 2011). No other information was found at the time of reporting regarding availability of spawning or overwintering habitats for fish, or potential presence of unique or off-channel habitats.

6.4 Leach Creek

Leach Creek is a tributary to Michel Creek that has not been influenced by mining (Table 6.1). All of the total stream length of 267 km is connected to the main stem (Table 6.3). Marten Creek and Bray Creek are tributaries to Leach Creek and have also not been affected by mining. The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality and the benthic invertebrate community were not evaluated by Windward et al. (2014; Table 6.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). Tissue selenium concentrations were categorized as “good”.

Fish and fish habitat surveys were conducted at two areas in the upper and lower portions of Leach Creek, respectively, in 2007 and 2008 (Golder 2015a). An area located 25 m upstream from the confluence with Michel Creek was surveyed again in 2012 (Golder 2015a). Stream morphology of the surveyed area in the upper watershed was characterized as cascade-pool with predominantly glide (40%), riffle (27%), and cascade (27%) habitats and substrate composed of cobble, boulder, and gravel (Golder 2015a; Photo 6.1). Mean bankfull width was 3.2 to 7.4 m (Golder 2015a). The surveyed area in the lower watershed had stream gradient and bankfull width of 1.5% and 18 m, respectively, with habitat dominated by glides (49%), riffles (39%), and pools (12%; Golder 2015a). The dominant substrates were gravel, cobble, and boulder. Riparian vegetation throughout the watershed was mixed mature forest (Golder 2015a).



Photo 6.1: Typical riffle habitat in Leach Creek (Golder 2015).

Westslope cutthroat trout, eastern brook trout, and mountain whitefish have been observed in Leach Creek (Golder 2015a). Leach Creek provides suitable habitat for all life stages of fish during the open-water season. Instream cover for fish includes woody debris (abundant), boulders, pools, and overhanging vegetation. Near the confluence of Michel Creek undercut banks provide additional instream cover. Isolated pockets of spawning gravels suitable for resident trout were present. Undercut banks and pools were considered suitable for overwintering. A pond located about 5 km upstream from Michel Creek contains eastern brook trout that may be resident (Minnow 2003).

6.5 Carbon Creek

Carbon Creek is not currently influenced by mining and is not expected to be affected in the future (Golder 2015a; Table 6.1). Of the total stream length of 21.1 km, 22% is connected to the main stem whereas the remainder is fragmented (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality, the benthic invertebrate community, and calcite levels were not evaluated by Windward et al. (2014; Table 6.2), but are likely comparable to other regional streams

that have not been affected by mining (e.g., “good”). Tissue selenium concentrations were categorized as “good”.

Fish and fish habitat surveys were conducted at two locations on Carbon Creek (lower and middle watershed) in 2008, 2012, and 2013 (Golder 2015a). A surveyed section (102 m) of the creek, located about 730 m upstream from the mouth at Michel Creek, was characterized as cascade-pool form, having a mixture of cascade habitat (82.8%), with glide (8.60%), riffle (6.6%), and pool (2.0%) habitat types (Golder 2015a). Stream gradient was 1.9% with bankfull width of 5.6 m. Another (100-m) section, located about 1.5 km upstream from the confluence with Michel Creek had stream morphology of riffle-pool form, with riffle (70.0%), cascade (17.0%), and glide (13.0%) habitat types. Average stream gradient was higher in this section (5.8%) and bankfull width was 5.0 m. Farther upstream, gradient significantly increased to between 18.3 and 22.3%, and was characterized by a step-pool morphology. Cobble and boulder are the dominant and subdominant substrates throughout the stream, with gravel also being abundant in some areas. Riparian vegetation includes mixed mature forest, coniferous mature forest and shrubs. Off-channel habitat was not observed (Golder 2015a).

The provincial database does not have records of fish historically captured in Carbon Creek (Golder 2015a). A gradient barrier is present mid watershed (about 4 km upstream from Michel Creek), consisting of a 27% gradient bedrock cascade 6 m in length, with a 31% gradient boulder cascade for 15 m slightly farther upstream, that together are suspected to preclude upstream movement of fish (Photo 6.2). No fish were observed upstream from these barriers over multiple surveys. Lower Carbon Creek provides suitable habitat for all life stages of fish during the open-water season. Both adult and juvenile westslope cutthroat trout have been captured or observed, as well as eastern brook trout. A variety of instream and overhead cover is present, such as large and small woody debris, overhanging vegetation, and some deep pools (e.g., 0.4 m). Areas with gravels suitable for spawning are present, and two redds were observed during the spring 2012 spawning survey. The downstream (4 km) fish-bearing portion is considered to provide suitable habitat for all life stages of fish during the open-water season and for overwintering. Spawning gravels suitable for resident trout were also observed (Golder 2015a).



Photo 6.2: Carbon Creek cascade barrier (Golder 2015).

6.6 Snowslide Creek

Snowslide Creek is not currently affected by mining and is not expected to be affected in the future (Table 6.1). Of the total stream length of 11.1 km, 32% is connected to the main stem (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality, calcite levels, and the benthic invertebrate community were not evaluated by Windward et al. (2014; Table 6.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). Tissue selenium concentrations were categorized as “good”.

Fish and fish habitat surveys were completed in 2008 and 2012 (Golder 2015a). A survey of the lower 3.8 km of Snowslide Creek characterized the stream as cascade-pool form, with habitats of cascade (71%), riffle (22%), glide (5.7%), and pool (2.2%) (Golder 2015a). Mean bankfull width was 3.6 m. Small cobble and large gravel were the dominant substrate types in cascade and riffle habitats, while medium and large gravel were present in glide and pool habitats. Riparian vegetation included coniferous, mixed

coniferous/deciduous and shrub forest or various successional stages. No off-channel habitat was observed. Two 100-m survey areas had gradients of 5% and 10% (Golder 2015a). The upper 1.9 km of Snowslide Creek is also dominated by cascade habitat (60%), with small amounts of riffle (4%), glide (4%), and pool (3%; Golder 2015a).

Between the upper and lower reaches described above is a potential barrier that has been characterized as both a 1.5-m vertical drop, and a 2-m bedrock cascade with 26% gradient (Golder 2015a; Photo 6.3). Slightly upstream from that is a bedrock falls downstream from which the stream goes subsurface due to a talus slope (521 m in length with a gradient of 52%); this is considered to be a permanent barrier to upstream fish migration. No fish have been found in the upper 1.8 km of the creek. The fish-bearing reach includes both westslope cutthroat trout and eastern brook trout. The dominant cover was overhanging vegetation, along with cobbles and boulders, and woody debris. There is suitable habitat for all life stages of fish during the open-water season. Pockets of spawning gravels suitable for salmonids were observed throughout. Overwintering potential was considered to be low to moderate (Golder 2015a).



Photo 6.3: Bedrock falls on Snowslide Creek (Golder 2015).

6.7 Wheeler Creek

Wheeler Creek is not affected by mining and is not expected to be affected in the future (Table 6.1). Of the total stream length of 60.8 km, 43% is connected to the main stem (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality, calcite levels, and the benthic invertebrate community were not evaluated by Windward et al. (2014; Table 6.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). Tissue selenium concentrations were categorized as “good”.

Fish and fish habitat surveys were conducted in 2007 and 2008, with more recent sampling in 2012 (Golder 2015a). The lower three reaches of Wheeler Creek (8.3 km in total) were dominated by cascade (42 to 55%) and riffle (24 to 45%) habitat types, with some glide (6.1 to 16%) and pool (3.1 to 6.0%) habitat types (Golder 2015a). Gradients were 4 to 6% with bankfull widths of 5.5 to 6.0 m, whereas the two reaches farther upstream had gradients of 6.6 to 18%. Cobble and gravel were the dominant substrate types. Riparian vegetation was primarily mixed deciduous and coniferous forest, although portions were dominated by shrubs (Golder 2015a).

The provincial database did not include fish distribution records for Wheeler Creek (Golder 2015a). Technical reports indicate westslope cutthroat trout and eastern brook trout have been captured or observed (Golder 2015a). About 8.5 km upstream from Michel Creek along the main stem is a 28% bedrock cascade, extending over 7 m, which acts as a permanent barrier to upstream fish migration. Temporary and semi-permanent barriers (e.g., steep-gradient, rock-debris cascades or falls) also limit access by fish to most of Little Wheeler Creek and an unnamed tributary to the north of Wheeler Creek (Photo 6.4). Surveys upstream from the barriers in 2012 did not yield fish. In the fish-bearing portion of Wheeler Creek, instream cover includes overhanging vegetation, undercut banks, boulder, and woody debris, although the small quantity of woody debris gave the fish-bearing reaches a poor rating for salmonid summer and winter rearing habitat. A small amount of channel braiding was noted in the most downstream reach. A 9-m side channel (Reach 2) and two 4-m side channels (Reach 3) were considered to be accessible during most flows while a 12-m side channel (Reach 2) is likely accessible only during high flows (Golder 2015a).



Photo 6.4: Example of steep-gradient cascade barrier in Wheeler Creek (Golder 2015).

The habitat of Wheeler Creek is considered suitable for rearing, holding and feeding for fish of various life stages, although pools were not considered deep enough (<1 m) to function for adult holding (Golder 2015a).. Small amounts of spawning gravels were observed, with greater abundance nearer to the Michel Creek confluence, but no redds were observed during spring spawning surveys. Little overwintering habitat is available in the creek (Golder 2015a).

Four wetlands were identified in upper portions of the Wheeler Creek basin, at one of which a western toad was observed (Teck undated).

6.8 Unnamed Tributary to Michel Creek

Unnamed Tributary is not affected by mining and is not expected to be affected in the future (Table 6.1). Of the total stream length of 10.1 km, 1% is connected to the main stem (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality, calcite, benthic invertebrate community, and tissue selenium concentrations were not evaluated by Windward et al. (2014; Table 6.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”).

Surveys of fish and fish habitat were conducted in 2012 and 2013 (Golder 2015a). A 131-m survey area, located about 500 m upstream from the confluence with Michel Creek was categorized as cascade-pool morphology, with predominantly cascade habitat (74%) and some glides (3.7%), pools (5.2%), and other features (17%). The average gradient was 10.7% in this section, with bankfull width of 2.0 m, and cobble and boulders being the dominant substrates. A 100-m section located about 3 km from the confluence with Michel Creek, was characterized as riffle-pool form with a lower average gradient of 4%, bankfull width of 2.8 m, and substrate consisting of gravel and fines. The riparian vegetation consisted of mixed coniferous and deciduous mature forest (Golder 2015a).

Two falls were identified within the lower portion of the creek, the first being about 130 m upstream from the confluence with Michel Creek and the second 140 m upstream from the first falls, which are both expected to limit the movement of fish from Michel Creek into Unnamed Michel Creek Tributary (Golder 2015a; Photo 6.5, Photo 6.6). Downstream from the lower falls, westslope cutthroat trout have been captured. Instream cover types included overhanging vegetation, boulders, undercut banks, and woody debris. There were isolated pockets of spawning substrate and there is moderate overwintering potential. Off-channel habitat was observed as a side channel habitat accessible to fish only at high flows. Based on multiple years of sampling, the creek upstream from the falls was classified as non-fish-bearing (Golder 2015a).

Two olive-sided flycatchers were observed in the headwater area of the Unnamed Tributary in 2013 (Golder 2015b).



Photo 6.5: First rock-fall barrier to fish movement on Unnamed Michel Creek Tributary (Golder 2015).



Photo 6.6: Second rock fall barrier to fish movement on Unnamed Michel Creek Tributary (Golder 2015).

6.9 Fir Creek

Fir Creek is not currently affected by mining and is not expected to be influenced in the future (Table 6.1). Of the total stream length of 19.1 km, 37% is connected to the main stem (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality, calcite, benthic invertebrate community, and tissue selenium concentrations were not evaluated by Windward et al. (2014; Table 6.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”).

Fish and fish habitat surveys were conducted in 2007, 2008, and 2012 (Golder 2015a). Surveyed areas of Fir Creek reflected relatively steep gradients (6.7 to 21%), with step-pool morphology, predominantly cascade (47 to 74%) and glide (19 to 47%) habitat, cobble-boulder-gravel substrate, and bankfull width of 3.6 to 5.6 m (Golder 2015a). Riparian vegetation was dominated by coniferous mature forest in the downstream reaches, and shrubs in the upstream reaches. Off-channel habitat was not observed (Golder 2015a).

Both westslope cutthroat trout and eastern brook trout have been reported in Fir Creek, although the most recent survey (2012) found only westslope cutthroat trout for which catch-per-unit effort was high relative to other streams surveyed in the region (Golder 2015a). A gradient barrier (>20% cascade) is present near the headwaters (Photo 6.7), about 7 km upstream from Michel Creek, which prevents upstream migration by fish and surveys upstream from the barrier have not yielded fish. Instream cover in the fish-bearing portion includes woody debris, boulders, and overhanging vegetation, with several deep pools present that may also provide overwintering habitat. The habitat was considered suitable for all life stages of fish. Some pockets of suitable spawning gravel were noted and two redds were confirmed in 2015 (Golder 2015a).

6.10 Alexander Creek

Alexander Creek has not been affected by mining activities (Table 6.1). All of the total stream length of 309 km is connected to the main stem (Table 6.3). The catchment has been disturbed by forestry activities and roads (Table 6.1).

Water quality was not evaluated by Windward et al. (2014; Table 6.2), but is likely comparable to other regional streams that have not been affected by mining (e.g., “good”). The benthic invertebrate community and tissue selenium concentrations were characterized as “good”. Low levels of calcite have been observed (calcite index: 0.48).



Photo 6.7: Gradient barrier defining upstream fish distribution limit in Fir Creek (Golder 2015).

Alexander Creek is frequently confined by valley walls in the lower reaches, broadens in the mid reaches and steepens and narrows again near the headwaters (Russell and Oliver 1996). Detailed fish and fish habitat surveys were conducted about 4 km upstream from the confluence with Michel Creek in 2006 and August 2012 (Lotic 2015a; Golder 2015a). The morphology of Alexander Creek was categorized as riffle-pool form (Lotic 2015a), having both riffle (58%) and cascade (24%) habitats (Golder 2015a). Stream gradient is about 2-3% (Lotic 2015a, Golder 2015a), with substrate dominated by cobble, boulder, and gravel (Lotic 2015a, Golder 2015a). Average bankfull width was about 16 m with riparian vegetation composed mainly of mature coniferous forest and mixed forest. Off-channel habitat was not observed (Golder 2015a).

A survey of a 50-m section of stream located about 7 km further upstream in 2012 characterized the stream section as having riffle and run habitat with 1% gradient, and predominantly cobble substrate with some sand (Minnow 2014).

Westslope cutthroat trout, bull trout, and eastern brook trout were captured in 2006 (Lotic 2015a). Historical fish habitat use information for Alexander Creek was not available in any of the data sources reviewed by Golder (2015), but 18 cutthroat trout and

one bull trout were captured in August 2012. Cover for fish included woody debris and deep pools (Golder 2015a) with trace amounts of undercut banks (Lotic 2015a). Habitat was considered suitable for all life stages of fish during the open-water season, and two deep pools were documented that may provide overwintering habitat for juvenile and adult fish (Golder 2015a). Isolated pockets of substrate suitable for cutthroat trout spawning were found. No barriers to fish passage were reported during the surveys, however a bedrock formation downstream from Highway 3 may be a barrier to fish movement during some flows (M. Robinson, Lotic Environmental, pers. comm.). In the lower watershed, major realignment of the stream channel has taken place to accommodate highway repair and gas pipeline (Russell and Oliver 1996).

Olive-sided flycatcher is a species at risk that was observed near Alexander Creek in a breeding bird survey conducted in 2012 (Minnow 2014a; Table 6.3).

6.11 Erickson Creek

About 26% of the total catchment area of Erickson Creek has been disturbed by mining activities (Table 6.1). Mining currently occurring within the permit boundary will continue into the near and far future. All of the total stream length of 82.4 is either fragmented or has been lost (Table 6.3). There are four ponds near the headwaters of Erickson Creek (Photo 6.8) and then a section of sub-surface flow before the stream flows above ground for the remaining 2.3 km to the confluence with Michel Creek (Lotic 2015a). With the exception of the upper-most pond, all ponds and reaches on Erickson Creek are influenced by EVO mine activities. There has also been historical disturbance from forestry activities and roads (Table 6.1).

Water quality was characterized as “poor” based on Windward et al. (2014), although concentrations of selenium, nitrate, and cadmium have been decreasing (Zadjlik and Minnow 2013; Table 6.2). The benthic invertebrate community was characterized as “poor” but tissue selenium concentrations were characterized as “good”. Moderate levels of calcite have been observed (calcite index: 1.77).

Active water treatment is scheduled for 2020 (EVO Phase I AWTF), which will treat water from Erickson, Bodie, and Gate Creeks (Teck 2014). Treated effluent will discharge into Erickson Creek. The second phase of AWTF is scheduled for Erickson Creek in 2024.



Photo 6.8: Most downstream pond in Erickson Creek headwaters (Lotic 2015a).

A fish and fish habitat survey was conducted in the lower watershed in 2009 and four areas along the creek and the four headwater ponds were surveyed in 2013 and 2014 (Lotic 2015a). The lower 290 m of the creek from the confluence with Michel Creek to a 2-m-high waterfall, is dominated by riffle-pool form with several cascades, a gradient of 4%, and bankfull width of 5.6 m. In this section of the creek, calcite is considered the dominant substrate, with a large amount of moss also growing in the stream (Photo 6.9). Riparian vegetation was predominantly coniferous forest. Only this lower section of creek is fish-bearing due to the barrier waterfall. Although westslope cutthroat trout, bull trout, and mountain whitefish have been reported in Erickson Creek historically, fish densities are low. There is no suitable spawning habitat due to calcite deposits and a lack of suitable spawning gravel. There is also poor cover. Overwintering habitat potential was considered moderate, based on water quality and water depths (Lotic 2015a).



Photo 6.9: Moss and calcified streambed in Erickson Creek (Lotic 2015a).

Upstream from the waterfall, stream gradient ranged from 3.5% to 13%, with bankfull widths of 5.2 to 5.9 m and predominantly cobble and gravel substrates and variable amounts of calcite dominating the substrate. Riparian vegetation varied from coniferous forest in the upstream and downstream sections to mixed forest in the middle section (Lotic 2015a).

Olive-sided flycatchers and spotted sandpipers were observed within the Erickson Creek basin in 2013 (Golder 2015b).

6.12 South Pit Creek

About 56% of the total catchment area of South Pit Creek has been disturbed by mining activities (Table 6.1). Mining is currently occurring within the permit boundary and will continue into the near future with reclamation (Table 6.1). All of the total stream length of 1.3 km is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).

Water quality was characterized as “poor” based on Windward et al. (2014) and increasing trends in the concentrations of selenium and sulphate were reported (Zadjlik and Minnow 2013; Table 6.2). Windward et al. (2014) did not report data for benthic invertebrate community, tissue selenium concentrations, or calcite.

South Pit Creek is not fish-bearing and habitat surveys have not been completed.

6.13 Milligan Creek

About 32% of the total catchment area of Milligan Creek has been disturbed by mining activities (Table 6.1). Mining is currently occurring within the permit boundary and will continue into the near future with reclamation (Table 6.1). Of the total stream length of 7.2 km, 61% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).

Water quality was characterized as “poor” (Windward et al. 2014) and increasing trends in the concentrations of selenium and sulphate were reported (Zadjlik and Minnow 2013; Table 6.2). A decreasing trend in the concentration of nitrate was observed. Windward et al. (2014) did not report data for benthic invertebrate community, tissue selenium concentration, or calcite.

Milligan Creek is not fish-bearing and no habitat surveys have been completed.

6.14 Gate Creek

About 56% of the total catchment area of Gate Creek has been disturbed by mining activities (Table 6.1). Mining is currently occurring within the permit boundary and will continue into the near and far future (Table 6.1). Of the total stream length of 5.9 km, 7% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).

Water quality was characterized as “poor” based on Windward et al. (2014) and increasing trends in the concentrations of selenium and sulphate, and a decreasing trend in the concentration of nitrate, were reported (Zadjlik and Minnow 2013; Table 6.2). No data are available for the benthic invertebrate community, tissue selenium concentration, or calcite.

Active water treatment is scheduled for 2020 (EVO Phase I AWTF), which will treat water from Erickson, Bodie, and Gate Creeks (Teck 2014). Treated effluent will discharge into Erickson Creek.

The lower portion of Gate Creek is a settling pond (6,562 m²), with fine substrate and maximum depth of 3 m (Lotic 2015a). Gate Creek and South Gate Creek both flow west from EVO mining areas around the Bodie spoil and under the railway tracks and Highway 3 before joining and entering into Gate Pond. Fish and fish habitat surveys were conducted in 2013 (Lotic 2015a). The main stem of Gate Creek was characterized as a very narrow (0.5 m width), shallow cascade with a gradient barrier. South Gate Creek was characterized as having cascade-pool morphology, mean gradient of 1.5%, and mean bankfull width of 2.07 m, with the dominant substrates being fines and gravel. No off-channel habitat was reported and riparian vegetation was not documented for this creek (Lotic 2015a).

Gate Pond has a small concrete culvert and flap valve (Photo 6.10) at the outlet which may act as a deterrent to fish passage but is not considered a barrier (Lotic 2015a). Fish access into the pond was easier prior to installation of the flap valve in 2011. Fish species captured or observed in Gate Pond and/or Gate Creek were longnose sucker, longnose dace, and westslope cutthroat trout (Lotic 2015a). Good cover is provided by aquatic vegetation along the shore and by the depth of the pond. The pond habitat was considered suitable for longnose sucker and density of this species was highest of all areas surveyed in an evaluation of fish species distribution in the Elk Valley in 2010 (Interior Reforestation 2014). The creek upstream from Gate Pond is considered fish-bearing for 730 m, because there are no barriers between the creek and the pond, but it was concluded to have limited fish use. South Gate Creek provides suitable habitat for all life stages of longnose sucker, but the lentic pond habitat would likely be preferred over the flowing creek. Spawning habitat suitable for cutthroat trout was not present (Lotic 2015a).

6.15 Bodie Creek

About 98% of the total catchment area of Bodie Creek has been disturbed by mining activities (Table 6.1). Mining is currently occurring within the permit boundary and will continue into the near and far future (Table 6.1). Of the total stream length of 12.5 km, 6% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).



Photo 6.10: Downstream view of the concrete culvert and flap valve for Gate Creek (Lotic 2015a).

Water quality was characterized as “poor” (Windward et al. 2014; Table 6.2). Increasing trends in the concentrations of selenium and sulphate have been reported, while trends for nitrate and cadmium have been decreasing (Zadjlik and Minnow 2013). The benthic invertebrate community was characterized as “poor” but tissue selenium concentrations were characterized as “good”. Low levels of calcite have been observed (calcite index: 0.06; Windward et al. 2014).

Active water treatment is scheduled for 2020 (EVO Phase I AWTF), which will treat water from Erickson, Bodie, and Gate Creeks (Teck 2014). Treated effluent will discharge into Erickson Creek.

Bodie Creek flows from the Bodie Rock Drain into Michel Creek. Fish and fish habitat surveys were conducted in 2001 and 2007, and in March, August, and October 2013 and June 2014 (Lotic 2015a). The lower 415 m of Bodie Creek, upstream from the confluence with Michel Creek, has been highly modified by channelization of flow around existing

buildings, Highway #3, and the CPR rail tracks (i.e., Warehouse 50 storage yard). The morphology of this reach is classified as riffle-pool form with an average gradient of 0.5%. A high sediment load from upstream mining, railway and highway results in the dominant substrate being fines. A hanging culvert separates this section of the creek from two EVO sediment ponds (1,554 and 3,078 m²), which are connected to each other by a 140 m long ditch. The upper 2 km of the creek, upstream from the sediment ponds, has a gradient that increases from 1% to 18% moving upstream with stream morphology shifting from riffle-pool form to step-pool form. Average bankfull width varies from 1.9 m to 3.3 m throughout the length of the creek. Riparian vegetation is predominantly grasses, with a few deciduous trees near the confluence with Michel Creek. No off-channel habitat was reported (Lotic 2015a).

Only the lower, channelized portion of the stream is considered fish-bearing (Lotic 2015a). The culvert located immediately downstream from the lower settling pond, is considered a barrier to upstream fish movement (Photo 6.11). Fish were salvaged from the sediment ponds in 2007 and none were observed when the ponds were dewatered for maintenance in 2014. The lower reach of Bodie Creek provides very little cover for fish, consisting mainly of overhanging grasses. The habitat is suitable for cyprinids (minnows) and catostomids (suckers), with some spawning potential particularly for longnose dace. Fish species captured or observed in the creek include westslope cutthroat trout, brook trout, mountain whitefish, longnose sucker, and longnose dace, with longnose dace (family Cyprinidae) found in highest density. Lack of cutthroat trout fry suggests lack of spawning within the creek itself, but moderate juvenile densities suggest it is utilized for off-channel rearing (Lotic 2015a).

No species at risk were observed in breeding bird and amphibian surveys conducted at the Bodie Creek setting ponds in 2012 (Minnow 2014a; Table 6.3).

6.16 Aqueduct Creek

About 33% of the total catchment area of Aqueduct Creek has been disturbed by mining activities (Table 6.1). Mining is currently occurring within the permit boundary and will continue into the near and far future. Of the total stream length of 6.9 km, 32% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 6.3). The catchment has also been disturbed by forestry activities and roads (Table 6.1).



Photo 6.11: CPR double culverts in Bodie Creek (Lotic 2015a).

Water quality was characterized as “good” based on Windward et al. (2014; Table 6.2). Tissue selenium concentrations in periphyton and benthic invertebrates were considered “poor” in terms of potential for effects to invertebrates or fish consumers of invertebrates, but were “good” in terms of potential effects to birds feeding on invertebrates. No data were available for benthic invertebrate community or calcite (Table 6.2).

Fish and fish habitat surveys were completed in 2005, in March and August-September, 2013, and in January, 2015 (Lotic 2015a). The downstream portion of Aqueduct Creek is a wetland (pond) habitat (1,720 m²) with a seasonal overland connection to Michel Creek through a culvert under the Great Northern Road, which runs parallel to the creek (Lotic 2015a; Photo 6.12). Substrate in the pond is composed of fine particles. There is good cover provided by overhanging and aquatic vegetation, large woody debris, and a 2-m deep area at the southern end of wetlands. The riparian vegetation was mixed forest, dominated by mature cottonwood. Longnose sucker are present and likely reside in the pond year-round, where there is suitable habitat present for spawning, feeding/resting, and overwintering of this species. The habitat is not

considered suitable for salmonid spawning and is likely used only seasonally by juvenile salmonids which can enter and exit the pond during high water (Lotic 2015a).



Photo 6.12: Wetland habitat in lower Aqueduct Creek (Lotic 2015a).

Extending about 1.5 km upstream from the wetland, Aqueduct creek has been modified by channelization through pasture land and residential development, resulting in extensive glides mixed with a riffle-pool form (Lotic 2015a). The gradient of this portion of the stream is fairly low, gradually increasing from 0.7% to 1.5% (moving upstream), with average bankfull width between 0.78 and 1.1 m and substrates dominated by gravels and fines. A sediment pond system that is a barrier to upstream fish movement was constructed in summer 2015 to be completed in summer 2016 about 1 km upstream from the wetland.

Farther upstream in the headwaters, Aqueduct Creek is dominated by a step-pool morphology with a gradient of 21%. Riparian vegetation changes from mixed forest with mature cottonwood near the mouth, to pastureland and development mid creek to mixed deciduous trees and grasses in the headwaters (Lotic 2015a).

Spring Creek (fish-bearing) and Qualtieri Creek (no fish have been observed) are small tributaries to Aqueduct Creek (Lotic 2015a). In Aqueduct Creek, although there is no barrier to fish movement for 1.0 km upstream from the wetland, only the lower 440 m of the system between Spring Creek and Michel Creek has been confirmed as fish-bearing (Lotic 2015a). Overhanging vegetation provides fair cover for fish, along with small quantities of small and large woody debris. This section supports all life stages of longnose suckers and cyprinids, including spawning, especially in the wetland area. Westslope cutthroat trout, bull trout, eastern brook trout and mountain white fish have also been observed in this lower section. Fish utilization of reaches farther upstream is likely limited by shallow water (dry in some seasons), highly modified channel morphology (e.g., ditching), and cover (Lotic 2015a).

Olive-sided flycatchers were observed near the headwaters of Aqueduct Creek in 2013 (Golder 2015b).

7.0 TRIBUTARIES TO MIDDLE ELK RIVER (MU4/MU5)


Tributaries in the Middle Elk River Watershed (MU4/MU5) are shown in Figure 7.1, and include:

- Grave Creek (Including Harmer Creek and EVO Dry Creek);
- Six-Mile Creek;
- Balmer Creek;
- Fennelon Creek;
- Goddard Creek;
- Otto Creek;
- Transmission Creek;
- Mine Creek (Hosmer Creek)

Matrices for these tributaries provide data related to watershed area and disturbances (Table 7.1), environmental quality descriptors (Table 7.2) and fish and aquatic habitat (Table 7.3). An orthophoto figure for each tributary is found in Appendix E. Each tributary is briefly described in the following sections.

7.1 Grave Creek (including Harmer Creek and EVO Dry Creek)

About 8% of the total catchment area of Grave Creek has been disturbed by mining activities (Table 7.1), and the BRE Project will extend mining operations into the EVO Dry Creek drainage. Harmer Creek is a major tributary of the Grave Creek system, and EVO Dry Creek, located on the north-east side of EVO, flows into Harmer Creek. Grave Creek upstream from the Harmer/Grave confluence is not mine influenced, nor are the headwaters of Harmer Creek. Of the total stream length of 158 km, 1% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 7.3). The catchment has also been disturbed by forestry activities and roads (Table 7.1). Grave Lake, which flows into Grave Creek from the north, is used as a recreational site.



Kilometers

MAP INFORMATION
Datum: NAD 1983 Map Projection: Zone 11N
Data Source: Department of Natural Resources Canada. All rights reserved.
Creation Date: Sept 2015

Legend

Water Flow Direction	Waterbody
Mid Elk River	Streams
Watershed Boundary	Mine Development Areas
Falls	Teck Coal Mine Operation
Community	Provincial Boundary

**Figure 7.1. DRAFT
Middle Elk River
Watershed Boundary**

Teck

1:125,000

Table 7.1: Watershed area and disturbances associated with Middle Elk River (MU4 and MU5). Data sources are described in Section 2.0.

Tributary Name	Catchment Area				Stressors																		Mine Alterations/Disturbances/Planned Mitigation	
	Total Catchment Area from Provincial Corporate Watershed Base Dataset (km ²) (1980s plus updates)	Total Catchment Area as of September 2015 (km ²)	% Change in Total Catchment Area	Teck Ownership Rights: % of Catchment Area and Type (Coal, Surface, Timber)	Mine-Related Disturbances in Catchment as of September 2015										Forestry Disturbances in Catchment				Road Disturbances in Catchment				Additional Planned Near/Far Future Mining Disturbances	Planned Mitigation/Fish Habitat Offsetting
					Timing of Disturbance: Historical (H), Current (C), Near Future (NF), Far Future (FF) or Not Expected (N)	Total Footprint Area Currently Disturbed (excludes reclaimed lands)		Reclaimed Land Area		Pit(s) (yes/no)	Waste Rock Storage (yes/no)	Rock Drain(s) (yes/no)	Channel Re-alignment or Diversion (yes/no)	Settlement Ponds/Catch Basins (number)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (km ²)	Equivalent Clearcut Area within Catchment Area from 1955-2010 (%)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (km ²)	Equivalent Clearcut Area within Catchment Area from 2011 to 2015 (%)	Road Density per km ²	Stream Crossings per km	Roads Within 100 m of a Stream	% Roads on Slopes Greater than 60%		
						km ²	%	km ²	%															
Grave Creek (including Harmer and EVO Dry creeks)	81.1	78.7	-3	1% Coal, 42% Surface and Coal, 16% Surface and Coal and Timber, 35% Surface and Timber	H, C, NF, FF	6.3	8	2.4	3	N	Y	N	N	2	11.8	15	0.3	0	1.9	1.2	0.7	0.2	The BRE Project will extend mining operations into the EVO Dry Creek drainage.	Fish habitat offsetting as part of the regional fish habitat management plan will be developed in consultation with DFO, FLNRO, and KNC representatives as part of the BRE project. Fish salvage and exclusion is planned for the EVO Dry Creek sediment ponds. From a regional water quality management perspective, the Harmer Compliance Point Selenium Evaluation Report identifies options for selenium management in Harmer Creek.
Six-mile Creek	5.1	5.1	0	94% Surface and Coal, 4% Surface and Timber	H	0.4	8	0.3	6	N	Y	N	Y	3	1.9	37	0.6	12	3.7	0.4	0.6	0.1	There are no additional near or far future mine disturbances planned.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Balmer Creek	4.2	4.1	-2	98% Surface and Coal	H	1.0	24	0.8	19	N	Y	N	N	0	1.5	36	0.0	0	5.0	1.6	0.6	0.2	There are no additional near or far future mine disturbances planned.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Fennelon Creek	5.7	2.6	-54	100% Surface and Coal	H, C, NF, FF	2.7	47	1.2	21	N	N	Y	N	0	1.4	25	0.0	0	4.3	1.0	0.5	0.3	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Goddard Creek	3.9	5.5	41	100% Surface and Coal	H, C, NF, FF	2.5	64	0.9	23	Y	Y	N	Y	4	0.1	3	0.0	0	3.4	4.3	0.9	0.5	Mining currently occurring within the permit boundary will continue into the near and far future.	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, selenium management to be determined.
Otto Creek	2.8	4.8	71	100% Surface and Coal	H, C, NF, FF	1.4	50	0.6	21	Y	Y	N	Y	5	0.1	4	0.0	0	5.2	3.6	0.9	0.6	Mining currently occurring within the permit boundary will continue into the near and far future.	Fish salvage and exclusion is planned for the sediment ponds. No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Transmission Creek	10.2	10.2	0	21% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	3.7	36	0.0	0	3.2	3.6	1.3	0.4	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.
Hosmer Creek (including Mine Creek)	33.1	33.1	0	26% Coal, 34% Surface and Coal	N	0.0	0	0.0	0	N	N	N	N	0	5.6	17	0.0	0	3.5	3.1	1.6	0.6	NA	No fish/aquatic habitat mitigation is planned. From a regional water quality management perspective, there is no planned mitigation.

NA - Not Applicable

Table 7.2: Environmental quality descriptors for Middle Elk River (MU4 and MU5). Data sources are described in Section 2.0.

Tributary Name	Teck Surface Water Quality Monitoring Station (Code, Permitted-P or Non-Permitted-NP, Years Data Available, and [EMS# if applicable])	Surface Water Chemistry (2011-2013) (from Windward et al. 2014) ^a									Median Concentrations and Trends Blue- and orange-highlighted cells indicate decreasing and increasing trends, respectively (from Zadjlik and Minnow 2013).										Total Loads (2012) (from Zadjlik and Minnow 2013) Bold load values are associated with 9+ months of data. Non- bold values should be interpreted with caution.						Surface Water Acute Toxicity (2015) (Q1-Q4)		Sediment Quality (2013) (Minnow 2014b)		Calcite (2013) (from Windward et al. 2014)	Benthic Invertebrate Community Structure (2012) (Windward et al. 2014)	Max SQ ^e for Selenium in Periphyton or Invertebrate Tissue (mean SQ in brackets if max SQ is >1) (Windward et al. 2014)								
		Water Quality Index (WQI) (see Section 2.0 and Table 2.4 for explanation about how WQI was computed and definitions for colour categories)						Dissolved Cadmium (ug/L) *Not included in WQI		Data Period for Selenium, Nitrate, and Sulphate Trends	Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)		Cadmium (mg/L)		Colour categories explained in Table 2.4 and text																						
		Median (# Samples >Level 1 BM 19 ug/L out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Selenium (µg/L)		Nitrate-N (mg/L)		Sulphate (mg/L)			Median (# Samples >Level 1 hardness-dependent BM out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median (# Samples >Level 1 BM of 481 mg/L out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median (# Samples >Level 1 hardness-dependent BM out of Total # Samples) - grey shade indicates more than 10% of samples exceeded benchmark	Mean	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Median Concentration 2012 (µg/L)	Trend (%/year) (p-value) - colour codes defined above	Data period for Cadmium Trend	Median Concentration 2012 (µg/L)	Trend (% year)	Selenium (ton/year)	% of Total Tributary Load	Nitrate (ton/year)	% of Total Tributary Load	Sulphate (ton/year)	% of Total Tributary Load	Rainbow trout (% Mortality)	Daphnia magna (% Mortality)	Chemistry	Toxicity Survival and growth of <i>Hyalella azteca</i> and <i>Chironomus riparius</i>	Calcite Index	% EPT ^b	% Ephemeroptera	Direct Effects to Invertebrates (periphyton consumption)	Fish Dietary Exposure (Invertebrate Consumption)	Bird Dietary Exposure (Invertebrate Consumption)
HACKUS		ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.15	76	32	0.5	0.8	0.6						
Grave Creek (including Harmer and EVO Dry creeks)	Harmer Pond Outlet EV_HC1 (P), 1990-2015 [E102682]	72	30.2 [58 of 62]	29.9	1.07 [0 of 62]	1.09	188 [0 of 62]	1.72	0.018 [0 of 62]	0.0191	2004- 2012	33	3.0 (0.0003)	1.14	NT	180.00	3.3 (0.0002)	2008- 2012	0.02	NT	0.6	4	28	0.9	3,698	6	0, ND, ND, ND	3, ND, ND, ND	Harmer Settling Pond (n=5 samples): 0/12 metals and 3/16 to 5/16 PAH above high SQG and reference concentratio ns all samples.	Harmer Settling Pond (HA7): No effect mean survival or growth <i>Chironomus riparius</i> , <i>Hyalella azteca</i>	0.17	46	30	0.5	0.6	0.4					
	GRCK	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.24	81	53	0.7	0.8	0.6						
	GRDS	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.54	61	29	0.6	0.7	0.5						
Six-mile Creek	EV_SM1 (P), 1990-2105 [E102681]	100	2.11 [0 of 49]	2.17	0.0321 [0 of 49]	0.141	59.9 [0 of 49]	59.9	0.00627 [0 of 49]	0.00727	2004- 2012	2.2	-2.2 (0.0130)	0.06	-10.9 (<0.0001)	64.73	2.3 (0.0006)	2007- 2012	0.02	NT	0.0	0.0	0.1	0.0	0.1	0.0	0, 0, 0, 0	0, 0, 0, 0	ND	ND	ND	ND	ND	ND	ND						
Balmer Creek	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND		ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND				
Fennelon Creek	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND		ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND				
Goddard Creek	EV_GC2 (P), 1990-2015 [E208043]	58	14.2 [80 of 80]	24.1	1.63 [8 of 70]	1.77	190 [0 of 70]	186	0.0191 [0 of 70]	0.0254	2004- 2012	12	-17.2 (<0.0001)	1.56	-11 (<0.0001)	175.00	-4.7 (0.0002)	2008- 2012	0.07	NT	0.1	0.4	4	0.1	4	0.0	0, 0, 0, 0	0, 0, 0, 0	Goddard Marsh (n=5 samples): 0/12 metals and 2/16 PAH above high SQG and reference concentratio ns in all samples.	Goddard Marsh (GO13): reduced mean survival of <i>Chironomus riparius</i> ; no effect <i>Hyalella azteca</i> survival or growth.	ND	ND	ND	ND	ND	ND					
Otto Creek	EV_OC1 (P), 1990-2015 [E102679]	100	1.72 [0 of 63]	1.72	0.0876 [0 of 63]	0.102	60.2 [0 of 63]	59.3	0.0123 [0 of 63]	0.014	2004- 2012	2.3	NT	0.12	-5.1 (0.0002)	60.60	5.4 (<0.0001)	2007- 2012	0.04	NT	0.0	0.0	0.3	0.0	0.3	0.0	0, 0, 0, 0	0, 0, 0, 0	ND	ND	0.02	38	13	0.3	0.4	0.3					
Transmission Creek	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.6	0.7	0.5					
Hosmer Creek (including Mine Creek)	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.6	0.7	0.5						

ND - No Data; NT - No trend
^a Based on monitoring data and site-specific targets established in EVWQP for nitrate, sulphate, cadmium, and selenium (see Appendices A and D in Windward et al. 2014).
^b EPT - Ephemeroptera (mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies).
^c HQ - Hazard Quotient computed as a ratio of tissue concentration to relevant benchmark (Windward 2014). Values above 1 indicate potential risk.
^d Quarterly monitoring data will be summarized in December 2015 version for locations monitored quarterly under Permit 107517.
^e Data for this location may be available but were not assessed by Windward et al. (2014).

Table 7.3: Aquatic habitat of Middle Elk River (MU4 and MU5). Data sources are described in Section 2.0.

Tributary Name	Annual Average Daily Flow (m³/s)	Ktunaxa traditional uses and cultural values	Stream Length based on Province's CWB Layer									Fish and Fish Habitat															Amphibian census location, date (and any observed species-at-risk)	Bird census location, date (and any observed species-at-risk)		
			Total (km) (1980s plus Updates)	Currently Connected to Main Stem		Currently Fragmented				Permanently Altered or Destroyed		Fish Species Present	Unique Fish Habitat Features (Yes/No/Unknown)	Barrier(s) Present (excludes infills)	Total Fish Habitat (1980s plus Updates)		Total Non-fish Bearing (1980s plus Updates)		Fish Habitat Currently Connected to Main Stem		Fish Habitat Currently Fragmented				Fish Habitat Permanently Altered or Destroyed				Riparian Habitat within Catchment Area	
						Potentially Reversible		Permanently Isolated													Potentially Reversible		Permanently Isolated							
				km	%	km	%	km	%	km	%				km	%	km	%	km	%	km	%	km	%	km	%			km	%
Grave Creek (including Harmer and EVO Dry creeks)	1.11		157.8	1.1	1	152.1	96	1.2	1	3.5	2	WCT, BT, MW, KO, RB, EB, LSU, RSC	Y	A, N	35.8	23	122.1	77	1.0	3	33.5	94	0.1	0	1.2	3	6.5	8	Harmer Pond (HA7), May 2012; Dry Creek, 2013 (western toad)	Harmer Creek, 2013 (olive-sided flycatcher)
Six-mile Creek	0.036		5.0	0.8	16	4.2	84	0.0	0	0.0	0	WCT, BT, MW, EB, LSU, LNC, RSC	U	A, N	0.6	12	4.4	88	0.6	100	0.0	0	0.0	0	0.0	0	0.3	6	ND	Near mouth, 2013 (olive-sided) flycatcher
Balmer Creek	ND		3.2	0.6	19	2.6	81	0.0	0	0.0	0	WCT, BT, MW, EB	U	0	0.6	19	2.6	81	0.6	100	0.0	0	0.0	0	0.0	0	0.2	5	ND	Near creek, 2013 (olive-sided flycatcher)
Fennelon Creek	ND		3.0	0.0	0	3.0	100	0.0	0	0.0	0	None	N	N	0.0	0	3.0	100	0.0	0	0.0	0	0.0	0	0.0	0	0.1	2	ND	ND
Goddard Creek	0.020		8.1	1.0	12	0.4	5	4.0	49	2.6	32	WCT, MW, EB, LSU, LNC, RSC	U	A	1.2	15	6.9	85	1.0	83	0.0	0	0.0	0	0.2	17	0.3	8	Goddard Marsh (GO13), May 2012.	Goddard marsh (GO13), June 2012 (bald eagle). Herons nested near the marsh in the past, but they moved their rookery about 1 km south prior to 2012 (Minnow 2014a, Golder 2015c)
Otto Creek	0.026		5.1	1.3	25	0.3	6	1.6	31	1.9	37	WCT, BT, MW, EB, LSU, LNC, RSC	U	P	1.3	25	3.8	75	1.3	100	0.0	0	0.0	0	0.0	0	0.1	4	Otto Creek (OTTO), May 2012 (Western toad).	Otto Creek (OTTO), June 2012
Transmission Creek	ND		25.3	25.3	100	0.0	0	0.0	0	0.0	0	WCT, EB	U	A, N	4.0	16	21.2	84	4.0	100	0.0	0	0.0	0	0.0	0	1.3	13	ND	ND
Hosmer Creek (including Mine Creek)	ND		81.6	81.6	100	0.0	0	0.0	0	0.0	0	WCT, MW, EB, LSU	U	0	18.8	23	62.8	77	18.8	100	0.0	0	0.0	0	0.0	0	4.1	12	ND	ND

ND - no data; WCT - westslope cutthroat trout (Blue-list conservation status); BT - bull trout (Blue-list conservation status); MW - mountain whitefish; KO - kokanee; RB - rainbow trout; EB - eastern brook trout; LSU - longnose sucker; RSC - redbside dace; LNC - longnose dace.

^a Available data set is missing flows for some low-flow months.

Water quality and the benthic invertebrate community monitored at the Harmer Pond outlet were characterized as “fair” and tissue selenium concentrations were characterized as “good” (Windward et al. 2014; Table 7.2). Increasing trends in the concentrations of selenium and sulphate were reported (Zadjlik and Minnow 2013; Figure 7.2). Calcite deposits are substantial in Dry Creek but diminish downstream in Harmer and Grave Creeks (Robinson and MacDonald.2014).




Figure 7.2: Selenium concentration trend over time (2004-2013) at the Harmer Pond outlet into Gate Creek (EV_HC1).

Grave Creek Fish Habitat

Fish and fish habitat surveys were conducted in June 2013 and 2014 in Grave Creek, as well as Harmer Creek and EVO Dry Creek (Lotic 2015a). A 100-m area was surveyed in each of three reaches in Grave Creek, all having morphology classified as riffle-pool, average gradients of 1.8% to 3.0%, bankfull widths of 5.6 to 7.4 m, and substrate dominated by cobble and boulder. The riparian vegetation was mixed forest (Lotic 2015a).

Fish species that have been captured or observed in Grave Creek itself include westslope cutthroat trout, rainbow trout, bull trout, eastern brook trout, kokanee, mountain whitefish, longnose sucker, and reddsideshiner (Lotic 2015a). The most downstream section of the creek had moderate to good spawning habitat for westslope cutthroat trout, including gravel, overhead cover, good flow, adequate water depth and good proximity to holding water; however no redds were observed. This section of the creek was considered suitable for use by all life stages during the open water season but overwintering potential

was considered poor due to low water temperatures. The section of creek downstream from the Grave Lake outlet includes four bedrock waterfalls (1-3 m in height) that were reported as possible barriers to upstream movement of fish (Lotic 2015a; Photo 7.1). Upstream from the confluence with the Grave Lake outlet, where Grave Creek enters the canyon, the only species of fish observed by Lotic (2015a) was westslope cutthroat trout which was attributed to the falls downstream as potential barriers. However, sampling in 1996 captured or inferred (based on captures in adjacent reaches and/or previously reported captures) the presence of westslope cutthroat trout, bull trout, and rainbow trout as far upstream in Grave Creek as the reach encompassing the mouth of Harmer Creek; however, access to reaches farther upstream was precluded by a hanging culvert with a 60 cm drop at the road crossing and a 7% gradient Interior Reforestation (1997). Headwater resident westslope cutthroat trout were confirmed upstream from the culvert in 1996 (Interior Reforestation 1997).



Photo 7.1: Upstream view of the 3 m high waterfall below the Grave Lake confluence in Grave Creek (Lotic 2015a).

In a spring survey by Lotic (2015a), areas were identified as having good potential spawning area for WCT but no redds were observed. Habitat was considered suitable for multiple life stages, and overwintering potential was good (Lotic 2015a).

There are no permanent barriers to fish movement between Grave Creek and Grave Lake, although connectivity is seasonally restricted. Rainbow trout and kokanee were stocked in Grave Lake prior to 2000, but not since then, so hybridization between rainbow trout and westslope cutthroat trout is not expected (Lotic 2015a).

Harmer Creek Fish Habitat

Harmer Creek flows into Grave Creek and its headwaters, upstream from the confluence with Dry Creek, are not exposed to any direct mining activities. A dam and spillway is located on Harmer Creek, about 0.5 km upstream from the confluence with Grave Creek, is a barrier to upstream fish movement (Lotic 2015a). The dam creates Harmer Pond, which is about 19,000 m² and has a maximum depth of 5 m (Photo 7.2).



Photo 7.2: Harmer Pond 0.5 km upstream of the confluence of Harmer Creek with Grave Creek (Lotic 2015a).

Harmer Pond appears to provide habitat suitable to support all life stages of fish year-round except for spawning, but no recent surveys have caught or observed any fish (Lotic 2015a). Cutthroat trout have been captured in the creek upstream from the pond so both the creek and pond are considered fish-bearing. Upstream from the pond to the confluence with Dry Creek is habitat suitable for all life stages, with fair cover, good overwintering potential, and confirmed spawning. However, use by fish might be limited by warm summer temperatures. There are no barriers preventing access by trout to the headwaters upstream of mining influence, where there is also diverse habitat for all life stages, including potential spawning and overwintering habitats (Lotic 2015a).

No species at risk were observed in breeding amphibian and bird surveys completed at Harmer Pond in 2012 (Minnow 2014a), but olive-sided flycatchers were observed near the confluence of Harmer and Grave Creeks in 2013 (Golder 2015b) (Table 7.3). American dippers were also observed near the confluence with EVO Dry Creek, as well as farther downstream in Harmer Creek, and a Harlequin duck was observed at Harmer Pond (Golder 2015b).

EVO Dry Creek Fish Habitat

The Dry Creek watershed has been directly influenced by mining, including the construction of the Dry Creek Spoil (Lotic 2015a). Sampling has been minimal due to safety concerns with the proximity of the spoil runout zone. Extensive calcite deposition has occurred all reaches of the creek, with evidence that calcite terracing has altered stream direction into forested riparian areas at certain locations, and limiting aquatic habitat in all reaches (Lotic 2015a). Although one cutthroat trout was captured in the lower 120 m of the creek, habitat was considered minimal for all life stages. A constructed settling pond (3,218 m²) was located upstream from this section of creek, and could be a barrier to smaller fish migration. A 2008 survey reported catching cutthroat trout in the pond (Lotic 2015a). Dry Creek continues upstream from the settling pond about 0.7 m to a series of ponds, and then about one km farther to the runout of the Dry Creek Spoil. Both sections were inferred to be fish bearing due to a lack of barrier, but no fish were captured or observed during recent surveys (Lotic 2015a).

A Western toad was observed near EVO Dry Creek in 2013 (Golder 2015b).

7.2 Six-Mile Creek

About 8% of the total catchment area of Six-Mile Creek has been disturbed by mining activities (Table 7.1). There are no additional mine disturbances planned for the near or

far future. Of the total stream length of 5 km, 16% is connected to the main stem (Table 7.3). The catchment has also been disturbed by forestry activities and roads (Table 7.1).

Water quality was characterized as “good” (Windward et al. 2014; Table 7.2). Decreasing trends in the concentrations of selenium and nitrate, and an increasing trend for sulphate, were reported (Zadjlick and Minnow 2013). Windward et al. (2014) did not include data for calcite, benthic invertebrate community condition, or tissue selenium concentrations.

Based on a survey conducted in June 2013, Lotic (2015a) reported that the lowest section of Six Mile Creek is channel habitat about 170 m long flowing into the Elk River (Photo 7.3). The upstream limit of this section is marked by a concrete spillway, which also represents the upstream limit of fish distribution in Six Mile Creek. A 100-m section was characterized as riffle-pool morphology, with mean gradient of 1.25%, average bankfull width of 1.6 m, and substrate dominated by gravels and fines (Lotic 2015a). The riparian vegetation was mainly shrubs. Immediately upstream from the spillway is the lower settling pond (8,745 m²), where the substrate is fines and supports macrophyte growth in the first two meters of littoral zone. Calcite was observed in the pond at low levels (CI value: 0.80). Upstream from the lower settling pond, the creek extends for about 2.75 km to the Lower Valley Road, where two additional settling ponds are located. In the surveyed section of the creek, a step-pool morphology was present with a very steep slope (42%) and a mean bankfull width of 2.21 m with some calcite present (CI range: 0.51 to 0.98). The riparian vegetation upstream from the spillway was mainly grasses and sedges (Lotic 2015a).

Fish presence was unknown for Six Mile Creek until the baseline assessment was completed in June 2013 and 2014 (Lotic 2015a). Fish species that were present in the lowest section of Six Mile Creek downstream from the concrete spillway included mountain whitefish, longnose dace, longnose sucker, redbelt shiner, westslope cutthroat trout, eastern brook trout, and bull trout. There was good cover (60%), provided primarily by overhanging vegetation, with some small woody debris. There were extensive areas of suitable spawning gravel for salmonids in the riffles and glides, although no fish were observed spawning during the surveys. Six Mile Creek had habitat potentially suitable for all life-stages of fish, including potential overwintering (Lotic 2015a). However, the overall small size of the stream makes it most suitable for small size classes. Low gradient and moderate velocity flows over riffles and glides provided excellent rearing habitat for fry and juvenile salmonids and multiple life stages of longnose suckers and longnose dace. It may provide refuge for Elk River fish during freshet but high water temperatures in the

summer ($>17.0^{\circ}\text{C}$) preclude use by salmonids at that time of year. No fish are present upstream from the spillway (Lotic 2015a).



Photo 7.3: Channel habitat in lower Six-Mile Creek (Lotic 2015a).

An olive-sided flycatcher and spotted sandpiper were observed near the mouth of Six-Mile Creek basin in 2013 (Golder 2015b).

7.3 Balmer Creek

About 24% of the total catchment area of Balmer Creek has been disturbed by mining activities, but no additional mine disturbances are planned for the near or far future (Table 7.1). Of the total stream length of 3.2 km, 19% is connected to the main stem whereas the remainder is fragmented (Table 7.3). The catchment has also been disturbed by forestry activities and roads (Table 7.1).

No data were available for water quality, benthic invertebrate community, tissue selenium concentrations, or calcite (Table 7.2).

A fish and fish habitat survey was conducted in June 2014 (Lotic 2015a). The most downstream portion of Balmer Creek extends from the confluence with the Elk River 280 m upstream to a significant gradient change. A 100-m section was characterized as riffle-pool morphology, with average gradient of 1.2%, average bankfull width of 1.92 m, and substrate dominated by fines and gravel. Farther upstream, the gradient increased to 34% with a step-pool morphology and a substrate dominated by boulders and cobble. Riparian vegetation was predominantly coniferous trees downstream and mixed forest upstream (Lotic 2015a).

Fish species that were captured or observed in lower Balmer Creek included mountain whitefish, westslope cutthroat trout, bull trout, and eastern brook trout (Lotic 2015a). Overhanging vegetation, small woody debris, and undercut banks provided a fair amount of cover (10%). There were a few small pockets of suitably sized gravel for westslope cutthroat trout spawning in the lower portion of the creek, and no suitable gravels for bull trout spawning (Lotic 2015a). Lower Balmer Creek provides good habitat for fry and juveniles but reduced overwintering potential. No fish were captured or observed upstream from the steep gradient change.

Olive-sided flycatchers were observed near Balmer Creek in 2013 (Golder 2015b).

7.4 Fennelon Creek

Fennelon Creek is located about 4.75 km north of Sparwood. About 47% of the total catchment area of Fennelon Creek has been disturbed by mining activities (Table 7.1). Mining is currently occurring within the permit boundary and will continue into the near and far future. All of the total stream length of 3 km is connected to the main stem (Table 7.3). The catchment has also been disturbed by forestry activities and roads (Table 7.1).

No data were available for water quality, benthic invertebrate community, tissue selenium concentrations (Table 7.2).

Fish and fish habitat inventories were conducted in June 2013 and 2014, as well as an overwintering survey in March 2013 (Lotic 2015a). The surveyed area had step-pool morphology with average gradient of 35%, bankfull width of 1.7 m, and substrate of cobble and gravel. Water depth was shallow, being only 0.1 m within the deepest pool observed in June 2013. No calcite deposition was observed. The riparian vegetation was mixed forest (Lotic 2015a).

A 2-m high waterfall located at the confluence with the Elk River represents a barrier to upstream fish passage (Lotic 2015a; Photo 7.4). About 30 m upstream from the Elk River confluence was another 4-m bedrock waterfall that is a second barrier to upstream fish

migration. No fish were captured or observed between the barriers in 2013 or 2014. Cover provided by overhanging vegetation and small woody debris was poor. There was no suitable spawning habitat for westslope cutthroat trout or bull trout due to inadequate substrate size and steep gradient. The overwintering survey completed in March 2013 found no overwintering potential, as the creek was frozen to the substrate, with no flowing water. Therefore, Fennelon Creek is considered non-fish bearing (Lotic 2015a).



Photo 7.4: View of 2 m tall waterfall at the confluence of Fennelon Creek with the Elk River (Lotic 2015a).

7.5 Goddard Creek

About 64% of the total catchment area of Goddard Creek has been disturbed by mining activities (Table 7.1). Mining is currently occurring within the permit boundary and will continue into the near and far future (Table 7.1). Of the total stream length of 8.1 km, 12% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 7.3). The catchment has also been disturbed by forestry activities and roads (Table 7.1).

Water quality has been characterized as “poor” based on Windward et al. (2015) but concentrations of selenium, nitrate and sulphate have shown decreasing trends (Zadjlick and Minnow 2013; Table 7.2; Figure 7.3). Benthic invertebrate community and calcite data were not reported by Windward et al. (2014).




Figure 7.3: Selenium concentration trend over time (2004-2013) in Goddard Creek at the confluence with the Elk River (EV_GC2).

The section of Goddard Creek closest to the confluence with the Elk River is lentic, and referred to as Goddard Marsh (4,872 m²; Lotic 2015a; Photo 7.5). Goddard Creek drains through multiple culverts into Goddard Marsh and there is permanent flow out of the marsh to the Elk River through a small channel that averages 1.84 m wide. Numerous studies of fish and aquatic habitat were conducted in Goddard Marsh between 2002 and 2013 (Lotic 2015a). The marsh measured 1.2 m deep in September 2013, with substrate dominated by fines. It is permanently inundated and supports abundant cattails and sedges, which grow throughout the marsh for up to about 40 m channel width, with mixed forest riparian vegetation bordering the cattail growth (Lotic 2015a).

Fish species captured or observed in Goddard Creek include longnose sucker, redbelly shiner, longnose dace, mountain whitefish, eastern brook trout, and westslope cutthroat trout (Lotic 2015a). The macrophytes, deep water and submerged woody debris provide abundant instream cover. Overhead cover was limited in the ponded water areas.



Photo 7.5: Goddard Marsh at the confluence of Goddard Creek and the Elk River (Lotic 2015a).

However, within the narrow, channelized sections of the marsh, cattails provided abundant overhead cover. There was no suitable spawning habitat for salmonids in the marsh due to the fine substrate, but the habitat is suitable for longnose sucker spawning. Goddard Marsh also provides good overwintering habitat, as adequate water depths were present and the marsh did not completely freeze (Lotic 2015a).

The upper end of the marsh ends at a constructed fish barrier, consisting of two v-notch concrete weirs, about 0.75 m high, with no plunge pools for passage over the weirs (Lotic 2015a). Upstream from the weirs there are a series of sediment ponds, a concrete/rip-rap spillway from the mid-settling pond, and the channel extending upslope from this to the headwaters. The channel had a step-pool morphology with an average gradient of 27%, and a substrate of boulders and fines. The reach is highly modified, with rip-rap lining the banks, so no bankfull width could be accurately measured. Riparian vegetation was largely absent, with some grass. There was no suitable spawning habitat for either westslope cutthroat trout or bull trout due to inadequate substrate size. This

section of Goddard Creek has little to no potential for fish use and is considered to be non-fish bearing (Lotic 2015a).

No species at risk were observed in breeding amphibian and bird surveys completed in Goddard Marsh in 2012 (Minnow 2014a; Table 7.3). Great blue herons nested near the marsh in the past, but moved their rookery about 1 km south prior to 2012 (Minnow 2014a, Golder 2015b). Columbia spotted frog and migratory birds, such as red-winged blackbirds have also been documented (Minnow 2014a).

7.6 Otto Creek

About 50% of the total catchment area of Otto Creek has been disturbed by mining activities and mining currently occurring within the permit boundary will continue into the near and far future (Table 7.1). Of the total stream length of 5.1 km, 25% is connected to the main stem whereas the remainder is fragmented or has been lost (Table 7.3). The creek has been highly modified by mining activities, including channelization between the rail line and tailings lagoons (Photo 7.6), and the creation of settling ponds (Lotic 2015a). The catchment has also been disturbed by forestry activities and roads (Table 7.1).



Photo 7.6: Channelized section of Otto Creek between the rail line and the tailings lagoons (Lotic 2015a).

Water quality and tissue selenium concentrations have been characterized as “good” whereas the benthic invertebrate community was characterized as “poor” (Windward et al. 2014; Table 7.2). A decreasing trend in nitrate concentrations and an increasing trend in sulphate concentrations were reported (Zadjlick and Minnow 2013; Table 7.2).

Surveys were completed in 2001, 2013 and 2014 (Lotic 2015a). The most downstream reach of the creek (total of 145 m) was characterized by riffle-pool morphology, with an average gradient of 1.9%, an average bankfull width of 3.4 m, and substrate was dominated by fines and gravel. Calcite was observed at low levels (CI: 0.30). Upstream from Highway 43 is a 900-m reach that includes a series of ponds/wetlands confined between the EVO rail loop and tailing lagoons, and separated by beaver dams. There were three wetland cells at the downstream end and four settling pond cells at the upstream end. Vegetation was dominated by coniferous trees and grasses.

Otto Creek has been differentiated into three reaches, two which are considered fish-bearing, whereas the third (Cossarini Creek) is considered non-fish bearing (Lotic 2015a). Fish species that were captured or observed in Otto Creek are longnose sucker, longnose dace, redbelly shiner, westslope cutthroat trout, bull trout, eastern brook trout, and mountain whitefish (Lotic 2015a). Lower Otto Creek had minimal spawning potential for salmonids due to substrate being either too fine or too coarse. No westslope cutthroat trout redds were observed. During the fall spawning survey, no bull trout or eastern brook trout redds were observed (Lotic 2015a). There was good cover provided by overhanging vegetation, and poor amounts of cover (10%) provided by both small and large woody debris. This reach provides habitat most suitable to fish in the size classes of juvenile salmonids (Lotic 2015a). The longnose dace fit that size range and do occupy similar habitats.

Upstream movement into the wetlands areas from the most downstream section is limited by the CPR rail culvert, however it is not a permanent barrier to upstream fish migration. There was no suitable spawning habitat for westslope cutthroat trout, but there was potentially suitable spawning habitat for longnose sucker and longnose dace along the margins of the ponds. The ponds would provide suitable holding and feeding water for adult sucker and dace (Lotic 2015a). A vegetated berm separates Otto Creek from Cossarini Creek, with a three-inch-diameter pipe running through the berm into the most upstream pond of Otto Creek. No fish were captured by electrofishing and minnow traps in Cossarini Creek (Lotic 2015a).

Western toad was the only species at risk identified in breeding amphibian and bird surveys completed in Otto Creek in 2012 (Minnow 2014a; Table 7.3).

7.7 Transmission Creek

Transmission Creek has not been disturbed by mining and is not expected to be influenced in the future (Table 7.1; Photo 7.7). All of the total stream length of 25.3 km is connected to the main stem (Table 7.3). The catchment has been disturbed by forestry activities and roads (Table 7.1).



Photo 7.7: Typical large woody debris/sediment wedge acting as a temporary barrier on Transmission Creek (Golder 2015).

Water quality, calcite, and the benthic invertebrate community were not evaluated by Windward et al. (2014) (Table 7.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). Tissue selenium concentrations were characterized as “good”.

A reconnaissance-level fish and fish habitat survey was completed for 100-m sections of the lower and upper creek in July 2012 and 2013 (Golder 2015a). The first surveyed area was located just upstream from Highway #3 (100 m section; Golder 2015a). The mean gradient was 9% and it was characterized by a cascade-pool morphology. The dominant substrates were cobble and gravel and the average bankfull channel width was 7.43 m. In

the upstream section of the creek, the mean gradient was 24% and the reach was characterized as step-pool morphology. Substrate was dominated by cobbles and boulders with some gravel present as well (Golder 2015a).

Both westslope cutthroat trout and eastern brook trout have been captured or observed in Transmission Creek (Golder 2015a). Only the downstream reach is fish-bearing. Overhanging vegetation was the dominant instream cover with trace amounts of small woody debris, boulders, and undercut banks. Both adult and juvenile trout were captured, indicating that rearing, holding, and feeding habitat is present. Upstream from the first survey location, the creek branches into the main branch and an unnamed tributary, both of which are considered non-fish bearing due to a series of temporary and permanent barriers to upstream migration (Golder 2015a; Photo 7.7). For example, over the first 460 m upstream from the first survey area were a series of woody debris and sediment wedge features that created falls ranging from 0.7 to 1.8 m. In addition, a temporary culvert barrier was noted, which had a 1.2-m outlet drop into a 0.2 m plunge pool and was considered impassible to fish (Golder 2015a). The first feature considered to be a permanent barrier to upstream fish migration was a 24% cascade, about 11 m long, located 56 m upstream from the culvert and 420 m downstream from the confluence with an unnamed tributary (Golder 2015a).

7.8 Mine Creek (Hosmer Creek)

Mine Creek has not been disturbed by mining and is not expected to be influenced in the future (Table 7.1). Mine Creek is a major tributary to Hosmer Creek, located close to the confluence of Hosmer Creek with the Elk River. All of the total stream length of 81.6 km is connected to the main stem (Table 7.3). The catchment has been disturbed by forestry activities and roads (Table 7.1).

Water quality, calcite, and the benthic invertebrate community were not evaluated by Windward et al. (2014) (Table 7.2), but are likely comparable to other regional streams that have not been affected by mining (e.g., “good”). Tissue selenium concentrations were characterized as “good”.

Fish and fish habitat surveys were conducted in 2007, 2012, and 2013 (Golder 2015a). This first surveyed section (85 m) of Mine Creek was lentic habitat located about 4 km upstream from the confluence with Hosmer Creek (Golder 2015a; Photo 7.8). The area was dominated by a series of pools and low flow glide areas, with a mean gradient of only 0.3%. The dominant substrate was fines with some gravel, and the average bankfull width was 3.20 m. A culvert was noted within the site. A second surveyed area (100 m) was located 6.6 km upstream from the confluence with Hosmer Creek. This section was



Photo 7.8: Lentic area in Mine Creek, upstream of the confluence with Hosmer Creek (Golder 2015).

dominated by riffle habitat (83.8%), with cascade, glide, and pool habitat units also present, classifying it as riffle-pool morphology. Dominant substrates included cobble, gravel and boulder and the bankfull channel width was 3.3 m. A third area (100 m) was surveyed about 8.7 km upstream from the confluence with Hosmer Creek, which was classified as cascade-pool morphology, dominated by cascade habitat (70.9%), with glide and riffle habitat units also present. The dominant substrate was cobble, gravel and boulder and the bankfull width was 3.93 m. Two unnamed tributaries join Mine Creek in the northern portion of the catchment, about 7 and 8 km upstream from the confluence with Hosmer Creek (Golder 2015a). Riparian vegetation was dominated by shrubs and grasses in the downstream sections and mixed mature forest in the upstream sections. No off-channel habitat was observed (Golder 2015a).

Eastern brook trout, bull trout and westslope cutthroat trout have been captured and observed within Mine Creek (Golder 2015a). The presence of juvenile and adult eastern

brook trout indicates that the lentic areas on Mine Creek provide rearing and holding/feeding habitat. The shallow water depths at the time of the survey in August 2013 precludes the use of the area as overwintering habitat for adult fish species, but may provide adequate overwintering habitat for juvenile eastern brook trout. In the upstream lotic areas, the presence of juvenile and adult fish species indicates suitable rearing, holding, and feeding habitat are present. Trace amounts of suitable resident trout spawning gravels were observed. Gradient barriers limit fish access to headwaters (Golder 2015a).

8.0 REFERENCES

- Allan, J.H. 1987. Fisheries investigations in Line Creek in 1987. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Allan, J.H. 1991. Fisheries investigations in Line Creek in 1991. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Allan, J.H. 1993. Fisheries investigations in Line Creek in 1993. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Allan, J.H. 1994. Fisheries investigations in Line Creek and South Line Creek in 1994. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Allan, J.H. 1995. Fisheries investigations in Line Creek in 1995. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Allan, J.H. 1996. Fisheries Investigations in Henretta Creek and Fish Pond Creek in 1995. Prepared for Fording Coal Ltd. Prepared by Pisces Environmental Consulting Services Ltd, Red Deer, AB.
- Allan, J.H. 2000. Fisheries investigations in Line Creek in 2000. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Allan, J.H. 2001. Fisheries investigations in Line Creek in 2001. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.
- Amos, L., and J. Wright. 2000. 1999 Fording River fish and fish habitat monitoring program. Prepared for Fording Coal Ltd., Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. 46 p. + 8 app.
- Amos, L., J. Wright. 2002. Fish and habitat assessments associated with coal bed methane development in the Upper Fording River drainage. Prepared for EnCana Corporation, Calgary, AB. 2002.

- Arnett, T., and Berdusco, J. 2009. Line Creek Operations 2008 Aquatic Health Monitoring Program. Consultant report prepared for Elk Valley Coal Corporation - Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 53 pp + 8 appds
- Arnett, T., and M.D. Robinson. 2010. 2007 – 2009 Line Creek Aquatic Health Monitoring Summary. Prepared for Teck Coal Ltd. – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 72 pp + 8 appds.
- BCCDC (British Columbia Conservation Data Centre). 2012. Search tools for conservation status available at <http://www.env.gov.bc.ca/cdc/access.html>.
- BCFISB (British Columbia Fisheries Information Services Branch). 2001. Reconnaissance (1:20,000) Fish and Fish Habitat Inventory: Standards and Procedures. Prepared for the Resources Inventory Committee. April.
- BCMOE (British Columbia Ministry of Environment). 2014. Permit 107517, Under the Provisions of the *Environmental Management Act*, Teck Coal Limited. Issued November 19, 2014.
- BCMOE (British Columbia Ministry of Environment). 2015. Fish Inventory Data Queries (FIDQ) Tool. Website: <http://www.env.gov.bc.ca/fish/fidq/queries.html>
- BCMOF (British Columbia Ministry of Forests). 1996. Channel Assessment Procedures Guidebook. Available at: <http://www.for.gov.bc.ca/tasb/legsregs/fpc/fpcguide/chanfld/cfld-toc.htm>. Accessed September 2014.
- BC Research. 1981. Greenhills Surface Coal Mining Project. Stage II Environmental Assessment, Volume I-III. Prepared for Kaiser Resources Ltd.
- Berdusco J. 2003. Brownie Creek and Tributaries Followup Fisheries and RMA Classification. Prepared for Fording River Operations, Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. July. 9 p.
- Berdusco, J., T. Arnett, and J. Bisset. 2007. Line Creek Operations 2006 Fisheries Investigation. Consultant report prepared for Elk Valley Coal Corporation – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC.
- Berdusco, J., and T. Arnett. 2008. Line Creek Operations 2007 Aquatic Health Monitoring Program. Consultant report prepared for Elk Valley Coal Corporation – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 57 pp + 8 appds.

- CCME. 2001a. CCME water quality index 1.0. Technical report. Canadian water quality guidelines for the protection of aquatic life. Canadian Council of Ministers of the Environment, Winnipeg, MB.
- CCME. 2001b. CCME water quality index 1.0. User's manual. Canadian water quality guidelines for the protection of aquatic life. Canadian Council of Ministers of the Environment, Winnipeg, MB.
- CCME. 2007. Sediment quality index 1.0 calculator (SeQI) [online]. Canadian Council of Ministers of the Environment, Winnipeg, MB. Available from: http://www.ccme.ca/ourwork/water.html?category_id=103.
- Cope, S., C.J. Schwarz, J. Bisset, and A. Prince. 2014. Upper Fording River Westslope Cutthroat trout (*Oncorhynchus clarkii lewisii*) population assessment – telemetry project annual report: 2013 (Interim Report 2). Report Prepared for Teck Coal Limited, Calgary, AB. Report Prepared by Westslope Fisheries Ltd., Cranbrook, B.C. 108 p. + 2 app.
- Edeburn, A. 2003. Brownie Creek and Tributaries Habitat Assessment. Prepared for Greg Sword, Fording Coal Ltd. Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. 3 p.
- Environment Canada. 2013. How Much Habitat is Enough? Third Edition. Environment Canada, Toronto, Ontario.
- Fording Coal Limited. 1985. Kilmarnock Creek dragline mining proposal environmental impact assessment. Report prepared for British Columbia Mine Development Steering Committee. Prepared by Fording Coal Limited, 61 p.
- Fording Coal Limited. 1991. Fording Coal Limited Enhancement of the Upper Fording River (Above Confluence with Henretta Creek). Prepared by Fording Coal Limited, 21 p, 1 app.
- Golder Associates Ltd. 2014a. Fish and Fish Habitat Baseline Report. Submitted to Teck Coal Limited, Fording River Operations, Elkford, BC. Annex G of Fording River Operations Swift Project Environmental Assessment Certificate Application. November.
- Golder Associates Ltd. 2014b. Fording River Operation Swift Project: Geomorphic Overview Characterization and Review. Technical Memorandum from J. Wright, R. Atkins, and J. Seabrook, Golder Associates Ltd., to G. Sword and W. Franklin, Teck Coal Limited. November 17, 2014. (Appendix G.VIII of Golder 2014a).

Golder Associates Ltd. 2014c. Site Conditions. Annex D.5 of the Elk Valley Water Quality Plan. Teck Coal Limited. July.

Golder Associates Ltd. 2014d. Fording River Operations Swift Project Environmental Assessment Certificate Application.

Golder Associates Ltd.. 2015a. Coal Mountain Phase 2 Project. Annex I – Fish and Fish Habitat Baseline Report. Submitted to Teck Coal Limited. August.

Golder Associates Ltd.. 2015b. Elkview Operations Baldy Ridge Extension Project. Annex J – Wildlife and Wildlife Habitat Baseline Report. Submitted to Teck Coal Limited, Elkview Operations. October.

Golder Associates Ltd.. 2015c. Elkview Operations Baldy Ridge Extension Project. Annex N – Landscape and Ecosystems Baseline Report. Submitted to Teck Coal Limited, Elkview Operations. July.

Goltz C. 1997. Fisheries investigations in Line Creek in 1997. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

Goltz C. 1998. Fisheries investigations in Line Creek in 1998. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

Goltz C. 1999. Fisheries investigations in Line Creek in 1999. Consultant report prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

Interior Reforestation Co. Ltd. 1997. Fish and Fish Habitat Inventory for Select Tributaries of the Upper Flathead and Elk Rivers. Prepared for Cresbrook Forest Industries Ltd., Sparwood Division, Cranbrook, BC. September.

Interior Reforestation Co. Ltd.. 2003... Turnbull South Pit and Spoil development Application, Fish and Fish Habitat Assessment. Prepared for Elk Valley Coal Corporation.

Interior Reforestation Co. Ltd. 2006. Letter to Mr. G. Sword of Elk Valley Coal Corporation, Fording River Operations, dated January 27, 2006. Re: 2005 Clode Ponds fish salvage project.

Interior Reforestation 2011. Elk River Fish Distribution Report. Surveyed area AND1 in August 2010.

- Johnston, N.T., and P.A. Slaney. 1996. Fish Habitat Assessment Procedures. Watershed Restoration Tech. Circ. No. 8. Ministry of Environment, Lands and Parks and Ministry of Forests, Vancouver, BC.
- KNRC (Kootenay Natural Resource Consulting. 2007. Fish Stream Identification, Greenhills Creek. Prepared for D. Charest, Elk Valley Coal Corporation. July 30, 2007.
- Lotic Environmental. 2015a. Elkview Operations Baldy Ridge Extension Project - Fish and Fish Habitat Baseline Assessment. Prepared by Lotic Environmental Ltd. for Teck Coal Ltd. July.
- Lotic Environmental. 2015b. Greenhills Operations Cougar Pit Extension Project - Fish and Fish Habitat Baseline Assessment. Prepared by Lotic Environmental Ltd. for Teck Coal Ltd. July.
- MacKenzie, W.H.. and J.R. Moran. 2004. Wetlands of British Columbia: A Guide to Identification. Land Management Handbook 52, BC Ministry of Forestry, Research Branch, Victoria, BC.
- Mackillop, D. 2012. Kootenay-Boundary BEC Update. Unpublished draft report by Deb Mackillop, Regional Ecologist, Kootenay-Boundary Region (March 2012).
- Matrix Solutions Inc. 2015. Wildlife Baseline Report: Greenhills Operations Cougar Pit Extension Project. Prepared for Teck Coal Limited Greenhills Operations. July.
- Minnow Environmental Inc. 2003. Selenium study of lentic areas in the Elk Valley (draft). Prepared for Elk Valley Selenium task Force, Mississauga, ON.
- Minnow Environmental Inc. 2014a. 2012 Biological Monitoring Program for Coal Mines in the Elk River Valley, B.C. Report Prepared for Teck Coal Limited, Sparwood, BC. March. Project #2456.
- Minnow Environmental Inc. 2014b. 2013 Sediment Sampling Program for the Coal Mines in the Elk River Watershed, BC. Prepared for Teck Coal Limited, Sparwood, BC. July. Project #2494.
- Minnow Environmental Inc. 2015a. Design for the Regional Aquatic Effects Monitoring Program (RAEMP), Elk River Watershed. Prepared for Teck Coal Limited, Sparwood, B.C. March 2015. Project #2529.
- Minnow Environmental Inc. 2015b. Phased Study Design for the Tributary Evaluation Program. Prepared for Teck Coal Limited, Sparwood, BC. May. Project #2563.



- Minnow Environmental Inc., Golder Associates Ltd. and Nautilus Environmental Inc. 2015. Sediment Toxicity Supporting Study Design, Elk River Watershed, B.C. Prepared for Teck Coal Limited. March. Project #3565.
- Mount, C., S. Norris, R. Thompson, and D. Tesch. 2011. GIS Modelling of Fish Habitat and Road Crossings for the Prioritization of Culvert Assessment and Remediation. Streamline Watershed Management Bulletin 14:7-13
- Norris, S., and C. Mount. In Draft. Fish Passage GIS Analysis. Version 2. Methodology and Output Data Specifications. B.C. Ministry of Environment.
- NorthSouth Consultants Inc. 2008. BP Canada Mist Mountain Project 2008 Baseline Aquatics Report: Surface Water Quality, Fisheries and Aquatic Habitat Assessment. Prepared for: Matrix Solutions Inc.
- NorthSouth Consultants Inc. 2007. BP Canada Mist Mountain Project 2007 Baseline Aquatics Report: Surface Water Quality, Fisheries and Aquatic Habitat Assessment. Prepared for: Matrix Solutions Inc.
- Paine, Ledge and Associates, and Minnow Environmental Inc. 2011. Three Year (2007-2009) Evaluation of Selenium Concentrations and Loads in the Elk River Watershed, BC. Prepared for Teck Coal Limited. Sparwood, BC. February.
- RL&L Environmental Services Ltd. 1994. Review of Fisheries and Habitat Investigations on Thompson Creek. Prepared for Fording Coal Ltd., Greenhills Creek. October.
- RL&L Environmental Services Ltd. 1995. Memo to R. Jones from Rick Pattenden. Re: Swift Creek Fish Inventory. 31 January.
- Robinson, M.D., and R.J. MacDonald. 2014. Teck Coal Ltd – 2013 Calcite Monitoring Program. Prepared for Teck Coal Ltd by Lotic Environmental Ltd. 13 pp + appendices.
- Robinson, M.D. and R.J. MacDonald. 2015. Teck Coal Ltd – Elk Valley 2014 Calcite Monitoring Program. Prepared for Teck Coal Ltd by Lotic Environmental Ltd.
- Robinson, M.D. and R.J. MacDonald. 2016. Teck Coal Ltd – Elk Valley 2015 Calcite Monitoring Program. Prepared for Teck Coal Ltd by Lotic Environmental Ltd.
- Robinson, M. Fish Biologist. Lotic Environmental Inc. Cranbrook, BC. Phone: 250-421-7802.
- Robinson, M., and J. Wright. 2005. Line Creek Fisheries Investigation 2004 Program. Prepared for Elk Valley Coal Corporation – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC.

- Robinson, M.D., Goodbrand, A., and R.J. MacDonald. 2014. Multi-scale Characterization of Bull Trout Spawning Locations in Line Creek. Prepared for Teck Coal Ltd by Lotic Environmental Ltd. 25 pp.
- Teck. 2014. Elk Valley Water Quality Plan. Submitted to the British Columbia Minister of Environment for approval on July 22, 2014.
- Teck. 2015. Fording River Operations Swift Project Environmental Assessment Certificate Application. Issued November 2014, Revised January 2015.
- Teck Coal Limited. Undated. Coal Mountain Operations – Phase 2: Wildlife Baseline Report (DRAFT).
- Teck Coal Limited. 2011a. Line Creek Operations Phase II Project Environmental Assessment Certificate Application. Issued December 2011. Section B.2.2.3
- Teck Coal Limited. 2011b. Line Creek Operations Phase II Project Fish and Fish Habitat Baseline Report. Submitted to British Columbia Environmental Assessment Office, Victoria, BC. December 2011.
- Teck, Her Majesty the Queen in Right of Canada, and Her Majesty the Queen in Right of British Columbia. 2011. Memorandum of Agreement for the Management of Select Coal Mining Impacts on Fish and Fish Habitat within the Elk Valley, BC.
- Tembec. 2005. Sustainable Forest Management Plan. Tembec Industries Inc. Forest Resource Management BC Division. Term: 2005-2010. Available at: http://forest.tembec.com/sites/forest.tembec.com/files/pdf/tembec_sfmp_final_sept_22_2005.pdf.
- Tschaplinski P.J., and R.G. Pike. 2010. Riparian management and effects on function. Pages 479-525 In R.G. Pike, T.E. Redding, R.D. Moore, R.D. Winkler and K.D. Bladon. Compendium of forest hydrology and geomorphology in British Columbia. BC Min. For. Range, For. Sci. Prog., Victoria, BC and FORREX Forum for Research and Extension in Natural Resources, Kamloops, BC. Land Manag. Handb. 66. Available at: www.for.gov.bc.ca/hfd/pubs/Docs/Lmh/Lmh66.htm.
- Windward Environmental LLC. 2014. Screening-Level Dietary Ecological Risk Assessment of Metals in the Elk River Watershed and Lake Koocanusa. Prepared for Teck Coal Ltd.
- Windward Environmental LLC, Minnow Environmental Inc. and CH2M HILL Ltd. 2014. Elk River Watershed and Lake Koocanusa British Columbia Aquatic Environment Synthesis Report. Prepared for Teck Coal Ltd. October.

Zajdlik & Associates, Inc and Minnow Environmental Inc. 2013. Three-year (2010-2012) evaluation of selenium, cadmium, sulphate and nitrate concentrations and loads in the Elk River Watershed, BC. (DRAFT). October.

APPENDIX A

TRIBUTARIES TO UPPER FORDING RIVER (MU1)






- | | |
|-------------------|--|
| Bridge | 10% Gradient Barrier |
| Culvert | 25% Gradient Barrier |
| Control Structure | Water Quality Monitoring |
| Rock Drain | Teck Coal Permit Boundary |
| Weir | Pre-mining Stream |
| Spillway | Catchment Area as of Sept. 2015 |
| Waterfall | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Fish Habitat Connected to Mainstem |
| Settling Ponds | Fish Habitat Fragmented (Potentially Reversible) |
| | Fish Habitat Fragmented (Permanently Isolated) |
| | Fish Habitat Permanently Altered or Destroyed |
| | Non-Fish Bearing |

**Figure A.1 DRAFT
Upstream Fording River**


0 375 750 1,500
Meters

Teck

1:34,000





**Figure A.2 DRAFT
Henretta Creek**



Teck

1:50,000





- | | |
|-------------------|--|
| Rock Drain | 10% Gradient Barrier |
| Bridge | 25% Gradient Barrier |
| Settling Ponds | Water Quality Monitoring |
| Culvert | Teck Coal Permit Boundary |
| Control Structure | Pre-mining Stream |
| Rock Drain | Catchment Area as of Sept. 2015 |
| Weir | CWB Catchment Area - 1980's with Updates |
| Spillway | Fish Habitat Connected to Mainstem |
| Waterfall | Fish Habitat Fragmented (Potentially Reversible) |
| | Fish Habitat Fragmented (Permanently Isolated) |
| | Fish Habitat Permanently Altered or Destroyed |
| | Non-Fish Bearing |

**Figure A.3 DRAFT
Fish Pond Creek**

0 250 500 1,000
Meters


Teck

1:20,000





- | | | | |
|---------------------|----------------------------|--|--|
| ✕ Culvert | ✕ 10% Gradient Barrier | Rock Drain | — Fish Habitat Connected to Mainstem |
| ✕ Control Structure | ✕ 25% Gradient Barrier | □ Settling Ponds | — Fish Habitat Fragmented (Potentially Reversible) |
| ✕ Rock Drain | ● Water Quality Monitoring | □ Teck Coal Permit Boundary | — Fish Habitat Fragmented (Permanently Isolated) |
| ✕ Weir | □ Bridge | □ Catchment Area as of Sept. 2015 | — Fish Habitat Permanently Altered or Destroyed |
| ✕ Spillway | | □ CWB Catchment Area - 1980's with Updates | — Non-Fish Bearing |
| ✕ Waterfall | | | |

**Figure A.4 DRAFT
Clode Creek**



Teck

1:20,000





- | | |
|---------------------|--|
| □ Bridge | ✕ 10% Gradient Barrier |
| ✕ Culvert | ✕ 25% Gradient Barrier |
| ✕ Control Structure | ● Water Quality Monitoring |
| ✕ Rock Drain | □ Teck Coal Permit Boundary |
| ✕ Weir | — Pre-mining Stream |
| ✕ Spillway | — Catchment Area as of Sept. 2015 |
| ✕ Waterfall | — CWB Catchment Area - 1980's with Updates |
| ▨ Rock Drain | — Fish Habitat Connected to Mainstem |
| ▨ Settling Ponds | — Fish Habitat Fragmented (Potentially Reversible) |
| | — Fish Habitat Fragmented (Permanently Isolated) |
| | — Fish Habitat Permanently Altered or Destroyed |
| | — Non-Fish Bearing |

**Figure A.5 DRAFT
Lake Mountain Creek**

0 265 530 1,060
Meters

Teck

1:24,000





- | | | | |
|---------------------|----------------------------|--|--|
| ✕ Culvert | ● Water Quality Monitoring | ▭ Teck Coal Permit Boundary | — Fish Habitat Connected to Mainstem |
| ✕ Control Structure | □ Bridge | ▭ Settling Ponds | — Fish Habitat Fragmented (Potentially Reversible) |
| ✕ Rock Drain | ✕ 10% Gradient Barrier | ▭ Catchment Area as of Sept. 2015 | — Fish Habitat Fragmented (Permanently Isolated) |
| ✕ Weir | ✕ 25% Gradient Barrier | ▭ CWB Catchment Area - 1980's with Updates | — Fish Habitat Permanently Altered or Destroyed |
| ✕ Spillway | Rock Drain | | — Non-Fish Bearing |
| ✕ Waterfall | | | |

**Figure A.6 DRAFT
Kilmarnock Creek**

0 250 500 1,000
Meters

Teck

1:34,000





- | | |
|-------------------|--|
| Bridge | 10% Gradient Barrier |
| Culvert | 25% Gradient Barrier |
| Control Structure | Water Quality Monitoring |
| Rock Drain | Teck Coal Permit Boundary |
| Weir | Catchment Area as of Sept. 2015 |
| Spillway | CWB Catchment Area - 1980's with Updates |
| Waterfall | Fish Habitat Connected to Mainstem |
| Rock Drain | Fish Habitat Fragmented (Potentially Reversible) |
| Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| | Fish Habitat Permanently Altered or Destroyed |
| | Non-Fish Bearing |

**Figure A.7. DRAFT
Swift Creek**

0 250 500 1,000
Meters


Teck

1:20,000





- | | |
|--------------------------|--|
| Bridge | 10% Gradient Barrier |
| Water Quality Monitoring | 25% Gradient Barrier |
| Rock Drain | Settling Ponds |
| Culvert | Pre-mining Stream |
| Control Structure | Teck Coal Permit Boundary |
| Rock Drain | Catchment Area as of Sept. 2015 |
| Weir | CWB Catchment Area - 1980's with Updates |
| Spillway | Fish Habitat Connected to Mainstem |
| Waterfall | Fish Habitat Fragmented (Potentially Reversible) |
| | Fish Habitat Fragmented (Permanently Isolated) |
| | Fish Habitat Permanently Altered or Destroyed |
| | Non-Fish Bearing |

**Figure A.8 DRAFT
Cataract Creek**



Teck

1:20,000





- | | |
|----------------------------|--|
| □ Bridge | ✕ 10% Gradient Barrier |
| ● Water Quality Monitoring | ✕ 25% Gradient Barrier |
| ✕ Culvert | — Fish Habitat Connected to Mainstem |
| ✕ Control Structure | — Fish Habitat Fragmented (Potentially Reversible) |
| ✕ Rock Drain | — Fish Habitat Fragmented (Permanently Isolated) |
| ✕ Weir | — Fish Habitat Permanently Altered or Destroyed |
| ✕ Spillway | — Non-Fish Bearing |
| ✕ Waterfall | — Settling Ponds |
| — Rock Drain | — Pre-mining Stream |
| | — Teck Coal Permit Boundary |
| | — Catchment Area as of Sept. 2015 |
| | — CWB Catchment Area - 1980's with Updates |

**Figure A.9 DRAFT
Porter Creek**

0 225 450 900
Meters

Teck

1:20,000





- | | |
|---|---|
| <ul style="list-style-type: none">BridgeRock DrainCulvertControl StructureRock DrainWeirSpillwayWaterfallSettling Ponds | <ul style="list-style-type: none">10% Gradient Barrier25% Gradient BarrierWater Quality MonitoringTeck Coal Permit BoundaryPre-mining StreamCatchment Area as of Sept. 2015CWB Catchment Area - 1980's with UpdatesFish Habitat Connected to MainstemFish Habitat Fragmented (Potentially Reversible)Fish Habitat Fragmented (Permanently Isolated)Fish Habitat Permanently Altered or DestroyedNon-Fish Bearing |
|---|---|

**Figure A.10 DRAFT
Chauncey Creek**

0 375 750 1,500
Meters

Teck

1:34,000





- | | |
|---------------------|--|
| □ Bridge | ✕ 10% Gradient Barrier |
| ✕ Culvert | ✕ 25% Gradient Barrier |
| ✕ Control Structure | □ Teck Coal Permit Boundary |
| ✕ Rock Drain | — Pre-mining Stream |
| ✕ Weir | ● Water Quality Monitoring |
| ✕ Spillway | □ Catchment Area as of Sept. 2015 |
| ✕ Waterfall | □ CWB Catchment Area - 1980's with Updates |
| □ Settling Ponds | — Fish Habitat Connected to Mainstem |
| ▨ Rock Drain | — Fish Habitat Fragmented (Potentially Reversible) |
| | — Fish Habitat Fragmented (Permanently Isolated) |
| | — Fish Habitat Permanently Altered or Destroyed |
| | — Non-Fish Bearing |

**Figure A.11 DRAFT
Ewin Creek**


0 500 1,000 2,000
Meters



Teck

1:50,000



- | | |
|--------------------------|--|
| Bridge | 10% Gradient Barrier |
| Water Quality Monitoring | 25% Gradient Barrier |
| Culvert | Settling Ponds |
| Control Structure | Catchment Area as of Sept. 2015 |
| Rock Drain | CWB Catchment Area - 1980's with Updates |
| Weir | Fish Habitat Connected to Mainstem |
| Spillway | Fish Habitat Fragmented (Potentially Reversible) |
| Waterfall | Fish Habitat Fragmented (Permanently Isolated) |
| Rock Drain | Fish Habitat Permanently Altered or Destroyed |
| | Non-Fish Bearing |
| | Teck Coal Permit Boundary |
| | Future Mine Development Line Creek Phase |





- | | | | |
|--|-------------------|--|--|
| | Rock Drain | | 10% Gradient Barrier |
| | Settling Ponds | | 25% Gradient Barrier |
| | Bridge | | Water Quality Monitoring |
| | Culvert | | Teck Coal Permit Boundary |
| | Control Structure | | Pre-mining Stream |
| | Rock Drain | | Catchment Area as of Sept. 2015 |
| | Weir | | CWB Catchment Area - 1980's with Updates |
| | Spillway | | Fish Habitat Connected to Mainstem |
| | Waterfall | | Fish Habitat Fragmented (Potentially Reversible) |
| | | | Fish Habitat Fragmented (Permanently Isolated) |
| | | | Fish Habitat Permanently Altered or Destroyed |
| | | | Non-Fish Bearing |

**Figure A.13 DRAFT
Greenhills Creek**





0 375 750 1,500
Meters



Teck

1:34,000


APPENDIX B

TRIBUTARIES TO LOWER FORDING RIVER (MU2)





- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Pre-mining Stream |
| Control Structure | 25% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Rock Drain | Rock Drain | CWB Catchment Area - 1980's with Updates |
| Weir | Settling Ponds | Fish Habitat Connected to Mainstem |
| Spillway | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Waterfall | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| | | Non-Fish Bearing |





Teck

1:85,000

APPENDIX C

TRIBUTARIES TO UPPER ELK RIVER (MU3)




- Bridge
- Water Quality Monitoring
- Culvert
- Control Structure
- Rock Drain
- Weir
- Spillway
- Waterfall

- 10% Gradient Barrier
- 25% Gradient Barrier
- Rock Drain
- WaterNetwork
- Settling Ponds
- Realigned channel
- Teck Coal Permit Boundary
- Pre-mining Stream

- Catchment Area as of Sept. 2015
- CWB Catchment Area - 1980's with Updates
- Fish Habitat Connected to Mainstem
- Fish Habitat Fragmented (Potentially Reversible)
- Fish Habitat Fragmented (Permanently Isolated)
- Fish Habitat Permanently Altered or Destroyed
- Non-Fish Bearing



Figure C.1 DRAFT

Br F Creek



Teck


1:20,000



- | | | |
|--------------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Water Quality Monitoring | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Culvert | Rock Drain | Fish Habitat Connected to Mainstem |
| Control Structure | WaterNetwork | Fish Habitat Fragmented (Potentially Reversible) |
| Rock Drain | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Weir | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Spillway | Teck Coal Permit Boundary | Non-Fish Bearing |
| Waterfall | Pre-mining Stream | |



Figure C.10 DRAFT

Thompson Creek



Teck


1:20,000



- | | | |
|--------------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Water Quality Monitoring | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Culvert | Rock Drain | Fish Habitat Connected to Mainstem |
| Control Structure | WaterNetwork | Fish Habitat Fragmented (Potentially Reversible) |
| Rock Drain | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Weir | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Spillway | Teck Coal Permit Boundary | Non-Fish Bearing |
| Waterfall | Pre-mining Stream | |



Figure C.2 DRAFT

Wolf Creek



Teck


1:20,000



- | | | | |
|--------------|----------------------------|-----------------------------|--|
| □ Bridge | ● Water Quality Monitoring | ✕ 10% Gradient Barrier | □ Catchment Area as of Sept. 2015 |
| ✕ Culvert | ✕ Control Structure | ✕ 25% Gradient Barrier | □ CWB Catchment Area - 1980's with Updates |
| ✕ Rock Drain | ✕ Rock Drain | ✕ Settling Ponds | □ Fish Habitat Connected to Mainstem |
| ✕ Weir | ✕ Spillway | ✕ Teck Coal Permit Boundary | □ Fish Habitat Fragmented (Potentially Reversible) |
| ✕ Waterfall | | ✕ Pre-mining Stream | □ Fish Habitat Fragmented (Permanently Isolated) |
| | | | □ Fish Habitat Permanently Altered or Destroyed |
| | | | □ Non-Fish Bearing |



Figure C.4 DRAFT

Wade Creek



Teck


1:20,000



- | | | |
|--------------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Water Quality Monitoring | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Culvert | Rock Drain | Fish Habitat Connected to Mainstem |
| Control Structure | WaterNetwork | Fish Habitat Fragmented (Potentially Reversible) |
| Rock Drain | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Weir | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Spillway | Teck Coal Permit Boundary | Non-Fish Bearing |
| Waterfall | Pre-mining Stream | |
| | MapbookInset | |



Figure C.5 DRAFT

Cougar Creek



Teck

1:20,000




- Bridge
- Water Quality Monitoring
- Culvert
- Control Structure
- Rock Drain
- Weir
- Spillway
- Waterfall

- 10% Gradient Barrier
- 25% Gradient Barrier
- Rock Drain
- WaterNetwork
- Settling Ponds
- Realigned channel
- Teck Coal Permit Boundary
- Pre-mining Stream

- Catchment Area as of Sept. 2015
- CWB Catchment Area - 1980's with Updates
- Fish Habitat Connected to Mainstem
- Fish Habitat Fragmented (Potentially Reversible)
- Fish Habitat Fragmented (Permanently Isolated)
- Fish Habitat Permanently Altered or Destroyed
- Non-Fish Bearing





Figure C.6 DRAFT



No-Name Creek




Teck



1:20,000





- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |






- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |

Figure C.9. DRAFT

Wolfram Creek





Teck

1:20,000

APPENDIX D


TRIBUTARIES TO MICHEL CREEK (MU4)



- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Bridge | Fish Habitat Connected to Mainstem |
| Weir | Rock Drain | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Settling Ponds | Fish Habitat Permanently Altered or Destroyed |
| | Realigned channel | Non-Fish Bearing |
| | Teck Coal Permit Boundary | |
| | Pre-mining Stream | |



Figure D.1 DRAFT

Michel Creek Upstream



Teck


1:50,000



- | | | |
|-------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Culvert | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Control Structure | Water Quality Monitoring | Fish Habitat Connected to Mainstem |
| Rock Drain | Rock Drain | Fish Habitat Fragmented (Potentially Reversible) |
| Weir | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Spillway | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Waterfall | Teck Coal Permit Boundary | Non-Fish Bearing |
| | Pre-mining Stream | |

Figure D.2

Corbin Creek



Teck

1:34,000









Figure D.3. DRAFT

Andy Good Creek



Teck


1:34,000



- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Water Quality Monitoring | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Bridge | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |



Figure D.4 DRAFT

Leach Creek



Teck


1:85,000



- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |



FigureD.5. DRAFT

Carbon Creek






Teck

1:20,000




- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |







- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Water Quality Monitoring | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Bridge | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |



Teck


1:50,000



- | | | |
|--------------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Water Quality Monitoring | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Culvert | Rock Drain | Fish Habitat Connected to Mainstem |
| Control Structure | WaterNetwork | Fish Habitat Fragmented (Potentially Reversible) |
| Rock Drain | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Weir | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Spillway | Teck Coal Permit Boundary | Non-Fish Bearing |
| Waterfall | Pre-mining Stream | |



Figure D.8 DRAFT

Unnamed Tributary to Michel



Teck

1:20,000



- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |

Figure D.9. DRAFT



Fir Creek

0 425 850 1,700

Meters

Teck


1:34,000



- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Pre-mining Stream |
| Control Structure | 25% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Rock Drain | Rock Drain | CWB Catchment Area - 1980's with Updates |
| Weir | Settling Ponds | Fish Habitat Connected to Mainstem |
| Spillway | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Waterfall | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| | | Non-Fish Bearing |

Figure D.10 DRAFT

Alexander Creek



Teck

1:85,000








Figure D.11

Erickson Creek



Teck


1:34,000



- | | | |
|--------------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Water Quality Monitoring | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Culvert | Rock Drain | Fish Habitat Connected to Mainstem |
| Control Structure | WaterNetwork | Fish Habitat Fragmented (Potentially Reversible) |
| Rock Drain | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Weir | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Spillway | Teck Coal Permit Boundary | Non-Fish Bearing |
| Waterfall | Pre-mining Stream | |





Figure D.12 DRAFT

South Pit Creek



Teck

1:20,000












Figure D.14 DRAFT





Gate Creek



Teck





1:20,000







APPENDIX E

TRIBUTARIES TO MIDDLE ELK RIVER (MU4/MU5)






- | | | |
|--------------------------|---------------------------|--|
| Bridge | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Water Quality Monitoring | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Culvert | Rock Drain | Fish Habitat Connected to Mainstem |
| Control Structure | WaterNetwork | Fish Habitat Fragmented (Potentially Reversible) |
| Rock Drain | Settling Ponds | Fish Habitat Fragmented (Permanently Isolated) |
| Weir | Realigned channel | Fish Habitat Permanently Altered or Destroyed |
| Spillway | Teck Coal Permit Boundary | Non-Fish Bearing |
| Waterfall | Pre-mining Stream | |





Figure E.2 DRAFT




Six Mile Creek





Teck

1:20,000








- | | | |
|---------------------|-----------------------------|--|
| ✕ Culvert | ✕ 10% Gradient Barrier | □ Catchment Area as of Sept. 2015 |
| ✕ Control Structure | ✕ 25% Gradient Barrier | □ CWB Catchment Area - 1980's with Updates |
| ✕ Rock Drain | □ Settling Ponds | — Fish Habitat Connected to Mainstem |
| ✕ Weir | □ Bridge | — Fish Habitat Fragmented (Potentially Reversible) |
| ✕ Spillway | ● Water Quality Monitoring | — Fish Habitat Fragmented (Permanently Isolated) |
| ✕ Waterfall | □ Teck Coal Permit Boundary | — Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | — Pre-mining Stream | — Non-Fish Bearing |




Figure E.5. DRAFT



Goddard Creek



Teck

1:24,000






- | | | |
|-------------------|---------------------------|--|
| Culvert | 10% Gradient Barrier | Catchment Area as of Sept. 2015 |
| Control Structure | 25% Gradient Barrier | CWB Catchment Area - 1980's with Updates |
| Rock Drain | Settling Ponds | Fish Habitat Connected to Mainstem |
| Weir | Bridge | Fish Habitat Fragmented (Potentially Reversible) |
| Spillway | Water Quality Monitoring | Fish Habitat Fragmented (Permanently Isolated) |
| Waterfall | Teck Coal Permit Boundary | Fish Habitat Permanently Altered or Destroyed |
| Rock Drain | Pre-mining Stream | Non-Fish Bearing |





FigureE.7. DRAFT

Transmission Creek



Teck

1:20,000



APPENDIX F
BIBLIOGRAPHIES

ANNOTATED BIBLIOGRAPHY OF LITERATURE RELEVANT TO ELK RIVER TRIBUTARIES

1. Aiello, J. 1980. The Rebirth of Corbin – The Byron Creek Collieries Story. The Canadian Mining and Metallurgical Bullentin. January.

- A brief history was presented of the mining activities in the Corbin Creek area on Coal Mountain by Byron Creek Collieries. Methods used to protect Corbin Creek and Michel Creek from mine-influenced water were described, including diversion ditches and settling ponds.

2. Allan, J.H. 1987. Fisheries Investigations in Line Creek in 1987. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- At the time of this report, the open pit mine located adjacent to Line Creek was operated by Crow's Nest Resources Limited. CNRL undertook investigations of the fish and fish habitat in Line Creek in 1983, 1986, and 1987 to evaluate the potential effects of mining. Fish presence and population sizes were estimated in five areas of Line Creek and the lower section of South Line Creek. Habitat utilization was evaluated by direct visual observation. The habitat inventory was a generalized estimate of the amount of each flow type, substrate type, and cover component recorded. Five species of fish were collected in the Line Creek study areas: bull trout, westslope cutthroat trout and mountain whitefish, and very rarely rainbow trout and eastern brook trout. Fish distribution in Line Creek is dependent on barriers and temperature regimes. Bull trout were the most common, and found throughout the watershed. Less is known about westslope cutthroat trout distribution but they were found in the upper reaches less often due to temperatures less than their optimal range. Mountain whitefish were limited to lower 3.7 km of Line creek due to a falls that was not passable by this species. The bull trout using Line Creek for spawning were part of a fluvial Elk River stock and are typically migratory, while the cutthroat trout population appeared to be a resident population. Mountain whitefish moved into Line Creek in the summer for feeding and emigrated to the Fording and Elk Rivers in the fall to spawn.
- Reviewed in Teck 2011b.

3. Allan, J.H. 1991. Fisheries investigations in Line Creek in 1991. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- In 1991, a quantitative physical habitat survey was completed in order to assess the potential impact of diverting Line Creek. The survey also included the downstream portion of South Line Creek, a major tributary to Line Creek. Bull trout spawning surveys were also conducted at the same time.

4. Allan, J.H. 1993. Fisheries investigations in Line Creek in 1993. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- In 1993, 1994, 1995, 1997, 1998, 1999, 2000 and 2001 (Reports 4-11), studies were carried out to investigate spawning populations of bull trout, rearing populations of bull

trout and the densities of westslope cutthroat trout in Line Creek, along with habitat surveys. Sections of Line Creek and South Line Creek near the mouth of Line Creek were altered to accommodate mine operations. A new main haul road crossing was constructed over Line Creek in 1990 and a new channel approximately 600 meters in length was created to the east of the original channel of Line Creek downstream of the haul road crossing. Flow from South Line Creek was diverted into the new channel in May of 1993 and the flow from Line Creek was introduced in August of 1993. A large, deep head pond was constructed between the haul road culverts and the mouth of South, Line Creek. In late 1990 an upper haul road crossing was constructed and use of a rock dump which has now buried approximately 3 km of Line Creek (approximately from stream km 8.4 to stream km 11.1) commenced. The rock dump will ultimately extend further up the valley of Line Creek. Population densities in Line Creek generally exceeded those from previous studies. The number of adult bull trout using Line Creek for spawning increased substantially, which was considered the most likely reason for the increase in population density. In 1995, following a substantial flood, the numbers of juvenile bull trout in Line Creek decreased significantly, however densities increased again in subsequent years.

5. Allan, J.H. 1994. Fisheries investigations in Line Creek and South Line Creek in 1994. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

6. Allan, J.H. 1995. Fisheries investigations in Line Creek in 1995. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

7. Goltz C. 1997. Fisheries investigations in Line Creek in 1997. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

8. Goltz C. 1998. Fisheries investigations in Line Creek in 1998. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

9. Goltz C. 1999. Fisheries investigations in Line Creek in 1999. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

10. Allan, J.H. 2000. Fisheries investigations in Line Creek in 2000. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

11. Allan, J.H. 2001. Fisheries investigations in Line Creek in 2001. Prepared for Luscar Ltd., Line Creek Mine, Sparwood, BC. Prepared by Pisces Environmental Consulting Services Ltd., Red Deer, AB.

- See description for report #4.

12. Allan, J.H. 1996. Fisheries Investigations in Henretta Creek and Fish Pond Creek in 1995. Prepared for Fording Coal Ltd. Prepared by Pisces Environmental Consulting Services Ltd, Red Deer, AB.

- This study estimated the fish densities in two artificially created pools in Henretta Creek, and also surveyed westslope cutthroat trout spawning and fry emergence/emigration in Fish Pond Creek. The Henretta Creek ponds were created to compensate for habitat losses incurred during the creation of Henretta Pit and were intended to primarily serve as overwintering habitat. Pond WCT densities in fall 1995 were lower than in 1993, but higher densities in 1993 may have been an artifact of a barrier preventing upstream movement of spawners in that year. Fish Pond Creek is a groundwater fed terrace tributary to the Fording River which was enhanced to improve WCT production. Fish Pond Creek was used more intensively for spawning in 1995 compared to historical years, which the authors attributed to recent enhancement of spawning habitat.

13. Allan, J.H. 1997. Trends in Fish Population at the Leve (Line) Creek Mine. Proceedings of the 21st Annual British Columbia Mine Reclamation Symposium, Cranbrook, BC.

- Fish populations in Line Creek were investigated since the mid 1970s. Line Creek supported populations of bull trout, westslope cutthroat trout and mountain whitefish. This paper used the results of previous fisheries investigations and records of mine operations provided by Line Creek Resources Limited to assess changes, and the probable cause of changes, in the fish populations of the Line Creek system. The number of spawning bull trout increased by 78% from 1983-1989 to 1991-1994, and was attributed to changes in angling regulations. Juvenile densities of trout species decreased in Line Creek and South Line Creek in 1995 due to a flood in June of that year. The study concluded that adult spawning was not measurably affected by Line Creek mine operations, but had been affected by changes to angling regulations.

14. Allan, J.H. 2000. A Perspective on People, Coal and Fish in the Southern Rocky Mountains. Prepared for Fording Coal Ltd., Luscar Ltd. Line Creek Mine and Elkview Coal Corporation. Prepared by Pisces Environmental Consulting Services Ltd. Red Deer AB.

- This paper briefly outlines the progression of coal mining in the Elk Valley, from early exploration to settlement to the effects of development both economically and environmentally. Also discussed was the impact of overharvesting on recreational

fisheries, highlighting the Elk River, Michel Creek, and Alexander Creek as some of the most attractive fishing locations in the valley.

15. Amos, L. and J. Wright. 2000. 1999 Fording River Fish and Fish Habitat Monitoring Program. Prepared for Fording Coal Ltd., Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. 46 p. + 8 app.

- The studies completed in 1999 indicated the fish population in the Fording River and its tributaries, near the Fording River Operations, were in a healthy, stable condition. In-stream mitigation works, designed to offset the impact of mining activities, were functioning as required and were providing evidence of beneficial uses to the westslope cutthroat trout population. This report examined spawning and fry emergence assessment, fish population survey, stream temperature monitoring, seasonal fish use and benthic invertebrate surveys for Henretta Creek, Fish Pond Creek, Upper Fording River and Lower Fording River in 1998 and 1999.
- Reviewed in Golder 2014a
- Several figures in main body and appendices are missing in PDF.

16. Amos, L., A. Edenum, J. Wright. 2002. Fish and Fish Habitat Monitoring in the Lower Henretta and Fish Pond Creek Drainages. Prepared for Fording River Operations: Final Report: Interior Reforestation Co. Ltd. Cranbrook, BC.

- Consistent with studies in 2000 and 2001, Henretta Creek water temperatures showed a net increase both upstream and downstream of Henretta Lake from July 1 to October 25. Paired samples T – tests indicated that mean daily temperatures for July through September were significantly ($P < .05$) higher at the downstream site. Warming within Henretta Lake may have beneficial effects on aquatic productivity. Minor habitat improvements were evident but habitat complexity was not fully realized because instream enhancements were not yet complete. Concentrations of most metals in water were less than the provincial and federal guidelines. The vegetative cover along the entire length of the channel met or exceeded anticipated restoration goals.

17. Amos, L., J. Wright. 2002. Fish and Habitat Assessments Associated with Coal Bed Methane Development in the Upper Fording River Drainage. Prepared for EnCana Corporation, Calgary, AB. 2002.

- Riparian management classification, and presence/absence and distribution of fish, were evaluated in the vicinity of the Greenhills saddle. In the Fording and its tributaries (Lake Mountain Creek, Clode Creek, Fish Pond Creek, and two unnamed tributaries), habitat data were collected along with information on resource values at proposed stream crossings associated with the construction of a CBM sales gas pipeline. Fish Pond Creek rated highest in overall habitat quality of any of the streams surveyed due to extensive enhancements. The surveys identified several issues in regards to the planned pipeline alignment. Suggestions for mitigation included re-alignment of the pipeline path in order to minimize the number of stream crossings, reduce the impact on already sparse forests and maintenance of valued habitat components.
- PDF missing several figures and the appendices.

18. Amos, L., J. Wright, A. Edeburn. 2003. Fish and Fish Habitat Monitoring in Henretta Lake and Reclaimed Channel. Prepared by: Interior Reforestation Co. Ltd. Prepared for: Fording Coal Ltd. Elkford, BC. March.

- This primary goal of this study was to ensure both short and long-term Fording Coal Limited program goals and objectives are met in a practical, cost-effective and scientifically defensible manner. This will allow for informed management decisions, identification of realistic alternatives and constraints, and further improvements to the biological and physical communities in the Lower Henretta Creek drainage. The program goals were to determine differences in thermal dynamics of Henretta Lake and downstream, investigate the potential for further enhancement structures, to determine if current modifications were meeting the requirements of fish and other aquatic resources. The recommended monitoring plan included riparian cover, water quality, habitat characteristics, and evaluation of westslope cutthroat trout populations, spawning potential, and presence of young-of-the-year. Temperature, water chemistry and low oxygen conditions were investigated, along with biological sampling and physical channel attributes. An increase in temperature downstream of Henretta Lake was reported but temperatures were still low compared to other creeks within geographic proximity. Although sufficient spawning habitat is present within the creek, higher temperatures are preferred by spawning WCT.
- Reviewed in Golder 2014a as IR 2003.

19. Arnett, T and J. Berdusco. 2009. 2008 Aquatic Health Monitoring Program - Line Creek Operations Annual Report. Prepared for Teck Coal Ltd. – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 53 pp + 8 appds.

- Annual results of a 3-year study (2007-2009) that is described under study #20.

20. Arnett, T. and M.D. Robinson. 2010. 2007 – 2009 Line Creek Aquatic Health Monitoring Summary. Prepared for Teck Coal Ltd. – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 72 pp + 8 appds.

- Evaluation of bedload sediment; periphyton (including the effects of water quality on productivity and community composition); benthic invertebrates; fish and fish habitat; and bull trout spawning in Line Creek. The study reported coarser bedload, increased nitrate concentrations, increased periphyton biomass and benthic invertebrate densities and fewer sensitive benthic invertebrate species (i.e. percent Ephemeroptera, Plecoptera and Trichoptera - %EPT) downstream of mining compared to upstream.
- Reviewed in Teck 2011b as Interior Reforestation 2010b.

21. British Columbia Department of Environment. 1976. Kootenay Air and Water Quality Study. Phase I. Assessment of Information Available to the End of 1974. Water Quality in Region 2, The Elk River Basin. February. File No. 0322512-1.

- Available data were not always harmonized with exploration activity, effluent analyses, river flow regime, and biological sampling so authors found it difficult in some cases to distinguish between effects of natural run-off and man-made disturbances (chiefly coal exploration and mining). Water quality parameters that most often exceeded recognized standards were suspended solids and turbidity, particularly during spring runoff. "There

was no evidence of acid mine drainage or of the discharge of toxic materials in significant amounts. These results were confirmed by a limited amount of aquatic biology data. The data indicated the presence of a relatively undisturbed population of invertebrates and a good fishery." Water data presented for some areas of Fording River, Michel Creek, Elk River and effluents from some mine-related (settling?) ponds.

22. BCMOE (British Columbia Ministry of Environment). 1978. Kootenay Air and Water Quality Study Phase II. Water Quality in the Elk and Flathead River Basins. Water Investigations Branch. October. File No. 0322512-1.

- This report finalized the study of water quality in the Elk and Flathead River basins. Data were collected between 1975 and 1976. The study identified suspended sediment as one of the major contributors to decreased water quality, which in turn disrupted invertebrate populations in affected areas. The study area included sites in the Fording River Basin, the Upper Elk River Basin, the Michel Creek Basin, the Lower Elk River Basin and the Flathead River Basin.
- PDF missing sections 4.6 to 9.

23. B.C. Research. 1981. Greenhills Surface Coal Mining Project. Stage II Environmental Assessment, Volume I-III. Prepared for Kaiser Resources Ltd.

- The Stage II Environmental Assessment provided a detailed description of the Greenhills coal project and detailed biophysical information to fulfill the Stage II requirements pursuant to the guidelines for coal development. The plan that was in place at the time of the study was for the production of 1.8 million tonnes of clean coal for a project life of 20 years. The assessment determined that the open pit mine may cause significant changes in flow in some or all of the streams originating in the project area, however would have little impact on the Fording and Elk Rivers. Greenhills Creek and other small tributaries would experience an increase in total suspended solids load. A settling pond was planned for Greenhills Creek to minimize this impact. Dissolved solids and nitrogen levels were also expected to increase. Fisheries resources in the Fording and Elk Rivers were not expected to be affected by the project, however the settling pond in Greenhills creek was expected to reduce the amount of spawning and rearing habitat available for westslope cutthroat trout in that creek.

24. Berdusco J. 2003. Brownie Creek and Tributaries Followup Fisheries and RMA Classification. Prepared for Fording River Operations, Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. July. 9 p.

- This report summarized a study conducted in response to recommendations following an overwintering study the previous year to assess fish presence in the creek. No fish were captured in Brownie Creek, nor in a tributary to Brownie Creek, despite electrofishing and angling efforts at all sites.
- PDF missing several figures.

25. Berdusco, J. and J. Wright. 2005. Fish Capture and Relocation in Line Creek Upstream of the Rock Drain, Prepared for Elk Valley Coal Line Creek Operations, Prepared By: Interior Reforestation.

- This report described a catch and relocation project initiated in July 2005 to relocate fish (bull trout and westslope cutthroat trout) from above the rock drain to downstream of the rock drain in Line Creek. Angling and electrofishing were used to capture the fish. The authors concluded that the relocation effort was successful. A total of 18 bull trout and 21 westslope cutthroat trout were relocated.
- Reviewed in Teck 2011b as Interior Reforestation 2005a.

26. Berdusco, J.R. and J. Wright. 2006. Fish and Fish Habitat Monitoring in the Henretta Creek Reclaimed Channel (2005). Prepared for Elk Valley Coal Corporation – Fording River Operations, Elkford, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, B.C. 22pp + 6 appendices

- The 2005 fish and fish habitat monitoring in Henretta Lake and the Henretta Creek Reclaimed Channel represented the sixth year of a seven year monitoring program required by the C-3 Reclamation Permit issued to Elk Valley Coal Corporation's Fording River Operations (FRO). Consistent with all reports completed previously, no Westslope cutthroat trout (WCT) spawning pairs or redds were observed in a spawning assessment conducted over the length of the reclaimed channel. As with previous reports, the water temperatures measured upstream and downstream of Henretta Lake were significantly different. No accurate westslope cutthroat trout population estimate was obtained through mark-recapture procedures as no tagged fish were recaptured. The detailed fish habitat assessments completed in 2005, in conjunction with a GPS survey of the thalweg position and habitat units, photo point comparison and overview assessment indicate that the complexity and quantity of habitat, riparian vegetation, and channel stability is steadily improving.

27. Berdusco, J., T. Arnett, and J. Bisset. 2007. Line Creek Operations 2006 Fisheries Investigation. Prepared for Elk Valley Coal Corporation – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC.

- This study focused on completion of population density estimates at two index sites, undertaking an assessment of bull trout spawning activity in 8.3 km of the Line Creek mainstem and 1.0 km of South Line Creek, and an analysis of recommended changes implemented during the field program of that year. Conclusions supported the need for careful identification and selection of environmental and biological indicators that provide measurable attributes of Line Creek. Trends seen in Line Creek are generally positive, and changes to the monitoring program were made to provide a better indication of Line Creek's overall ecosystem health, and to identify any relationships to mining activity.
- Reviewed in Teck 2011b as Interior Reforestation 2007.

28. Berdusco, J. and T. Arnett. 2008. Line Creek Operations 2007 Aquatic Health Monitoring Program. Prepared for Elk Valley Coal Corporation – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 57 pp + 8 appds.

- Annual results of a 3-year study (2007-2009) that is described under study #20.

29. Arnett, T and J. Berdusco. 2009. 2008 Aquatic Health Monitoring Program - Line Creek Operations Annual Report. Prepared for Teck Coal Ltd. – Line Creek Operations,

Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. 53 pp + 8 appds.

- Annual results of a 3-year study (2007-2009) that is described under study #20.

30. Carscadden, K. and S. Rogers. 2011. Genetic analysis of Westslope Cutthroat Trout from the Upper Fording River Drainage, British Columbia. Report Prepared for Golder Associates Ltd., Calagary, AB.

- Westslope cutthroat trout were caught from Swift Creek and Dry Creek and adipose fins were collected in order to determine whether any hybridization between WCT and rainbow trout had occurred. Microsatellite fingerprint analysis concluded that WCT in Swift Creek and Dry Creek were genetically pure and showed no signs of hybridization with rainbow trout.
- Reviewed in Teck 2011b.

31. Chapman, P.M., R. Berdusco and R. Jones. 2008. Selenium Investigations in the Elk River valley, B.C. – 2008 Update. Report prepared for Elk Valley Selenium Task Force. Prepared by Golder Associates Ltd., Burnaby, B.C.

- This report evaluated the current information on selenium concentrations water in the Elk Valley watershed. Se has been measured in benthic invertebrates, fish muscle and bird eggs but although levels are elevated, they have not increased significantly over several years at the time of this report. Using Line Creek as an example, concentrations detected downstream of mining operations have not appeared to have an impact on fish and water bird viability. Despite this, significant effort has been invested in predicting future Se loads, factors affecting cycling once it reaches the environment, and mitigation plans to manage the release of Se, including treatment options to reduce the load.

32. Edeburn, A. 2003. Brownie Creek and Tributaries Habitat Assessment. Prepared for Greg Sword, Fording Coal Ltd. Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. 3 p.

- This letter described a cursory fish habitat assessment completed in Brownie Creek and its tributaries on November 28, 2002. A key objective of this study was to determine fish habitat present, particularly overwintering suitability for the assumed fish species present, westslope cutthroat trout. Two sites were surveyed, one in the lower watershed and one in the upper watershed. The author recommended further biological sampling to determine fish presence, in particular above a gradient cascade thought to be a potential barrier to upstream fish migration.

33. Edeburn, A. and J. Wright. 2003. Line Creek Fisheries Investigation 2002 Program. Prepared for Elk Valley Coal Corporation - Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC.

- The overall intent of the fisheries investigations program was to provide a measure of the success of the environmental protection and management program at Line Creek mine with regard to aquatic resources. The scope of the assessment in 2002 focused on completing population estimates at 2 index sites in Line Creek and undertaking an assessment of bull trout spawning activity in 8.4 km of the Line Creek mainstem and 610

m of South Line Creek. Based on this study the authors proposed several recommendations for potential improvements to the program. These recommendations included the use of permanently marked index sites in the field, the use of consistent annual sampling methodologies, and scheduling sampling to minimize variability in flow conditions. Additional recommendations to improve the strength of the program included fish aging to complement previous data collected, adding another index site in reach 1, and completion of a Fish Habitat Assessment Procedure (FHAP) for each reach of Line Creek. Completion of a westslope cutthroat trout redd count and a benthic invertebrate survey to provide an indication of general aquatic health was also suggested.

- Reviewed in Teck 2011b as Interior Reforestation 2003.

34. Ferguson, K.D., and S.M. Leask. 1988. The Export of Nutrients from Surface Coal Mines. Environment Canada, West Vancouver, British Columbia. Regional Program Report 87-12. March.

- Historical nitrogen and phosphorus data were reviewed from 1980 to 1985 for the five (now Teck) surface coal mines in Elk Valley. The focus was evaluation of nitrogen losses from explosives use (range 0.1% to 4.3% among operations and years), with values being generally higher downstream of the mining activity.

35. Fording Coal Limited. 1985. Kilmarnock Creek Dragline Mining Proposal Environmental Impact Assessment. Report Prepared for British Columbia Mine Development Steering Committee. Prepared by Fording Coal Limited, 61 p.

- This report summarized the Kilmarnock Creek Dragline mining proposal (commencing 1987 for 2.5 years), its impacts, and the planned mitigation. The project involved the mining of a 1.1 km portion of the Kilmarnock Creek Valley, which was made possible by diversion of Kilmarnock Creek by a cut-off dam and large steel culverts skirting the perimeter of the pit. At the time of the project, the project area consisted of post logging condition with numerous access roads and drillsites. The authors concluded that the project would have a significant impact on groundwater, surface water quality, fisheries, and to a lesser degree wildlife in the Kilmarnock area. Impact management was planned to offset the disturbance. The mitigation included increased settling pond capacity in Brownie Creek, a cutoff dam above the mining area, large diameter steel culverts for diversion, stream re-establishment and the revised role of Kilmarnock settling ponds. There was also construction of offsite overwintering pools and summer rearing areas for westslope cutthroat trout. Creeks surveyed for physical habitat included Henretta Creek, Brownie Creek, Swift Creek, Lake Mountain Creek, Kilmarnock Creek and the Fording River.
- PDF missing sections two and three.

36. Fording Coal Limited. 1991. Fording Coal Limited Enhancement of the Upper Fording River (Above Confluence with Henretta Creek). Prepared by Fording Coal Limited, 21 p, 1 app.

- This report described the enhancements made to the Upper Fording River in a 400 to 450 m portion upstream from the confluence with Henretta Creek. It was enhanced to improve riffle-glide habitat. Enhancements included the installation of root wads, rock spurs, single rocks, rock groups, and rock clusters.
- PDF is missing appendices

37. Frenette, J.L. 2008. An Evaluation of Benthic Invertebrate Communities as an Indicator of Stream Ecosystem Health below Active Coal Mines in the Elk River Watershed. Master of Science Thesis, Royal Roads University. May.

- Water and benthic invertebrate samples were collected in September 2006 and 2007. Sampling areas were upstream and downstream of mining discharges on Michel Creek, and in reference streams (Alexander, Wheeler, Leach). In areas downstream of coal mines, several invertebrate community metrics were below, or at the low end, of the range observed at reference areas: #EPT individuals, # and % E and the number and percent of EPT taxa. Correlations were observed with some community metrics and concentrations of one or more water quality variables associated with mining (e.g., selenium, sulphate, nitrate, nitrite) as well as various habitat variables (e.g., elevation, water depth, discharge, substrate).
- Reviewed in Teck 2011b.

38. Golder Associates. 2015. Elkview Operations Baldy Ridge Extension Project. Annex J – Wildlife and Wildlife Habitat Baseline Report. Submitted to Teck Coal Limited, Elkview Operations. October.

- This reports described the terrestrial wildlife species and important wildlife habitats present near the Baldy Ridge Expansion Project at Elkview Operations that are potentially affected by the project. Particular attention was paid to searching for species of conservation concern, which were defined as those that were either federally listed as “Special Concern”, “Threatened” or “Endangered”, or provincially “blue” or “red-listed”. During the 11 field surveys that were conducted, 136 unique vertebrate species out of the 272 that were identified as potentially present were observed in the study area. Results were presented on area maps showing the locations of sightings of key species. The presence and absence of species of conservation concern were documented, however results were not presented in tabular form showing the precise locations of each of the species, so information could not be accurately interpreted with respect to site-specific presence/absence.

39. Hooton, R., Andrusak, H. and Bull, C.J. 1971. A Survey of the Elk River and Its Tributaries. British Columbia Fish and Wildlife Branch.

- Until shortly before the report was written, most local mining was underground, and there had been few obvious harmful effects to fish, wildlife and recreation. In the late 1960s, surface areas started to be disturbed for exploration and mining purposes as well as other mine related activities such as coal washing and coking, leading to periodic influences on Michel Creek. This study was undertaken to document the presence, distribution and relative abundance of sport fish and food fish populations in the Elk River Watershed, along with the watersheds use for fishing and other recreational activities. The effects of mining were also examined. Steams sampled included Boivin, Bingay, Forsyth, Quarrie, Aldridge, Bleasdel, Cadorna, Tobermory, Line and Todhunter Creeks. At least one-half mile of each stream was walked to make “cursory inventory”, including discharge, riparian vegetation, river velocity, river substrate and fish species that could be observed or angled.
- Reviewed in Teck 2011b and Golder 2014a.

40. Interior Reforestation Co. Ltd. 2006. Memo – 2005 Clode Ponds Fish Salvage Project. Prepared for Elk Valley Coal Corporation, Fording River Operations, Elkford, B.C. 6 p.

- Westslope cutthroat trout were captured via boat and backpack electrofishing, and seine netting from the Clode Ponds system in June 2005 to be released to predetermined release sites within the Fording River Watershed. Effort and capture results were recorded to determine total fish captured by reach, date and method. Biological endpoints (length and weight) were measured on a subset of fish, and abnormalities and deformities were recorded. Age-class was determined using length categories. Fish were released into Fish Pond Creek, the Fording River at Swift Creek, the Fording River at Chauncey Creek, and the Fording River upstream of the Fording Bridge. A total of 758 fish were captured in the salvage effort, 675 from boat electrofishing, 52 from backpack electrofishing and 31 from seining. Out of the total number of fish caught, 177 were subsampled and 2 of these fish had external deformities. The authors speculated that one of these deformities may have been due to selenium exposure (sideways spinal curvature and protruding eyeball). Overall, a low frequency of deformities were observed. Over the current salvage operation and another in 2004, 5,956 fish were removed from Clode Ponds. In light of a decreased catch per unit effort and the placement of a permanent upstream migration barrier between Clode Pond and the Fording River, the salvage was considered complete.
- Reviewed in Golder 2014a.

41. Interior Reforestation Co. Ltd. 2010. Elk River Watershed Historic Fish Distribution Project. Phase 1: Desktop Study, Prepared for Teck Coal, Prepared by Interior Reforestation

- In this report, species composition was assessed at a rudimentary level by contrasting mining influenced exposed sites with sites not located downstream of mines or reference sites of similar habitat to consider the effect of habitat preference. Habitat considered only lotic types in the Elk River, upstream of Elko. One hypothesis was that the observed species composition within the Elk River is a function of habitat preference. A second was that the migratory barrier at Elko has prevented colonization by some fish species. Finally, a third hypothesis suggested that the lack of certain species (e.g., Cottidae spp.) may be a result of mining activities, specifically increased water-borne selenium concentrations. This report presented the results of the desktop study that reviewed available data on fish distributions throughout the Elk River system since the 1970s. Whenever possible, the study used fisheries information previously compiled for Teck as part of past projects; the majority of which was fish distribution data obtained through the British Columbia Ministry of Environment's (MOE) Fisheries Inventory Summary System (FISS 2009). Pairs of tributaries were created with sub-basins that were similar in geographic location, drainage area and elevation, and all analyses were based on the assumption that a sub-basin pair represented similar stream habitat. The sub-basin pairs were Lower Elk River:Wigwam River, Michel Creek:Alexander Creek, and Upper Elk River:Fording River. The results of this study suggest that the Elk River (lower) sub-basin had the highest species diversity. This site was shown to be unique when averaged over the entire data record, but did not vary over time any differently than Wigwam sub-basin in a paired comparison. The differences between the exposed and reference watersheds were thought to be more likely due to habitat differences

rather than exposure, and suggests that the fish community structure and spatial variability are more likely due to the first two hypotheses.

- Reviewed in Golder 2015a.

42. KNRC (Kootenay Natural Resource Consulting. 2007. Fish Stream Identification, Greenhills Creek. Prepared for D. Charest, Elk Valley Coal Corporation. July 30, 2007.

- Between July 9th and 12th, 2007 Kootenay Natural Resource Consulting (KNRC) completed a fish stream identification (FSID) on Greenhills Creek to help facilitate the continued planning of Elk Valley Coal Corporation's Greenhills Operation mine development. The stream was surveyed from its confluence with the settling pond upstream approximately 5.8 km until the stream becomes covered by spill rock from a mine spoil. The survey consisted of electrofishing, visual observations, research and professional judgment. This information was used to classify the stream based on riparian assessment procedures. The lower portion of the creek was identified as fish-bearing (S3/S2), while the upper portion was not (S6). The west-fork of Greenhills Creek was also surveyed and determined to be fishbearing (S3/S4).

43. Lister, D.B. and Associates Ltd. and Kerr Wood Leidal Associates Ltd. 1980. Fording River Aquatic Environment Study. Prepared for Fording Coal Ltd., Elkford, B.C. 77 pp. + app.

- This report described the results and conclusions of a study of the Fording River and its Yellowstone cutthroat trout (now westslope cutthroat trout). The study area included the Fording River from the confluence with Henretta Creek downstream to a point 5 km downstream from the Kilmarnock Creek mouth, as well as several tributaries draining the mine area in the vicinity of the Clode Settling Pond. A major focus was a 0.9 km section of the Fording River which was diverted in 1977 to permit construction of the South Tailings Pond. The study consisted of a biophysical survey (habitat, obstructions), fish population estimates, fish size and age, benthic invertebrates, substrate sampling, temperature and discharge. Principal findings included decreased abundance of benthic invertebrates important for trout diets, especially in the diversion and other highly mine-influenced areas. Water quality was generally good in the areas studied, although suspended solids were significantly higher in mine-impacted areas particularly during spring freshet. Plans to reduce erosion and sediment output were recommended. Population estimates identified fish in all the sampling locations, but density was greater in areas not affected by mining, possibly due to habitat suitability. Several obstructions to fish migration were identified. A groundwater fed pond that was deepened and enlarged in November 1978 provided a significant overwintering habitat for fish. The tributaries around Clode Settling Pond lacked suitable cover and thus were utilized very little for spawning and rearing.

44. McDonald, L. Undated. Nutrient/Algal Growth Trends in the Elk River Basin. Impacts on Algal Growth of Various Nutrient Inputs, 1979-1983. Interim Report. Prepared for M.K. Baillargeon, Regional Waster Manager, Kootenay Region, B.C. Ministry of Environment.

- The study examined the relationship between nutrient levels and periphytic algal growth in the Elk River Basin due to concern over increased nitrate concentrations from bulk

explosive use and concentration of dissolved phosphorus originating from municipal sewage discharges. Significant increases in algal biomass were observed downstream of sewage discharges and were attributed to increased phosphorus as algal growth is phosphorus limited. Both nitrogen and phosphorus concentrations were very low in the unimpacted reaches of the Elk River, although phosphorus was still limiting. The high nitrate loads from surface mining produced a much stronger phosphorus limitation, and increased the maximum potential algal growth.

45. McDonald, L.E. 1983. Nutrient/Algal Growth Trends in the Elk River Basin interim Report. Impacts of Algal Growth of Various Nutrient Inputs 1979-1983. Prepared for M.K. Baillargeon, B.C. Ministry of Environment.

- This report examined the relationship between nutrient levels and periphytic algal growth in the Elk River Basin. The focus was on nuisance growth due to increased nitrate concentrations from bulk explosives used in the surface coal mining process along with increased phosphorus from municipal sewage discharges. Nutrients were monitored along the Elk River and at the mouth of major tributaries (Fording River and Line Creek, Michel Creek). Major increases in algal biomass were observed downstream of sewage discharges, suggesting algal growth was phosphorus limited. Increased nitrogen from coal mining produced a much greater phosphorus limitation, increasing the potential for algal growth downstream of phosphorus inputs.

46. McDonald, L. 1987. The Impact of Surface Coal Mining and Municipal Sewage Discharges on Nutrients and Algal Growth in the Elk River Basin 1981-1986. Prepared for J. McLaren, British Columbia Ministry of Environment and Parks. December.

- This study stated that nutrient concentrations in the Elk River upstream of Michel Creek were low prior to commencement of coal mining in early 1970s and algal production was and (at the time of the study continued to be) low based on limited phosphorus. However, soluble nitrogen levels in Elk River at Hosmer had increased 3-4 times since 1975 due to explosives use in coal mining. Lower Michel Creek was considered enriched by phosphorus loads from Leach and Wheeler Creeks (unknown origin) during freshet (April to July), resulting in heavy algal growth in Michel Creek and the Elk River downstream of (nitrogen sources) Erickson and Bodie Creeks. After freshet, when phosphorus loads from Leach and Wheeler were less, algal growth continued to be quite heavy downstream of Sparwood, likely due to phosphorus loadings from municipal sewage at Sparwood and Fernie.

47. McDonald, L. 2008. Coalbed Gas Baseline Water Quality Survey. Environmental Resource Information Project, Coalbed Gas Strategy. Prepared for Ministry of Energy, Mines, and Petroleum Resources. November.

- Sampling areas included Leach, Wheeler, Michel, Coal, Morrissey and North Lodgepole creeks of the Elk River watershed. Concentrations of TDS, component ions (e.g., calcium, magnesium, sulphate, etc.), nitrate, selenium, and other constituents were highest in Michel Creek compared to the other (non-mine-influenced) areas. Algal growth was considered "extensive" in most streams as a result of naturally occurring phosphorus, as well as soluble nitrogen input from bulk explosive use by mines. Triplicate benthic macroinvertebrate samples were taken on one occasion at each site using a 33 cm diameter Hess sampler. Michel Creek had the highest macroinvertebrate density of all sites in the five coal fields sampled.

48. McDonald, L.E., and M.M. Strosher. 1998. Selenium Mobilization from Surface Coal Mining in the Elk River Basin, British Columbia: A Survey of Water, Sediment, and Biota. Ministry of Environment, Lands and Parks, Cranbrook, British Columbia. September.

- Selenium concentrations in water, sediment, periphyton and fish were evaluated in the Elk River, the Fording River and Michel Creek. Water samples were also taken at the following tributaries: North Greenhills diversion, Swift Creek, Greenhills Creek, Eagle Settling Pond, Cataract Creek, Porter Creek, Harmer Creek, Line Creek, Bodie Creek, Erickson Creek, Spring Creek and Corbin Creek. The study concluded that significant quantities of selenium, mostly in the dissolved form, were being mobilized by surface coal mining into the Elk River System. At the time of the study there had been no known reports of selenium toxicity (embryo deformities or reproductive failure in fish as well as birds) in the Elk River Watershed, but no specific investigations had been undertaken to directly address this question. Areas downstream of mining had selenium concentrations in water up to 100 to 200 times those of reference areas, but only 2 to 5 times reference levels in bottom sediments, periphyton and invertebrates. Selenium concentrations in fish were also elevated in westslope cutthroat trout below coal mine tributary sources.

49. Morris, K.J., Cope, R.S. and Amos, L.P. 1997. Fish and Fish Habitat Inventory for Select Tributaries of the Upper Flathead and Elk Rivers. Interior Reforestation Co. Ltd. Prepared for Crestbrook Forest Industries Ltd. Cranbrook, B.C. September.

- The objective of this project was to undertake a reconnaissance level fish and fish habitat inventory to describe watershed wide fish distribution, habitat characteristics and provide stream and riparian classifications. In the Elk River watershed, Grave and South Line Creeks were sampled.

50. Nordin, R.N. 1982. The Effect on Water Quality of Explosives Use in Surface Mining. Volume 2: The Effect on Algal Growth. British Columbia Ministry of Environment, Victoria, B.C. File 64.0903. December.

- Nitrate concentrations were observed to increase in the Fording River from <0.1 mg/L upstream of Fording Mine to as much as 10 mg/L within the mine site at low flow. Moderate increases in biomass (based on weight and chlorophyll-a) and changes in community composition were observed downstream of mining. Lack of major increase in standing crop was attributed to the system being phosphorus limited.

51. North/South Consultants Inc. 2007. BP Canada Mist Mountain Project 2007 Baseline Aquatics Report: Surface Water Quality, Fisheries and Aquatic Habitat Assessment. Prepared for: Matrix Solutions Inc.

- BP Canada Energy Company (BP) has proposed a three- to five-year technical appraisal program to assess the feasibility of a coal bed gas (CBG) development in southeast British Columbia (BC). The Study Area is comprised of portions of the Flathead River, Michel Creek and Elk River watersheds. The overall objective of the aquatic baseline study was to characterize current conditions in representative watercourses in the Study Area. This report summarizes the aquatic program conducted to the end of 2007, which included water quality and fish and fish habitat surveys in August and September of

2007. Creeks included Flathead River, McEvoy Creek (Flathead River), Marten Creek, Fir Creek, Michel Creek, Robert Creek, Wheeler Creek (Michel Creek), Coal Creek, Morrissey Creek and North Lodgepole Creek (Elk River). Stream flows in August and September 2007 were typical or below average due to less than usual precipitation during the summer of 2007, and represented a baseflow or near-baseflow condition.

52. North/South Consultants Inc. 2008. BP Canada Mist Mountain Project 2008 Baseline Aquatics Report: Surface Water Quality, Fisheries and Aquatic Habitat Assessment. Prepared for: Matrix Solutions Inc.

- Second year (2008) of study #51. Surface water and sediment quality was considered protective of aquatic life at most of the sites, and suitable fish habitat was present at all sites in at least one of the open water seasons. Benthic invertebrates and periphyton were considered productive.
- Reviewed in Golder 2015a.

53. Oliver, G.G. 1999. Fish and Fish Habitat Assessment at Designated Sites in the Upper Fording River. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC. Prepared for Fording Coal Ltd. (Fording River Operations), Elkford, B.C. 47 pp. + app.

- An assessment of the enhancement structures in Fish Pond Creek and the Upper Fording River above Henretta Creek was completed in August 1998 to determine habitat suitability and functionality. Spawning surveys were also conducted in Fish Pond Creek, Henretta Creek, and Clode Pond outlet channels along with summer utilization surveys in lower Henretta Creek and the Fording River in June to August 1998. Additional information was collected on benthic invertebrates in Fish Pond Creek, water temperature at all sites. Overall, structures placed in Fish Pond Creek were more suitable than those in the Upper Fording River, with the highest densities of fish being at those structures with the greatest complexity. Overall the highest fish densities were in Fish Pond Creek.
- Reviewed in Golder 2014a

54. Pisces Environmental Consulting Services Ltd. 1992. Impact Assessment for fisheries for a proposed diversion of Line Creek. Prepared for Line Creek Resources Limited. Sparwood, BC. 19 pp.

- This report documents an impact assessment into the potential effects on fisheries resources of the diversion of a portion of Line Creek to accommodate several settling ponds. Previous studies on fish presence, habitat and habitat utilization were reviewed. Assessment concluded that the most sensitive fisheries component that might be affected by the diversion was the spawning, incubation and rearing life phases for bull trout and westslope cutthroat trout, however mitigation plans were sufficient to compensate for the loss.

55. Robinson, M., and J. Wright. 2005. Clode Pond Fish Salvage. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC.

- Westslope cutthroat trout were captured from the Clode Ponds system in 2004 with the goal of maximizing fish salvage and reducing the risks associated with the Turnbull

South Pit construction, spatially and temporally distribute fish to predetermined release sites, record effort and catch results by reach date and method, collect biological data (length, weight, age) on a subset of fish to compliment long term data sets, tag selected fish to observe movement, determine size at age, subsample for abnormalities and deformities and to block fish passage into Clode Ponds. Fish were collected by seining, boat electrofishing and backpack electrofishing. A total of 5,198 fish were collected, and of those 781 were subsampled for biological endpoints, and those fish that were greater than 190 mm (254) were tagged. Fish were relocated to the West Exfiltration, Fish Pond Creek, Fording River at Swift Creek, Fording River at Chauncey Creek, and the Fording River Bridge. Only 12 fish were found with external deformities that the authors concluded could be related to selenium exposure (fin damage most prevalent). The salvage operation was considered as success with minimal mortality observed.

- Reviewed in Golder 2014a.

56. Robinson, M., and J. Wright. 2005. Line Creek Fisheries Investigation 2004 Program. Prepared for Elk Valley Coal Corporation – Line Creek Operations, Sparwood, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, BC.

- This report documented the 2004 Fisheries Investigation Program through the Line Creek Operations. Fish densities were estimated and spawning assessments were undertaken in both Line Creek and in the lower section of South Line Creek. A slightly lower number of redds were observed compared to previous years, but the authors attributed that to increased seasonal discharge rather than reduction in population or escapement. The authors underscored the importance of careful identification and selection of environmental and biological indicators that provide measurable attributes if system behavior. This is essential in order to properly interpret fisheries changes that may be observed.

57. Robinson, M.D., Goodbrand, A., and R.J. MacDonald. 2014. Multi-Scale Characterization of Bull Trout Spawning Locations in Line Creek. Prepared for Teck Coal Ltd by Lotic Environmental Ltd. 25 pp.

- This study applied an approach that accounted for a range of spatial scales in assessing factors influencing bull trout spawning in Line Creek. The objective of this study was to characterize the factors that describe bull trout spawning habitat site selection at various spatial scales from the watershed to a sub-channel unit scale. The information will be used to inform Teck Coal's design and decisions regarding fish habitat offsetting projects. Results suggested that moving from the watershed scale to channel unit scale, the interaction between groundwater and surface water remains an important characterization of spawning habitat. Line Creek is a stream with a relatively high proportion of groundwater contributing to total streamflow, providing stable thermal conditions throughout the incubation period. There does not appear to be selection of spawning locations that rely upon groundwater discharge at smaller spatial scales (reach to channel unit). However, this study clearly demonstrates that spawning habitat site selection is likely driven by velocity and depth between channel units, and vertical hydraulic gradient within channel units.

58. Russell, E.J.A and G.G. Oliver. 1996. Fish-Stream Identification in the Upper Elk River Drainage. Prepared for: W.T. Westover, B.C. Ministry of Environment, Lands and Parks. Cranbrook, BC. January.

- This study investigated the presence/absence of fish, the identity of fish species present and reported on the specific habitat components of Alexander, Crossing, Chauncey, Ewin and Todhunter Creeks. Data were collected between August 22 and September 29, 1995. Four main criteria were identified for each watercourse: the number of stream reaches, the stream channel widths and gradients, the fish distribution throughout each stream length and the identification of species present. The streams were sorted into two broad categories: fish-bearing and non fish-bearing, with an additional category of suspect fish-bearing added to the report in the aftermath of extreme flooding and stream damage in June 2005. (NOTE Copy of paper received appears to be incomplete.)

59. Townsend Environmental. 2004. Fish Stream Identification of Grace Creek 2004. Prepared for Elk Valley Coal Corporation, Line Creek Operations, Sparwood, B.C.

- A fish stream identification assessment and report was completed for Grace Creek in May 2004. Westslope cutthroat trout were known to be present, and the study confirmed presence, but this limited to the lowest reach closest to the confluence with the Fording River. A culvert crossing associated with the rail line was deemed impassible to fish due to its considerable length (100 m). The authors recommended that further sampling be conducted in the fall.
- Reviewed in Teck 2011b.

60. Westslope Fisheries Ltd. 2003. Elk River Westslope Cutthroat trout radio telemetry study 2000-2002. Prepared for Columbia-Kootenay Fisheries Renewal Partnership. Cranbrook, B.C. 2003.

- Native species of cutthroat trout have experienced drastic declines in their distributions due to over-harvest, habitat degradation, and the introduction of non-native salmonids. In order to manage, protect and possibly enhance Elk River Westslope cutthroat trout in the face of increasing angler effort, industrial activities, and urban development, baseline data including fish distribution and habitat use are required. Despite geographic differences among spawning locations (mainstem margin, off-channel, ephemeral and perennial tributaries), there appeared to be little variation in both the microhabitats and characteristics of redds. Elk River cutthroat trout spawned in stable stream channels with a mean velocity of 0.62 m/s, an abundance of large woody debris (LWD), and undercut banks. Most redds were near the tailout of pools or glides.
- PDF missing appendices.

61. Wood, J.A. and R.J. Berdusco. 1999. Fording River revisited. A Review of Environmental Projects at Fording Coal Limited's Operations at Fording River over the Last 25 Years. Proceedings of the 23rd Annual British Columbia Mine Reclamation Symposium, Kamloops, B.C., 1999. Pages 58 – 70.

- At the time of this study, Fording Coal Ltd. had completed a number of environmental projects at the Fording River Operations including major river diversions, water quality control settling ponds, stream enhancements for westslope cutthroat trout, and wildlife habitat restoration. The study looked to assess the effectiveness of these projects after they had been in place for over 20 years. The projects included the Fording River Diversion (1976), the Henretta Dragline Project (1990), Fish Pond Creek and Upper

Fording River Habitat Replacement, Stormwater Settling Ponds, and Riparian Vegetation.

62. Wright, J. A., J.R. Berdusco and J.E. Bisset. 2005. Fish and Fish Habitat Monitoring in the Henretta Creek Reclaimed Channel (2004). Prepared for Elk Valley Coal Corporation – Fording River Operations, Elkford, BC. Prepared by Interior Reforestation Co. Ltd., Cranbrook, B.C. 24pp + 6 appendices.

- The 2004 fish and fish habitat monitoring represented Year 5 of the proposed seven year monitoring program required by the C-3 Reclamation Permit issued to Elk Valley Coal Corporation's Fording River Operations' (FRO). In 2004, changes in the invertebrate populations were variable between sites, dependent on the metric being analyzed. At the control site there were notable reductions in invertebrate densities, and moderate reductions in the percent composition of mayflies, stoneflies and caddisflies (%EPT) and mayflies alone (% Ephemeroptera). Upstream of Henretta Lake, benthic metrics were generally comparable with 2003 results. No spawning pairs were observed during the 2004 spawning and redd assessment. This is consistent with previous studies and observations of low utilization prior to and after mining. The detailed fish habitat assessments completed in 2004 indicate that the complexity and quantity of habitat and channel stability upstream and downstream is improving since 2003. Physical features in Henretta Lake have not changed for the most part. Continued channel adjustment (downcutting, narrowing, bar formation) which provides deeper pools and increased habitat features (glides, undercuts, woody debris accumulation), along with continued growth and establishment of healthy riparian vegetation, has further contributed to channel stability and complexity.

63. Wright, J., L. Amos and A. Edeburn. 2001. 2000 Fish and Fish Habitat Monitoring in the Henretta Creek Drainage. Report prepared for Fording Coal Ltd., Elkford, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, BC. -58 pp.

- As with previous reports, water temperature was higher downstream of Henretta Lake compared to upstream. Temperatures in the current report were considered within the preferred range for westslope cutthroat trout. The lack of spawning or redd construction during the 2000 survey provides no definitive proof of spawning or redd construction in the Henretta Creek reclaimed channel, despite the availability of suitable spawning gravels from initial restoration efforts by FCL. As the freshet flows coincide with the expected peak spawning times in the drainage, fish may have avoided use of the reclaimed channel as a natural mechanism to avoid the currently dynamic streambed and subsequently improve spawning success by relocating to other possible spawning locations. Average fish density in Henretta Creek during the 2000 survey was 6.7 fish/100m², which exceeds that of pre-impact studies. Fish habitat enhancement structures were installed within the stream portion of Henretta Creek to provide resting and holding cover for fry, juvenile and adult WCT, and most were meeting their intended purposes. Henretta Lake was still in the early stages of succession, with submergent, floating and emergent vegetation still lacking. At this stage, Henretta Lake also faces challenges with a lack of dissolved oxygen, particularly in December, January and February, confining fish to the top 2 m of the lake where water temperatures are well below optimum during these months.
- PDF missing several figures and the appendices.

REPORTS LACKING INFORMATION SPECIFIC TO TRIBUTARIES IN THE TEP

1. Amos, K., J. Wright, and L. Amos. 2000. Reconnaissance (1:20,000) fish and fish habitat inventory of the Upper Elk River Watershed WSCL 349-248100. Phases IV to VI. Prepared for Crestbrook Forest Industries Ltd. Cranbrook, B.C. Prepared by Interior Reforestation Co. Ltd. Cranbrook, B.C. May. 20 p. + 4 app.
2. Canton, S.P., A. Fairbrother, A.D. Lemly, H.M. Ohlendorf, L.E. McDonald, D.D. MacDonald. 2008. Experts workshop on the evaluation and management of selenium in the Elk Valley, British Columbia. Prepared for Ministry of Environment Environmental Protection Division. Nelson, B.C. August. 46 p. + 3 app.
3. COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2006. COSEWIC assessment and update status report on the Westslope Cutthroat trout *Oncorhynchus clarkii lewisi* (British Columbia population and Alberta population) in Canada. COSEWIC, Ottawa. Available: www.cosewic.gc.ca/. (December 2008).
4. Dessouki, T.C.E., and A. Ryan. 2010. Canada-British Columbia Water Quality Monitoring Agreement. Water Quality Assessment of the Kootenay, Elk, and St. Mary Rivers. Prepared for B.C. Ministry of Environment and Environment Canada. March.
5. Elphick, J.R., H.C. Bailey, B.K. Lo, J. Berdusco and G. Sword. 2009. Effect of Selenium on Early Life-stage Development of Westslope Cutthroat Trout. Report prepared for the Elk Valley Selenium Task Force. Prepared by Nautilus Environmental, Burnaby, British Columbia. -66 pp.
6. G3 Consulting Ltd. 2006. Coalbed gas environmental resource project. Fish population and habitat study review: Crowsnest Coalfield. Prepared for Ministry of Energy, Mines and Petroleum Resources, Victoria, B.C. January. 17 p + 2 app.
7. Hagen, J. and J.T.A. Baxter. 2009. 2008 Westslope Cutthroat trout population abundance monitoring of classified waters in the East Kootenay region of British Columbia. Prepared for BC Ministry of Environment, Cranbrook, BC, 34 p.
8. McDonald, L. 2009. Survey of Selenium in Water, Zooplankton and Fish in Lake Kootanusa, BC, 2008. Report Prepared for Environmental Protection, Kootenay Region, BC MOW on the Behalf of the Elk Valley Selenium Task Force, July 2009.
9. Robinson, M.D. 2011. Elk River fish distribution and longnose dace tissue assessment. Prepared for the Elk Valley Selenium Task Force. Prepared by Interior Reforestation Co. Ltd. 21 pp +app.
10. Swain, L.G. 2007. Canada-British Columbia Water Quality Monitoring Agreement. Water Quality Assessment of Elk River at Highway 93 near Elko (1968-2005). Prepared for B.C. Ministry of Environment and Environment Canada.
11. Swain, L.G. 2007. Canada-British Columbia Water Quality Monitoring Agreement. Water Quality Assessment of Elk River at Sparwood (2002-2005). Prepared for B.C. Ministry of Environment and Environment Canada.
12. Valdal, E.J. 2006. Cumulative effects of landscape disturbance on westslope cutthroat trout in the Upper Kootenay River Watershed: Implications for management and conservation. Master of Science Thesis. University of Calgary, Calgary, A.B. April.
13. Wilkinson, C.E. 2009. Sportfish population dynamics in an intensively managed river system. M.Sc. Thesis. University British Columbia, Vancouver, B.C. 140 p. + 3 app.